

# Particle Detectors

**Summer Student Lectures 2008**

**Werner Riegler, CERN, [werner.riegler@cern.ch](mailto:werner.riegler@cern.ch)**

- **History of Instrumentation ↔ History of Particle Physics**
- **The 'Real' World of Particles**
- **Interaction of Particles with Matter**
- **Tracking Detectors, Calorimeters, Particle Identification**
- **Detector Systems**




# The 'Real' World of Particles

## Elektro-Weak Lagrangian


$$\begin{aligned}
 L_{GSW} = & L_0 + L_H + \sum_l \left\{ \frac{g}{2} \bar{L}_l \gamma_\mu \bar{\nu} L_l \bar{A}^\mu + g' \left[ \bar{R}_l \gamma_\mu R_l + \frac{1}{2} \bar{L}_l \gamma_\mu L_l \right] B^\mu \right\} + \\
 & + \frac{g}{2} \sum_q \bar{L}_q \gamma_\mu \bar{\nu} L_q \bar{A}^\mu + \\
 & + g' \left\{ \frac{1}{6} \sum_q [\bar{L}_q \gamma_\mu L_q + 4 \bar{R}_q \gamma_\mu R_q] + \frac{1}{3} \sum_{q'} \bar{R}_{q'} \gamma_\mu R_{q'} \right\} B^\mu
 \end{aligned}$$

$$\begin{aligned}
 L_H = & \frac{1}{2} (\partial_\mu H)^2 - m_H^2 H^2 - h \lambda H^3 - \frac{h}{4} H^4 + \\
 & + \frac{g^2}{4} (W_\mu^+ W^\mu + \frac{1}{2 \cos^2 \theta_w} Z_\mu Z^\mu) (\lambda^2 + 2 \lambda H + H^2) + \\
 & + \sum_{l, q, q'} (\frac{m_l}{\lambda} \bar{l} l + \frac{m_q}{\lambda} \bar{q} q + \frac{m_{q'}}{\lambda} \bar{q}' q') H
 \end{aligned}$$

## Higgs Particle



$p p \rightarrow H^0$   
 $\quad \quad \quad \searrow \swarrow$   
 $\quad \quad \quad \gamma \gamma$




W. Riegler/CERN

# The 'Real' World of Particles

$$\begin{array}{l}
 1 \\
 0
 \end{array}
 \begin{pmatrix} e^- \\ \nu_e \end{pmatrix}
 \begin{pmatrix} \mu^- \\ \nu_\mu \end{pmatrix}
 \begin{pmatrix} \gamma^- \\ \nu_\gamma \end{pmatrix}
 \begin{array}{l}
 \text{Electromagnetic, Weak} \\
 \text{Weak}
 \end{array}$$

$$\begin{array}{l}
 \frac{2}{3} \\
 -\frac{1}{3}
 \end{array}
 \begin{pmatrix} u \\ d \end{pmatrix}
 \begin{matrix} 15-45 \text{ MeV} \\ 5-8.5 \text{ MeV} \end{matrix}
 \begin{pmatrix} c \\ s \end{pmatrix}
 \begin{matrix} 1-1.4 \text{ GeV} \\ 80-155 \text{ MeV} \end{matrix}
 \begin{pmatrix} t \\ b \end{pmatrix}
 \begin{matrix} 175 \text{ GeV} \\ 4-4.5 \text{ GeV} \end{matrix}
 \begin{array}{l}
 \text{Electromagnetic, Weak, Strong} \\
 \text{Electromagnetic, Weak, Strong}
 \end{array}$$

Spin  $\frac{1}{2}$  Particles


$p \sim uud$ , 
  
 $n \sim udd$ 
  
 $\pi \sim u\bar{d}, \bar{u}d, \frac{1}{\sqrt{2}}(u\bar{u} - d\bar{d})$ 
  
 $K \sim u\bar{s}, d\bar{s}, \bar{d}s, d\bar{s}$ 
  
 $\Lambda \sim uds$


Spin 1 Particles


EM:  $\gamma$  - Photon  $\uparrow$  QED
   
 Weak:  $W^\pm, Z^0$   $\downarrow$  Electroweak
   
 Strong:  $g$  - Gluon QCD


$$\begin{matrix}
 1 & \left( \underline{e} \right) & \left( \underline{\mu} \right) & \left( \underline{\gamma} \right) & \frac{2}{3} & \left( \underline{u} \right) & \left( \underline{c} \right) & \left( \underline{t} \right) \\
 0 & \left( \underline{\nu_e} \right) & \left( \underline{\nu_\mu} \right) & \left( \underline{\nu_\tau} \right) & -\frac{1}{3} & \left( \underline{d} \right) & \left( \underline{s} \right) & \left( \underline{b} \right)
 \end{matrix}$$


Electromagnetic Interaction  $\gamma$ -Photon

Scattering:   $e^- + e^- \rightarrow e^- + e^-$

Anihilation:   $e^+ + e^- \rightarrow \mu^+ + \mu^-$

Anihilation:   $e^+ + e^- \rightarrow B_0 + \bar{B}_0$


Bremsstrahlung:   $e^- + Atom \rightarrow e^- + \gamma + Atom$

Pair Production:   $\gamma + Atom \rightarrow e^+ + e^- + Atom$

$$\begin{matrix}
 1 \\
 0
 \end{matrix}
 \begin{pmatrix}
 \underline{e} \\
 \underline{\nu_e}
 \end{pmatrix}
 \begin{pmatrix}
 \underline{\mu} \\
 \underline{\nu_\mu}
 \end{pmatrix}
 \begin{pmatrix}
 \underline{\tau} \\
 \underline{\nu_\tau}
 \end{pmatrix}
 \frac{2}{3}
 \begin{pmatrix}
 \underline{u} \\
 \underline{d}
 \end{pmatrix}
 \begin{pmatrix}
 \underline{c} \\
 \underline{s}
 \end{pmatrix}
 \begin{pmatrix}
 \underline{t} \\
 \underline{b}
 \end{pmatrix}$$


Weak Interaction  $W^\pm, Z^0$

Neutral Current:




$$e^- + \bar{\nu}_n \rightarrow e^- + \bar{\nu}_n$$

Muon Decay:




$$\mu^- \rightarrow \nu_\mu + e^- + \bar{\nu}_e$$

Neutron Decay:



$$n \rightarrow p + e^- + \bar{\nu}_e$$

H Production:



1	$\begin{pmatrix} e \\ \nu_e \end{pmatrix}$	$\begin{pmatrix} \mu \\ \nu_\mu \end{pmatrix}$	$\begin{pmatrix} \tau \\ \nu_\tau \end{pmatrix}$	$\frac{2}{3}$	$\begin{pmatrix} u \\ d \end{pmatrix}$	$\begin{pmatrix} c \\ s \end{pmatrix}$	$\begin{pmatrix} t \\ b \end{pmatrix}$
0				$-\frac{1}{3}$			

Strong Interaction

$g$  Gluons




Proton



Self Interaction





"Confinement"



.... Strong Interaction.....

$e^+ + e^- \rightarrow$  jets in Detector



e.g. Two jets of Hadrons are 'spraying' away from the Interaction Point.

Over the last century  
this 'Standard Model' of  
Fundamental Physics was discovered  
by studying

Radioactivity

Cosmic Rays

Particle Collisions (Accelerators)

A large variety of Detectors and  
experimental techniques have been  
developed during this time.

"Material Culture of Particle Physics"



## Scales

$$E = ma^2$$

$$E = mb^2$$

$$E = mc^2 \leftarrow \text{Energy} \hat{=} \text{Mass}$$

⋮


$$m(\text{electron}) = 9.1 \cdot 10^{-31} \text{ kg}$$

$$m_e c^2 = 8.19 \cdot 10^{-14} \text{ J}$$

$$= 510\,999 \text{ Electron Volt (eV)}$$


$$= 0.511 \text{ MeV}$$

$$1 \text{ Electron Volt} = e_0 \cdot 1V = 1.603 \cdot 10^{-19} \text{ J}$$



1 Electron Volt = Energy an Electron gains as it traverses a Potential Difference of 1V

# Build your own Accelerator



## Scales

Visible Light:

$$\lambda = 500 \text{ nm}, \quad h\nu \sim 2.5 \text{ eV}$$

Excited States in Atoms:

$$1 - 100 \text{ keV} \quad \text{"X-Rays"}$$

Nuclear Physics:

$$1 - 50 \text{ MeV}$$

E.g:  ${}_{39}^{90}\text{Y} \rightarrow \beta^- \rightarrow e^-$  with  $E_a = 2.283 \text{ MeV}$

$$E_k = mc^2(\gamma - 1) \quad mc^2 \sim 0.511 \text{ MeV}$$

$$\gamma = \frac{E_k}{mc^2} + 1 \sim 5.5$$

$$\beta = \frac{v}{c} = \sqrt{1 - \left(\frac{mc^2}{E_k + mc^2}\right)^2} \sim 0.98 \rightarrow \text{Highly Relativistic}$$

$$E_{\text{kin}} = mc^2 \rightarrow mc^2(\gamma - 1) = mc^2 \rightarrow \gamma = 2 \rightarrow \beta = 0.87$$

E.g:  ${}_{95}^{241}\text{Am} \rightarrow \alpha$  with  $E_{\text{kin}} = 5.486 \text{ MeV}$ ,  $m_{\alpha}c^2 = 3.75 \text{ GeV}$

$$\gamma \sim 1.0015 \quad \beta \sim 0.054 \rightarrow 16.2 \cdot 10^6 \text{ m/s}$$

Particle Physics:

$$1 - 1000 \text{ GeV} \quad (\text{LHC } 14 \text{ TeV})$$

Highest Measured Energy:

$$10^{20} \text{ eV} \quad (\text{Cosmic Rays})$$

# Basics

9

## Lorentz Boost:

$$\mu^- \rightarrow e^- + \bar{\nu}_e + \gamma_\mu \quad \tau = 2.2 \cdot 10^{-6} \text{ s}$$

E.g. Produced by Cosmic Rays (p, He, Li ...) colliding with air in the upper atmosphere  $\sim 10 \text{ km}$

$$s = v \cdot \tau \sim c \cdot \tau = 660 \text{ m}$$

But we see Muons here on Earth

$$E_\mu \sim 2 \text{ GeV}, m_\mu c^2 = 105 \text{ MeV} \rightarrow \gamma \sim 19$$

$$\text{Relativity: } \bar{\tau} = \tau \cdot \gamma$$

$$s = c \cdot \bar{\tau} = 12.5 \text{ km} \rightarrow \text{Earth}$$


$$\text{Pions: } \pi^+, \pi^- \quad \tau \sim 2.6 \cdot 10^{-8} \text{ s}, m_\pi c^2 = 135 \text{ MeV}$$

$$2 \text{ GeV} \rightarrow s = 115 \text{ m}$$

Pions were discovered in Emulsions exposed to Cosmic Rays on high Mountaintops.

## Basics


E.g.  $\pi^- (ud) \rightarrow \mu^- + \bar{\nu}_\mu \quad (> 99.9\%)$



$$\tau = 2.6 \cdot 10^{-8} \text{ s}$$

$\pi^-$

$$\vec{p} = 0, E = m_\pi c^2$$



$$\vec{p}_1 + \vec{p}_2 = 0, E_\mu + E_\nu = E$$

$$0 = \frac{m_\mu v_1}{\sqrt{1 - \frac{v_1^2}{c^2}}} + \frac{m_\nu v_2}{\sqrt{1 - \frac{v_2^2}{c^2}}} \quad \left. \vphantom{\frac{m_\mu v_1}{\sqrt{1 - \frac{v_1^2}{c^2}}}} \right\} v_1, v_2$$

$$m_\pi c^2 = \frac{m_\mu c^2}{\sqrt{1 - \frac{v_1^2}{c^2}}} + \frac{m_\nu c^2}{\sqrt{1 - \frac{v_2^2}{c^2}}}$$

$E_\mu, E_\nu$  are uniquely defined


→ Two Body Decay gives "sharp"

Energies of the Decay Particles

## Basics


11

1920ies:  $\beta^-$  Radioactivity




But:  $e^-$  shows a continuous Energy Spectrum

→ W. Pauli proposed an "invisible" Particle →  $\bar{\nu}_e$



For  $> 2$  Body decay, the Energy Spectrum of the decay particles depends on the Nature of the Interaction. Kinematics alone doesn't define the Energies.

# Stopping Pions and measuring the decay electron Spectrum:



Energy Spectrum (3 Body Decay)




↓  
"sharp" Energy (2 Body Decay)

## Basics

Invariant Mass:

LAB:



Relativity:  $\tilde{a} = \begin{pmatrix} a_0 \\ \vec{a} \end{pmatrix}$   $\tilde{b} = \begin{pmatrix} b_0 \\ \vec{b} \end{pmatrix}$   $\tilde{a} \tilde{b} = a_0 b_0 - \vec{a} \cdot \vec{b}$

$$E = mc^2 \gamma, \quad \vec{p} = m \vec{v} \gamma$$

$$\tilde{p} = \begin{pmatrix} \frac{E}{c} \\ \vec{p} \end{pmatrix}, \quad \tilde{p}_1 = \begin{pmatrix} \frac{E_1}{c} \\ \vec{p}_1 \end{pmatrix}, \quad \tilde{p}_2 = \begin{pmatrix} \frac{E_2}{c} \\ \vec{p}_2 \end{pmatrix}$$

$$\tilde{p} = \tilde{p}_1 + \tilde{p}_2 \quad \text{Energy + Momentum Conservation}$$

$$\tilde{p}^2 = (\tilde{p}_1 + \tilde{p}_2)^2 \rightarrow \tilde{p} \tilde{p} = \tilde{p}_1 \tilde{p}_1 + \tilde{p}_2 \tilde{p}_2 + 2 \tilde{p}_1 \tilde{p}_2$$

$$\underline{M^2 c^2 = m_1^2 c^2 + m_2^2 c^2 + 2 \left( \frac{E_1 E_2}{c^2} - p_1 p_2 \cos \theta \right)}$$


- Measuring Momenta and Energies OR
- Measuring Momenta and identifying Particles gives the Mass of the original Particle



## Basics

14

E.g: Discovery of  $V^0$  Particles






"If 1 is a Proton and 2 is a Pion  
the Mass of the  $V^0$  particle is ...."

Identification is the Experiment by  
looking at the specific Ionization .....  
(see later)

# $\mu$ -Lifetime

The muon (any unstable Particle) doesn't have an inner 'clock', i.e. nothing that tells it' age.



We look for a Distribution  $P(t)$  where drawing a time  $t$  from  $P(t)$  and subtracting a random Number  $T$  gives again the same Distribution  $P(t)$  for  $t > 0$

$p(T)$ : Arbitrary Distribution

$$P_2(t) = \int_0^{\infty} p(\tau) P_1(t-\tau) d\tau$$

$$P_2(t) := c_1 P_1(t)$$

only if  $P_1(t-T) = P_1(t) \cdot P_1(-T)$

$\rightarrow$   $P(t) = c_1 e^{-c_1 t}$   $\rightarrow$  Exponential Distribution

$$\gamma = \int_0^{\infty} t c_1 e^{-c_1 t} dt = \frac{1}{c_1} \quad \text{Average Lifetime}$$


$$P(t) = \frac{1}{\gamma} e^{-\frac{t}{\gamma}} \quad \gamma = \text{"Life time"}$$

"A Particle has a lifetime  $\gamma$ " means:

The Probability that it Decays at time  $t$  after starting to measure it (independent of what happened before) is  $P(t) = \frac{1}{\gamma} e^{-\frac{t}{\gamma}}$


Electron e  $m_e = 0.511 \text{ MeV}$   $\gamma = \infty$

Myon  $\mu$   $m_\mu = 105.7 \text{ MeV}$   $\gamma = 2.2 \cdot 10^{-6} \text{ s}$ ,  $c\gamma = 659 \text{ m}$




$$\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu \quad (> 99.9\%)$$

Tauon  $\tau$   $m_\tau = 1777 \text{ MeV}$   $\gamma = 2.9 \cdot 10^{-13} \text{ s}$ ,  $c\gamma = 87 \mu\text{m}$



$$\tau^- \rightarrow \mu^- + \bar{\nu}_\mu + \nu_\tau \quad (\sim 17\%)$$



$$\tau^- \rightarrow e^- + \bar{\nu}_e + \nu_\tau \quad (\sim 17\%)$$



$$\tau^- \rightarrow \pi^- + \nu_\tau \quad (\sim 11\%)$$



$$\tau^- \rightarrow \pi^0 + \pi^- + \nu_\tau \quad (\sim 25\%)$$

⋮

Due to the larger Mass, more Decay Possibilities are open to the  $\tau$

→ the Lifetime is smaller ....

<http://pdg.lbl.gov>

~ 180 Selected Particles

$\eta, W^\pm, Z^0, g, e, \mu, \tau, \nu_e, \nu_\mu, \nu_\tau, \pi^\pm, \pi^0, \eta, f_0(600), g(700),$   
 $\omega(782), \eta'(958), f_0(980), a_0(980), \phi(1020), h_1(1170), b_1(1235),$   
 $a_1(1260), f_2(1270), f_1(1285), \eta(1295), \pi(1300), a_2(1320),$   
 $f_0(1370), f_1(1420), \omega(1420), \eta(1440), a_0(1450), g(1450),$   
 $f_0(1500), f_2'(1525), \omega(1650), \omega_3(1670), \pi_2(1670), \phi(1680),$   
 $g_3(1690), g(1700), f_0(1710), \pi(1800), \phi_3(1850), f_2(2010),$   
 $a_4(2040), f_4(2050), f_2(2300), f_2(2340), K^\pm, K^0, K_S^0, K_L^0, K^*(892),$   
 $K_1(1270), K_1(1400), K^*(1410), K_0^*(1430), K_2^*(1430), K^*(1680),$   
 $K_2(1770), K_3^*(1780), K_2(1820), K_4^*(2045), D^\pm, D^0, D^*(2007)^0,$   
 $D^*(2010)^\pm, D_1(2420)^0, D_2^*(2460)^0, D_2^*(2460)^\pm, D_s^\pm, D_s^{*\pm},$   
 $D_{s1}(2536)^\pm, D_{s1}(2573)^\pm, B^\pm, B^0, B^*, B_s^0, B_c^\pm, \eta_c(1S), J/\psi(1S),$   
 $\chi_{c0}(1P), \chi_{c1}(1P), \chi_{c2}(1P), \psi(2S), \psi(3770), \psi(4040), \psi(4160),$   
 $\psi(4415), \Upsilon(1S), \chi_{b0}(1P), \chi_{b1}(1P), \chi_{b2}(1P), \Upsilon(2S), \chi_{b0}(2P),$   
 $\chi_{b2}(2P), \Upsilon(3S), \Upsilon(4S), \Upsilon(10860), \Upsilon(11020), p, n, N(1440),$   
 $N(1520), N(1535), N(1650), N(1675), N(1680), N(1700), N(1710),$   
 $N(1720), N(2190), N(2220), N(2250), N(2600), \Delta(1232), \Delta(1600),$   
 $\Delta(1620), \Delta(1700), \Delta(1905), \Delta(1910), \Delta(1920), \Delta(1930), \Delta(1950),$   
 $\Delta(2420), \Lambda, \Lambda(1405), \Lambda(1520), \Lambda(1600), \Lambda(1670), \Lambda(1690),$   
 $\Lambda(1800), \Lambda(1810), \Lambda(1820), \Lambda(1830), \Lambda(1890), \Lambda(2100),$   
 $\Lambda(2110), \Lambda(2350), \Sigma^+, \Sigma^0, \Sigma^-, \Sigma(1385), \Sigma(1660), \Sigma(1670),$   
 $\Sigma(1750), \Sigma(1775), \Sigma(1915), \Sigma(1940), \Sigma(2030), \Sigma(2250), \Xi^0, \Xi^-,$   
 $\Xi(1530), \Xi(1690), \Xi(1820), \Xi(1950), \Xi(2030), \Omega^-, \Omega(2250)^-,$   
 $\Lambda_c^+, \Lambda_c^0, \Sigma_c(2455), \Sigma_c(2520), \Xi_c^+, \Xi_c^0, \Xi_c'^+, \Xi_c'^0, \Xi(2645),$   
 $\Xi_c(2780), \Xi_c(2815), \Omega_c^0, \Lambda_b^0, \Xi_b^0, \Xi_b^-, t\bar{t}$

There are many more

Particle	Mass (meV)	Life time $\tau$ (s)	$c\tau$
$\gamma$	0	$\infty$	$\infty$
$\pi^\pm (u\bar{d}, d\bar{u})$	140	$2.6 \cdot 10^{-8}$	7.8 m
$K^\pm (u\bar{s}, \bar{u}s)$	494	$1.2 \cdot 10^{-8}$	3.7 m
$K^0 (d\bar{s}, \bar{d}s)$	497	$5.1 \cdot 10^{-8}$ $8.9 \cdot 10^{-11}$	15.5 m 2.7 cm
$D^\pm (c\bar{d}, \bar{c}d)$	1869	$1.0 \cdot 10^{-12}$	315 $\mu\text{m}$
$D^0 (c\bar{u}, \bar{c}u)$	1864	$4.1 \cdot 10^{-13}$	123 $\mu\text{m}$
$D_s^\pm (c\bar{s}, \bar{c}s)$	1969	$4.9 \cdot 10^{-13}$	147 $\mu\text{m}$
$B^\pm (u\bar{b}, \bar{u}b)$	5279	$1.7 \cdot 10^{-12}$	502 $\mu\text{m}$
$B^0 (b\bar{d}, \bar{b}d)$	5279	$1.5 \cdot 10^{-12}$	462 $\mu\text{m}$
$B_s^0 (s\bar{b}, \bar{s}b)$	5370	$1.5 \cdot 10^{-12}$	438 $\mu\text{m}$
$B_c^\pm (c\bar{b}, \bar{c}b)$	$\sim 6400$	$\sim 5 \cdot 10^{-13}$	150 $\mu\text{m}$
$p (uud)$	938.3	$> 10^{33} \text{ y}$	$\infty$
$n (udd)$	939.6	885.7 s	$2.655 \cdot 10^8 \text{ km}$
$\Lambda^0 (uds)$	1115.7	$2.6 \cdot 10^{-10}$	7.89 cm
$\Sigma^+ (uus)$	1189.4	$8.0 \cdot 10^{-11}$	2.404 cm
$\Sigma^- (dds)$	1197.4	$1.5 \cdot 10^{-10}$	4.434 cm
$\Xi^0 (uss)$	1315	$2.9 \cdot 10^{-10}$	8.71 cm
$\Xi^- (dss)$	1321	$1.6 \cdot 10^{-10}$	4.91 cm
$\Omega^- (sss)$	1672	$8.2 \cdot 10^{-11}$	2.461 cm
$\Lambda_c^+ (udc)$	2285	$\sim 2 \cdot 10^{-13}$	60 $\mu\text{m}$
$\Xi_c^+ (usc)$	2466	$4.4 \cdot 10^{-13}$	132 $\mu\text{m}$
$\Xi_c^0 (dcs)$	2472	$\sim 1 \cdot 10^{-13}$	29 $\mu\text{m}$
$\Omega_c^0 (ssc)$	2698	$6.0 \cdot 10^{-14}$	19 $\mu\text{m}$
$\Lambda_b (uab)$	5620	$1.2 \cdot 10^{-12}$	368 $\mu\text{m}$

"Secondary Vertices"

From the 'hundreds' of Particles listed by the PDG there are only  $\sim 27$  with a life time  $c\tau > \sim 1\mu\text{m}$  i.e. they can be seen as 'tracks' in a Detector.

$\sim 13$  of the 27 have  $c\tau < 500\mu\text{m}$  i.e. only  $\sim\text{mm}$  range at GeV Energies.  
 $\rightarrow$  "short" tracks measured with Emulsions or Vertex Detectors.

From the  $\sim 14$  remaining particles

$$e^{\pm}, \mu^{\pm}, \gamma, \pi^{\pm}, K^{\pm}, K^0, p^{\pm}, n$$

are by far the most frequent ones

A particle Detector must be able to identify and measure Energy and Momenta of these 8 particles.


$e^\pm$	$m_e = 0.511 \text{ MeV}$	} EM
$\mu^\pm$	$m_\mu = 105.7 \text{ MeV} \sim 200 m_e$	
$\gamma$	$m_\gamma = 0, Q = 0$	
$\pi^\pm$	$m_\pi = 139.6 \text{ MeV} \sim 270 m_e$	} EM, Strong $\sim 3.5 m_\pi$
$K^\pm$	$m_K = 493.7 \text{ MeV} \sim 1000 m_e$	
$p^\pm$	$m_p = 938.3 \text{ MeV} \sim 2000 m_e$	
$K^0$	$m_{K^0} = 497.7 \text{ MeV} \quad Q=0$	} Strong
$n$	$m_n = 939.6 \text{ MeV} \quad Q=0$	

The Difference in *Mass, Charge*


*Mass, Charge, Interaction*

is the key to the Identification







- Electrons ionize and show Bremsstrahlung due to the small mass
- Photons don't ionize but show Pair Production in high  $Z$  Material. From then on equal to  $e^\pm$
- Charged Hadrons ionize and show Hadron Shower in dense Material.
- Neutral Hadrons don't ionize and show Hadron Shower in dense Material
- Myons ionize and don't shower




**ATLAS**

**CMS**




**LHCb**


**ALICE**



W. Riegler/CERN

Vertex Detector  
Inner Tracking Chamber  
Time Projection Chamber  
Electromagnetic Calorimeter  
Hadron Calorimeter  
Muon Detectors














-  Vertex Detector
-  Inner Track Chamber
-  Time Projection Chamber
-  Electromagnetic Calorimeter
-  Superconducting Magnet Coil
-  Hadron Calorimeter
-  Muon Detection Chambers
-  Luminosity Monitors



Fig. 1 - The ALEPH Detector

$\gamma, e^+, \tau^+, k^+$   
 $k^0, p, n, \mu^+$

# ALEPH




# CMS A Compact Solenoidal Detector for LHC




$$Z \rightarrow e^+ e^-$$

Two high momentum charged particles depositing energy  
in the Electro Magnetic Calorimeter



$$Z \rightarrow \mu^+ \mu^-$$


Two high momentum charged particles traversing all calorimeters and leaving a signal in the muon chambers.






$$Z \rightarrow t^+ t^- \rightarrow m^+ n^- e^- n$$

1 or 2 secondary vertices, high momentum electron,  
high momentum muon, missing momentum.




$$Z \rightarrow q \bar{q}$$

Two jets of particles




$$Z \rightarrow q \bar{q} g$$

Three jets of particles





Two secondary vertices with characteristic decay particles giving invariant masses of known particles.

Bubble chamber like – a single event tells what is happening. Negligible background.




## ALEPH Higgs Candidate





**Undistinguishable background exists. Only statistical excess gives signature.**

# Cosmic Shower of Muons



# Higgs Boson at CMS



Particle seen as an excess of two photon events above the irreducible background.


## Conclusion:

Only a few of the numerous known particles have lifetimes that are long enough to leave tracks in a detector.

Most of the particles are measured through the decay products and their kinematic relations (invariant mass). Most particles are only seen as an excess over an irreducible background.

Some short lived particles (b,c –particles) reach lifetimes in the laboratory system that are sufficient to leave short tracks before decaying → identification by measurement of short tracks.

In addition to this, detectors are built to measure the 8 particles



$e^{\pm}, \mu^{\pm}, \gamma, \pi^{\pm}, K^{\pm}, K^0, p^{\pm}, n$

Their difference in mass, charge and interaction is the key to their identification.