



Off-shell Higgs for LHC Run 2


- experimental view and short/mid term wish-list

Yanyan Gao (University of Edinburgh)
Jian Wang (Universite Libre de Bruxelles)


LHC HXSWG "off-shell Higgs" open meeting October 24, 2014

What has been achieved

- Phenomenology ideas
 - Kauer, Passarino JHEP 1208 (2012) 116
 - · Caola, Melnikov PRD 88 (2013) 054024; Campbell et al, JHEP 1404 (2014) 060
- CMS/ATLAS results on Higgs width constraints using ZZ->4l and ZZ->2l2nu channels
 (PLB 736 (2014) 64, ATLAS-CONF-2014-042)



 $\Gamma_{H}/\Gamma_{H}^{SM} < 5.4 @ 95\% CL$



 $\Gamma_{\text{H}}/\Gamma_{\text{H}}^{\text{SM}} < 4.8{\sim}7.7 \ @ \ 95\% \ \text{CL}$ (depending on gg->VV background K factor)

To sharpen the Higgs width constraints

- Analyses will surely be repeated in Run 2
 - H->ZZ->4l channel is statistical limited
- Constraints on the Higgs width with VBF only
 - No loop contribution model independent (?)
 - 20/fb @ 8TeV, VBF ZZ->2l2nu channel sets a limit of ~40*SM Higgs width (CMS-PAS-HIG-14-002); VBF WW could be competitive here
 - Better S/B; theoretical uncertainties better understood (e.g. background K factor); but still ggH contamination
 - Much larger statistics in Run 2; could combine all ZZ,WW channels

gg->VV @ NLO and jet binning

- K factor of gg->VV background and consequently interference unknown
 - Currently using signal K factor with residual uncertainties (Passarino, EPJC 74 (2014) 2866, Bonvini et al, PRD 88 (2013) 3, 034032)
 - Prospect of approximation and uncertainties
 - What do we do for 13 TeV
- Reliable description of jet multiplicity/pT(VV)
 - Jet binning/veto unavoidable in H->WW analysis, in order to suppress top backgrounds
 - ggH+2jet contamination in VBF channel
 - Suggestion about uncertainty estimation?

To constrain Higgs properties in EFT

- New ideas using off-shell production?
 - Yes, they are popping up, e.g. 1406.6338, etc
 - Status of MC tools to facilitate analyses
 - Dedicated talks today
- What is feasible/ should be prioritized in the Run 2
- In long run, what is the additional/complementary physics reach compared to onshell measurements?