

## Estimation of the performance of a HERMES-type gas target internal to the LHC

*Tuesday 15 September 2015 09:30 (30 minutes)*

A storage cell target is capable of producing a high areal density at minimum gas flow into a vacuum system, e.g. that of the LHC. It may be fed with polarized hydrogen atoms for the study of single-spin asymmetries in ultra-relativistic fixed-target pp collisions, or similar light-ion reactions with spin. Another application could be to inject heavy noble gases like Xe ( $M \approx 131$ ) in conjunction with a Pb beam to investigate Heavy Ion collisions. In a recent study [1], the performance of a storage cell target combined with LHC beams has been estimated under realistic assumptions. The luminosities obtained are for pp collisions of the order  $10^{33}/\text{cm}^2\text{s}$ , and for HI collisions of the order of the Pb-Pb collider design value of  $10^{27}/\text{cm}^2\text{s}$ . The assumptions and results will be discussed in the talk.

[1] C. Barschel, P. Lenisa, A. Nass, and E. Steffens, *Advances in High Energy Physics - special AFTER@LHC issue*, Article ID 463141. Accepted for publ. July 5, 2015.

**Primary author:** Dr STEFFENS, Erhard (Univ. of Erlangen-Nürnberg)

**Presenter:** Dr STEFFENS, Erhard (Univ. of Erlangen-Nürnberg)

**Session Classification:** Session 4