



Contribution ID: 857

Type: **Oral contribution**

Study of the Solar Modulation of Local Interstellar Protons with AMS-02, PAMELA, Neutron Monitors and Voyager 1

Saturday, 1 August 2015 11:30 (15 minutes)

Galactic Cosmic Rays (GCRs) entering the heliosphere are disturbed by the magnetic field of the Sun, which varies with a period of 11 years. The solar modulation affects the GCR fluxes up to few tens of GeV, modifying the shape and the intensity of the local interstellar spectrum (LIS). The time variation of the galactic cosmic protons at Earth can be studied indirectly on ground with the Neutron Monitors (NMs) and directly from space with AMS-02 (2011-now) and PAMELA (2006-2010).

A new parametrization of the LIS will be presented, based on the latest data from AMS-02 and Voyager 1. Using the framework of the force-field approximation, the solar modulation parameter will be extracted from the time-dependent proton fluxes measured by PAMELA. The results will be compared with the modulation parameter inferred by NMs.

Collaboration

– not specified –

Registration number following "ICRC2015-I/"

737

Primary author: Mr CORTI, Claudio (University of Hawai'i at Manoa (US))

Co-authors: CONSOLANDI, Cristina (University of Hawai'i at Manoa (US)); WHITMAN, Kathryn (University of Hawai'i at Manoa (US)); BINDI, Veronica (University of Hawai'i at Manoa (US))

Presenter: Mr CORTI, Claudio (University of Hawai'i at Manoa (US))

Session Classification: Parallel SH 03 SEP II

Track Classification: SH-TH