



Contribution ID: 654

Type: **Oral contribution**

The origin of the ankle in the UHECR spectrum, and of the extragalactic protons below it

Tuesday 4 August 2015 15:45 (15 minutes)

The sharp change in slope of the ultra-high energy cosmic ray spectrum around $10^{18.6}$ eV (the ankle), combined with evidence of a light but extragalactic component near and below the ankle and intermediate composition above, has proved exceedingly challenging to understand theoretically.

We propose a mechanism whereby photo-disintegration of ultra-high energy nuclei in the region surrounding a UHECR accelerator naturally accounts for the observed spectrum and inferred composition at Earth. We discuss the conditions required to reproduce the spectrum above $10^{17.5}$ eV and the composition, which – in our model – consists below the ankle of extragalactic protons and the high energy tail of Galactic Cosmic Rays, and above the ankle of surviving nuclei from the extended source. Predictions for the spectrum and flavors of neutrinos resulting from this process will be presented, and also implications for candidate sources.

Collaboration

– not specified –

Registration number following "ICRC2015-I/"

388

Author: Prof. FARRAR, Glennys (New York University)

Co-authors: Prof. ANCHORDOQUI, Luis (Lehman College and CUNY); Dr UNGER, Michael (NYU and KIT)

Presenter: Prof. FARRAR, Glennys (New York University)

Session Classification: Parallel CR17 EAS spec

Track Classification: CR-TH