



Contribution ID: 1020

Type: Oral contribution

Time-dependent modelling of particle acceleration and non-thermal emission in Eta Carina

Saturday 1 August 2015 12:15 (15 minutes)

Eta Carina is the only colliding-wind binary for which non-thermal emission is detected from hard X-rays to high-energy gamma rays. Although the physical conditions in the shock region change on timescales of hours to days, the variability seen at GeV energies is quite weak and on significantly longer timescales. The gamma-ray spectrum exhibits two spectral features that can be interpreted as emission originating from the shock and post-shock regions of the two distinct stellar winds. Here we report on the first time-dependent modelling of the non-thermal emission in Eta Carina in this scenario, which aims to explain the level of gamma-ray emission as well as the variability with orbital phase. We used a time-dependent acceleration scheme to derive the spectrum of relativistic particles in the adiabatic shock of the companion star, including the effects of cosmic ray pressure modification. We find that emission from primary electrons is unlikely to be responsible for the gamma-ray emission, and that gamma-ray emission from accelerated protons interacting with the dense shocked wind material can explain the observations. The significant pair production absorption in the stellar radiation fields leads to a drop in the predicted gamma-ray emission above several hundred GeV - consistent with the upper limits as reported by H.E.S.S.

Collaboration

– not specified –

Registration number following "ICRC2015-I/"

571

Primary authors: Dr PARKIN, Elliot Ross (University of Leeds); HINTON, Jim (University of Leicester); OHM, Stefan (DESY); ZABALZA, Victor (University of Leicester)

Presenter: ZABALZA, Victor (University of Leicester)

Session Classification: Parallel GA09 Binaries

Track Classification: GA-TH