



Contribution ID: 64

Type: **Poster contribution**

## Understanding the anisotropy of cosmic rays at TeV and PeV energies

*Thursday, July 30, 2015 3:30 PM (1 hour)*

The anisotropy in cosmic-ray arrival directions in the TeV-PeV energy range shows both large and small-scale structures. While the large-scale anisotropy may arise from diffusive propagation of cosmic rays, the origin of the small-scale structures remains unclear. We perform three-dimensional Monte-Carlo test-particle simulations, in which the particles propagate in both magnetostatic and electromagnetic turbulence derived from a three-dimensional isotropic power spectrum. However, in contrast to earlier studies, we do not use a backtracking method for the computation of the particle trajectories, and hence anisotropy must build up from a large-scale isotropic (or dipole) boundary condition. It has been recently argued that the turbulent magnetic field itself generates the small-scale structures of the anisotropy if a global cosmic-ray dipole moment is present. Our code is well suited to test that hypothesis. We also investigate the impact of a finite phase velocity of interstellar turbulence.

### Collaboration

– not specified –

### Registration number following "ICRC2015-I"

82

**Primary author:** Mr RETTIG, Robert (Universitaet Potsdam)**Co-author:** POHL, Martin (DESY)**Presenter:** POHL, Martin (DESY)**Session Classification:** Poster 1 CR**Track Classification:** CR-TH