



Contribution ID: 1105

Type: Oral contribution

Non-linear Cosmic Ray propagation close to the acceleration site

Tuesday, 4 August 2015 11:00 (15 minutes)

Recent advances on gamma-ray observations from SuperNova Remnants and Molecular Clouds offer the possibility to study in detail the properties of the propagation of escaping Cosmic Rays (CR). However, a complete theory for CR transport outside the acceleration site has not been developed yet. Two physical processes are thought to be relevant to regulate the diffusion: the growth of waves caused by streaming instability, and possible wave damping mechanisms that reduce the growth of the turbulence. Only a few attempts have been made so far to incorporate these mechanisms in the theory. In this talk I present recent advances in this subject. In particular, I show results obtained by solving the coupled equations for the diffusion of CRs and the evolution of waves. I discuss the importance of streaming instabilities and wave damping in different ISM phases and for different CR energies.

Collaboration

– not specified –

Registration number following "ICRC2015-I"

899

Primary author: Dr NAVA, Lara (HUJI)**Co-authors:** MARCOWITH, Alexandre (LUPM Université Montpellier); Dr MORLINO, Giovanni (Gran Sasso Science Institute); Dr GABICI, Stefano (CNRS/APC); Prof. PTUSKIN, Vladimir (IZMIRAN)**Presenter:** Dr NAVA, Lara (HUJI)**Session Classification:** Parallel CR16 TH prop**Track Classification:** CR-TH