



NEUTRINOS FROM CLUSTERS OF GALAXIES AND RADIO CONSTRAINTS

FABIO ZANDANEL

(GRAPPA INSTITUTE – UNIVERSITY OF AMSTERDAM)


F.ZANDANEL@UVA.NL

WITH IRENE TAMBORRA (GRAPPA), STEFANO GABICI (APC PARIS)
& SHIN'ICHIRO ANDO (GRAPPA) --- A&A 578, A32, 2015

ICRC 2015 – DEN HAAG – JULY 30, 2015


CLUSTERS OF GALAXIES

LARGEST GRAVITATIONALLY BOUND STRUCTURES IN THE UNIVERSE WITH $10^{14} - 10^{15} M_{\odot}$ AND MPC RADII



MINIATI & BERESNYAK (2015)


SHOCKS, TURBULENCE, AND GALAXY OUTFLOWS
→ **COSMIC RAY PROTONS**




BLASI, GABICI & BRUNETTI (2007)

HADRONIC PROTON-PROTON INTERACTIONS

E.G., PFROMMER ET AL. (2008), PINZKE & PFROMMER (2010), ... SEE BRUNETTI & JONES (2014) FOR A REVIEW



NEUTRINOS FROM CLUSTERS



MURASE, AHLERS & LACKI (2013)

CAN THE
SUMMED
OUTPUT OF
ALL GALAXY
CLUSTERS IN
THE UNIVERSE
EXPLAIN THE
ICECUBE
NEUTRINOS?

AARTSEN ET AL. (2014)


E.G., BEREZINSKY ET AL. (1997), MURASE ET AL. (2008), KOTERA ET AL. (2009),
MURASE & BEACOM (2013), MURASE ET AL. (2013), ...

CONSTRAINTS

- **NON-DETECTIONS OF CLUSTERS IN GAMMA RAYS**

REFERENCE: UPPER LIMITS ON
PERSEUS (MAGIC COLL. 2012) AND COMA (ZANDANEL & ANDO 2014)

- **OBSERVED DIFFUSE RADIO EMISSION IN CLUSTERS**



FERETTI ET AL. (2012)

CONSERVATIVE CONSTRAINT:
**NOT OVERSHOOT OBSERVED
CLUSTER RADIO COUNTS!**

REFERENCE: NVSS SURVEY
(GIOVANNINI ET AL. 1999)

WHY CONSERVATIVE?

**WE DO KNOW THAT MOST
EMISSION IS LIKELY NOT FROM
SECONDARIES**

(BRUNETTI ET AL. 2012, ZANDANEL ET AL. 2014)

SIMPLE APPROACH

CALCULATE TOTAL GAMMA-RAY AND NEUTRINO FLUXES FOR ALL CLUSTERS IN THE UNIVERSE USING THEIR **MASS FUNCTION**

WE ADOPT A **GAMMA-RAY LUMINOSITY – MASS RELATION** RESPECTING CONSTRAINTS FROM RADIO TO GAMMA-RAYS

GAMMA LUMINOSITY:

$$L_\gamma = A_\gamma \int \rho_{\text{CR}} \rho_{\text{ICM}} dV$$

RADIO LUMINOSITY:

$$L_f = A_f \int \rho_{\text{CR}} \rho_{\text{ICM}} \frac{\epsilon_B}{\epsilon_B + \epsilon_{\text{CMB}}} \left(\frac{\epsilon_B}{\epsilon_{B_c}} \right)^{\frac{\alpha_p - 2}{4}} dV$$

IGNORING MAGNETIC
FIELD SPATIAL STRUCTURE

$$\frac{L_\gamma}{L_f} = \frac{A_\gamma}{A_f} \frac{\epsilon_B + \epsilon_{\text{CMB}}}{\epsilon_B} \left(\frac{\epsilon_{B_c}}{\epsilon_B} \right)^{\frac{\alpha_p - 2}{4}}$$

ASSUMING
 $B \gg B_{\text{CMB}}$


$$\frac{L_\gamma}{L_f} \approx \frac{A_\gamma}{A_f}$$

WE STUDY THE CASES WITH:

$B \gg B_{\text{CMB}}$, $B = 1 \mu\text{G}$ AND $B = 0.5 \mu\text{G}$


$\alpha_p = 1.5, 2, 2.2$ AND 2.4

RADIO COUNTS



ZANDANEL, TAMBORRA, GABICI & ANDO (2015)


TOTAL GAMMAS AND NEUTRINOS




ZANDANEL, TAMBORRA, GABICI & ANDO (2015)

RADIO LOUD/QUIET CLUSTERS

NOT ALL HOST
RADIO EMISSION




E.G., CASSANO ET AL. (2013)



ZANDANEL, TAMBORRA, GABICI & ANDO (2015)

TOTAL GAMMAS AND NEUTRINOS




ZANDANEL, TAMBORRA, GABICI & ANDO (2015)

ANISOTROPY POWER SPECTRUM

FULL SPATIAL MODELING OF CRS, GAS AND MAGNETIC FIELD

(FOLLOWING ZANDANEL, PFROMMER & PRADA 2014) → WE CAN ESTIMATE THE

POWER SPECTRUM OF GAMMA-RAY ANISOTROPIES



ZANDANEL, TAMBORRA, GABICI & ANDO (2015)

SUMMARY & CONCLUSIONS

- WE CONSIDER RADIO CONSTRAINTS FOR THE FIRST TIME
 - CLUSTERS $\leq 10\%$ OF ICECUBE FLUX FOR $\alpha_p = 2$
 - MUCH LESS FOR $\alpha_p > 2$, BUT CAN EXPLAIN DATA FOR SPECULATIVE VALUES OF $\alpha_p < 2$
- CONTRIBUTION TO DIFFUSE GAMMA-RAY BACKGROUND ALWAYS NEGLIGIBLE + ANISOTROPIES
 - ICECUBE CAN TEST MOST OPTIMISTIC SCENARIOS STACKING NEARBY SOURCES

OUR CONCLUSIONS ARE VALID FOR ALL SOURCES INJECTING CRS INTO THE INTRA-CLUSTER MEDIUM!