



Contribution ID: 44

Type: **Poster contribution**

## CORSIKA modification for rigidity dependent primary selection based on Geomagnetic cutoff rigidity for GRAPES-3 simulations

*Thursday 30 July 2015 15:30 (1 hour)*

For the analysis of the GRAPES-3 Muon data, large scale Monte Carlo simulations are required. These simulations are performed using the CORSIKA simulation package developed by the KIT group. However, the geomagnetic cutoff rigidity varies with direction, therefore, a constant threshold for selection of primary energy results in generation of a large number of events that are subsequently rejected due to their rigidity being below the cutoff value in some directions. We have implemented an efficient mechanism in CORSIKA to select only those primary cosmic rays that lie above the cutoff rigidity in a given direction resulting in rejection of those primary cosmic rays that would have otherwise been rejected subsequently. Results based on actual simulations of GRAPES-3 Muon data have shown that by using this rigidity based cut, the actual computation time was reduced by a factor of two without compromising the reliability of the results.

### Collaboration

– not specified –

### Registration number following "ICRC2015-I"

54

**Author:** Mr BALAKRISHNAN, Hari Haran (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India)

**Co-authors:** Dr OSHIMA, Akitoshi (Chubu University, Engineering Science Laboratory, Kasugai-shi, Aichi 487-8501, Japan); Mr CHANDRA, Anuj (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Mr JAIN, Atul (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mu); Mr RAO, B Srinivasa (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Mr P, Jagadeesan (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Dr NAYAK, Pranaba K (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Dr MOHANTY, Pravata K (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Prof. KAWAKAMI, Saburo (Graduate School of Science, Osaka City University, Osaka 558-8585, Japan); Mr MORRIS, Sam D (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Prof. AHMAD, Shakeel (Department of Physics, Aligarh Muslim University, Aligarh 202002, India); Prof. DUGAD, Shashi R (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road,

Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Prof. GUPTA, Sunil K (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India); Prof. HAYASHI, Yohio (Graduate School of Science, Osaka City University, Osaka 558-8585, Japan)

**Presenter:** Mr BALAKRISHNAN, Hari Haran (HECR Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India and GRAPES-3 Experiment, Cosmic Ray Laboratory, Ooty 643 001, India)

**Session Classification:** Poster 1 CR

**Track Classification:** CR-TH