

Monte-Carlo event generator CompHEP (version 4.5)

Alexander Sherstnev
University of Oxford and
Moscow State University

For CompHEP Collaboration

Outline

- Motivation behind CompHEP
- CompHEP in nutshell
- Parallel calculations
- Implementation of Les Houches formats
- CompHEP-Interfaces package
- HepML
- Final remarks

Motivations

Increasing collider energy and luminosity require calculations of processes with more and more particles in the final state with better and better precision (NLO, NNLO, NLL resummation)

1. SLAC/LEP I – basically 2 fermion physics;
2. LEP II – basically 4 fermion physics;
3. TEVATRON, LHC, and (I)LC – 4,5,6 and even 8 fermions + additional photons and/or gluons (jets);
 - Single top in the t-channel mode – 5 fermions;
 - Top pair production with decays – 6 fermions;
 - Strongly interacting Higgs sector in hadron collisions – 6 fermions

$$pp \rightarrow W^- W^+ + jj$$

- Yukawa coupling – 8 fermions

$$pp \rightarrow t \bar{t} + H$$

Motivations

Large number of Feynman diagrams and large number of subprocesses (hadron colliders) require automation

Goals:


- Automation of tree level diagram calculations
- “Unification” of symbolical and numerical calculation (in UI) – a full computational chain for phenomenologists
- Interfacing to other generators (for showering and hadronization) and further (full simulation)
- Interfacing to NLO codes: cross section calculators, mass spectrum calculators

Community answer: CompHEP, GRACE, MadGraph, AlpGen, Omega/WHIZARD, Sherpa/Amegic, etc.

CompHEP

(Computation in High Energy Physics)

Incomplete list of processes simulated with CompHEP in the past (3 papers have ~1000 citations):



CompHEP conception

- CompHEP constructs tree level Feynman diagrams for a given parton process ($u, U \rightarrow e, E, m, M, G$ or $p, p \rightarrow e, E, m, M, j$)
- Symbolical calculations of the Feynman diagrams squared
- Automated preparation of binary for numerical calculations by Monte-Carlo technique (based on C code): cross sections, distributions, events
- Model independence: Model description files are input for CompHEP. CompHEP can work with 0, 1/2, 1-spin particles, Majorana and Dirac spinors, 3- and 4- vertices with fields, derivatives of fields, functions of parameters
- “Universal”, build-in symbolic calculator: CompHEP can calculate N^2 tree level diagrams for any process $1, 2 \rightarrow M$. N and M are limited by computer resources only
- User-friendly interface: GUI for both symbolic and numerical parts, comprehensive build-in help (F1), batch scripts

CompHEP Model

**CompHEP Model defines particles and their interactions.
Technically CompHEP model is a set of 5 text files (tables)**

- A set of fundamental particles: names, mass/width, spin, charges
- Numerical model parameters: mass/width values, couplings, mixing parameters, etc.
- Constrains: relations between the parameters
- Lagrangian: a set of all interaction vertexes
- Composite particles: proton, artificial useful particle combinations

CompHEP SM Model

[test_trunk_bb : tcsh] CompHEP version 4.5... CompHEP version 4.5... CompHEP version 4.5... CompHEP version 4.5...

CompHEP version 4.5.0rc6

Variables

Clr	Rest	Del	Size	Name	Value	>	Comment	<
EE				Elementary charge (alpha=1/127.9, on-shell, MZ)	0.31345			
GG				Strong coupling constant (Z pnt, alp=0.1172pm0)	1.21358			
SW				sin of the Weinberg angle (MZ point -> MW=79.9)	0.48076			
s12				Parameter of C-K-M matrix (PDG2002)	0.2229			
s23				Parameter of C-K-M matrix (PDG2002)	0.0412			
s13				Parameter of C-K-M matrix (PDG2002)	0.0036			
MZ				mass of Z boson	91.1876			
wZ				width of Z boson	2.43631			
wW				width of W boson	2.02798			
Mm				mass of muon	0.10566			
Mtau				mass of tau-lepton	1.77699			
Mc				mass of c-quark	1.65			
Ms				mass of s-quark	0.117			
Mtop				mass of t-quark	174.3			
wtop				width of t-quark	1.54688			
Mb				mass of b-quark	4.85			
MH				mass of Higgs	115			
wH				width of Higgs	0.0061744			

F1 F2 Top Bottom GoTo Find Zoom ErrMes

CompHEP version 4.5.0rc6

Particles

Clr	Rest	Del	Size	Full name	P	aP	2*spin	mass	width	color	aux	>	LaTeX(A)	<
				gluon	G	G	2	0	0	8	G	G		G
				photon	A	A	2	0	0	1	G	A		A
				Z boson	Z	Z	2	MZ	wZ	1	G	Z		Z
				W boson	W+	W-	2	MW	wW	1	G	W+		W+
				neutrino	ne	Ne	1	0	0	1	L	\nu^e		\nu^e
				electron	e	E	1	0	0	1		e		e
				mu-neutrino	mm	Nm	1	0	0	1	L	\nu^\mu		\nu^\mu
				muon	m	M	1	Mm	0	1		m		m
				tau-neutrino	nl	Nl	1	0	0	1	L	\nu^\tau		\nu^\tau
				tau-lepton	l	L	1	Mtau	0	1		l		l
				u-quark	u	U	1	0	0	3		u		u
				d-quark	d	D	1	0	0	3		d		d
				c-quark	c	C	1	Mc	0	3		c		c
				s-quark	s	S	1	Ms	0	3		s		s
				t-quark	t	T	1	Mtop	wtop	3		t		t
				b-quark	b	B	1	Mb	0	3		b		b
				Higgs	H	H	0	MH	wH	1		H		H

F1 F2 Top Bottom GoTo Find Zoom ErrMes

CompHEP version 4.5.0rc6

Constraints

Clr	Rest	Del	Size	Name	>	Expression	<
C1						$\sqrt{1-SW^2}$	
c12						$\sqrt{1-s12^2}$	
c23						$\sqrt{1-s23^2}$	
c13						$\sqrt{1-s13^2}$	
Vud						$c12*c13$	
Vus						$s12*c13$	
Vub						$s13$	
Vcd						$-s12*c23-c12*s23*s13$	
Vcs						$c12*c23-s12*s23*s13$	
Vcb						$s23*c13$	
Vtd						$s12*s23-c12*c23*s13$	
Vts						$-c12*s23-s12*c23*s13$	
Vtb						$c23*c13$	
MW						$MZ*CW$	

F1 F2 Top Bottom GoTo Find Zoom ErrMes

CompHEP version 4.5.0rc6

Lagrangian

Clr	Rest	Del	Size	P1	P2	P3	P4	>	Factor	<
C				b	W+				$-EE*\sqrt{2}*Vcb/(4*SW)$	G(m)
C				b	W+.f				$i*EE*\sqrt{2}*Vcb/(4*MW*SW)$	Mb*
C				c	A				$-2*EE/3$	G(m)
C				c	G				GG	G(m)
C				c	H				$-EE*Mc/(2*MW*SW)$	1
C				c	Z				$-EE/(12*CW*SW)$	(3-
C				c	Z.f				$i*EE*Mc/(2*MW*SW)$	G5
C				d	W+				$-EE*\sqrt{2}*Vcd/(4*SW)$	G(m)
C				d	W+.f				$-i*EE*Mc*\sqrt{2}*Vcd/(4*MW*SW)$	(1-
C				s	W+				$-EE*\sqrt{2}*Vcs/(4*SW)$	G(m)
C				s	W+.f				$i*EE*\sqrt{2}*Vcs/(4*MW*SW)$	Ms*
D				c	W-				$-EE*\sqrt{2}*Vcd/(4*SW)$	G(m)
D				c	W-.f				$i*EE*Mc*\sqrt{2}*Vcd/(4*MW*SW)$	(1+
D				d	A				EE/3	G(m)
D				d	G				GG	G(m)
D				d	Z				$-EE/(12*CW*SW)$	2*S
D				t	W-				$-EE*\sqrt{2}*Vtd/(4*SW)$	G(m)
D				t	W-.f				$i*EE*Mtop*\sqrt{2}*Vtd/(4*MW*SW)$	(1+
D				u	W-				$-EE*\sqrt{2}*Vud/(4*SW)$	G(m)
E				e	A				EE	G(m)
E				e	Z				EE/(4*CW*SW)	(1-

F1 F2 Top Bottom GoTo Find Zoom ErrMes

Basic and user-defined CompHEP Models

- Simple training models: QED, Effective 4-fermion Fermi model
- SM in two different gauges: unitary gauge and t 'Hooft-Feynman gauge. Flavour simplified SM model (#-model)
- SUSY Models: unconstrained MSSM (again in two gauges); SUGRA model; GMSB model

New (user-defined) Models

- Simple way (if your model is relatively simple): add new particles/params/vertices
- For more complicated models: LanHEP – a program for generation of Feynman rules for user-defined model (**developed by A.Semenov**)
 - Works with super-multiplets and superpotential
 - Generates all needed files for CompHEP (also FeynArts and LaTeX format)
 - Several options for self-checking (charge conservation, BRST invariance, etc.)
 - Has been used for CompHEP SUSY models and many other BSM models

CompHEP BSM Lagrangians

- **Complete Leptoquark model.** Includes Yukawa couplings for all types of LQ, gauge couplings and anomalous gauge couplings for vector LQ (by request)
- **Top quark Lagrangian with anomalous couplings** as follows from the dimension 6 effective operators (by request)
- **Excited fermion Model** (by request)
- **Complete two-Higgs-doublet Model with conserved or broken CP invariance** (by request)
- **RS1 model and effective 4 particle Lagrangian for RS below KK threshold**
- **UED** model (Matchev et al.)
- **Minimal Higgsless Model** (Chivukula et al.)
- **Exotics:** Muonic photon; para-photon; E6 isosinglet quark; Z' , W' bosons; doubly charged Higgs, color octet pseudoscalars, Inert Douplet Model, etc.

Symbolic calculations

The image shows four windows of the CompHEP 4.5.0rc6 software interface. The top-left window displays the 'Abstract' section, describing the package's purpose for calculating decay and collision processes. The top-right window shows the 'Model' selection screen, currently set to 'SM, Feynman gauge', with a menu for entering decay or scattering processes. The bottom-left window lists 'List of structure functions' with 17 options, including PDF sets like cteq6m and cteq6l1. The bottom-right window shows the 'List of composite particles' with three entries: j1 (u, U, d, D, G), j2 (u, U, d, D, s, S, c, C, G), and j3 (u, U, d, D, s, S, c, C, b, B, G). A command line at the bottom of the bottom-right window shows 'Enter Final State: p, p -> e, E, m, M, j2'. The interface includes a menu bar at the bottom with options like F1-Help, F2-Man, F5-Switches, F6-Results, and F9-Quit.

Abstract

CompHEP package is created for calculation of decay and high energy collision processes of elementary particles in the tree approximation. The main idea put into the CompHEP was to make available passing from the Lagrangian to the final distributions effectively, with the high level of automatization. Use the F2 key to get the information about interface facilities and the F1 key to get online help.

- * QED
- Effective 4-fermion
- SM, unitary gauge
- SM, Feynman gauge**
- MSSM, unitary gauge
- MSSM, Feynman gauge
- SUGRA, Feynman gauge
- GMSB, Feynman gauge
- _SM_ud
- _SM_qQ
- CREATE NEW MODEL

F1-Help F2-Man F5-Switches F6-Results F9-Quit

Model: SM, Feynman gauge

Abstract

CompHEP package is created for calculation of decay and high energy collision processes of elementary particles in the tree approximation. The main idea put into the CompHEP was to make available passing from the Lagrangian to the final distributions effectively, with the high level of automatization. Use the F2 key to get the information about interface facilities and the F1 key to get online help.

- * Enter Decay Process
- Enter Scattering Process
- Edit Beams Table
- Edit Str. Functions Table
- Edit Model
- Delete Model

F1-Help F2-Man F5-Switches F6-Results F9-Quit

Model: SM, Feynman gauge

List of structure functions

- 0: OFF
- 1: WWA ($m=0.000511$ Ch=-1 Q=100)
- 2: Laser Photons
- 3: ISR(100 Beamstr.:OFF)
- 10: PDF:cteq6m(proton)
- 11: PDF:cteq6m(anti-proton)
- 12: PDF:cteq6l1(proton)
- 13: PDF:cteq6l1(anti-proton)
- 14: PDF:cteq6d(proton)
- 15: PDF:cteq6d(anti-proton)
- 16: PDF:cteq5m1(proton)
- 17: PDF:cteq5m1(anti-proton)

PgDn

Enter 1st Beam: p

Enter 1st Beam Energy (GeV) : 7000.000000

Enter 2nd Beam: p

Enter 2nd Beam Energy (GeV) : 7000.000000

Enter PDF number : 12

Model: SM, Feynman gauge

List of composite particles (switch to the particle list by F3)

- Name: j1 (u, U, d, D, G)
- Name: j2 (u, U, d, D, s, S, c, C, G)
- Name: j3 (u, U, d, D, s, S, c, C, b, B, G)

Enter Final State: p, p -> e, E, m, M, j2

Applications Places System Thu 30 Oct, 09:37 Alexander Sherstnev

Symbolic calculations (2)

CompHEP version 4.5.0rc6

Model: SM, Feynman gauge

List of (anti)particles

G(G)	gluon	A(A)	photon	Z(Z)	Z boson
W+(W-)	W boson	ne(Ne)	neutrino	e(E)	electron
nm(Nm)	mu-neutrino	m(M)	muon	nl(Nl)	tau-neutrino
l(L)	tau-lepton	u(U)	u-quark	d(D)	d-quark
c(C)	c-quark	s(S)	s-quark	t(T)	t-quark
b(B)	b-quark	H(H)	Higgs		

Enter Final State: p,p -> e,E,j1
 Exclude diagrams with H
 Keep diagrams with

CompHEP version 4.5.0rc6

Model: SM, Feynman gauge

Process: p,p -> e,E,j1

Feynman diagrams

64 diagrams in 16 subprocesses are constructed.
 0 diagrams are deleted.

NN	Subprocess	Del	Rest
1	u,U -> G,e,E	0	4
2	u,G -> e,E,u	0	4
3	d,D -> G,e,E	0	4
4	d,G -> e,E,d	0	4
5	U,u -> G,e,E	0	4
6	U,G -> e,E,U	0	4
7	D,d -> G,e,E	0	4
8	D,G -> e,E,D	0	4
9	s,S -> G,e,E	0	4
10	c,C -> G,e,E	0	4
11	S,s -> G,e,E	0	4
12	C,c -> G,e,E	0	4

PgDn

F1-Help F2-Man F3-Model F5-Switches F6-Results F7-Del F8-UnDel F9-Quit

CompHEP version 4.5.0rc6

Delete, On/off, Restore, Latex

F1-Help, F2-Man, PgUp, PgDn, Home, End, #, Esc

CompHEP version 4.5.0rc6

Model: SM, Feynman gauge

Process: p,p -> e,E,j1

Feynman diagrams

64 diagrams in 16 subprocesses are constructed.
 0 diagrams are deleted.

Squared diagrams

160 diagrams in 16 subprocesses are constructed.
 0 diagrams are deleted.
 160 diagrams are calculated.
 0 Out of memory

Write results
 C-compiler
 Enter new process

F1-Help F2-Man F3-Model F4-Diagrams F5-Switches F6-Results F9-Quit

Numerical calculations

- **Customize numerical MC generator:** The most complicated part: do proper phase space sampling (regularizations + kinematics), set necessary kinematic cuts, Q^2 , PDF set, etc. Main goals – to improve efficiency of MC calculation and describe physics task. User may change model parameters and set kinematic cuts
- **Calculate full cross section and distributions:** CompHEP uses an improved version of the adaptive VEGAS algorithm for MC calculations. User may order different variables (P_T , inv. mass, rapidity, etc.) for histogramming
- **Generate events:** As soon as CompHEP customized, events can be generated for the subprocess. User set a number of the events They are kept to text files.

If the process consists of some subprocesses, the procedure is applied to the each subprocess.

Numerical calculations (2)

The image shows four windows of the CompHEP version 4.5.0rc6 software. Each window displays the same process: $p, p \rightarrow e, E, j1$ (16 subprocesses) and $(\text{sub})\text{Process: } u, U \rightarrow G, e, E$. The Monte Carlo session is 1(begin).

Top-left window: Shows a menu for "Initial state" with options: Subprocess, Initial state, Model parameters, Constraints, QCD scale, Width scheme: Fixed, Cuts, Kinematics, Regularization, and Numerical Session.

Top-right window: Shows the "Initial state" details:
Beam particle 1: proton
Beam particle 2: proton
Str.Fun.1: PDF:cteq611
Str.Fun.2: PDF:cteq611
1 particle momentum[GeV] = 7000
2 particle momentum[GeV] = 7000

Bottom-left window: Shows the "Model parameters" menu with options: Model parameters, Change parameter, and a list of parameters: EE= 0.31345, SW= 0.48076, NZ= 91.188, wZ= 2.4363, Mc= 1.65, Ms= 0.117, GG= 1.2136.

Bottom-right window: Shows the "QCD scale" menu with options: QCD scale, QCD alpha, and QCD Lambda6= ???
Q(GeV) = m45
Alpha(Q) plot

Each window has a menu bar at the bottom: F1-Help F2-Man F4-Diagrams F6-Results F9-Quit (with F7-Plot in the top-right window).

Numerical calculations (3)

CompHEP version 4.5.0rc6

Process: p,p -> e,E,j1 (16 subprocesses)
(sub)Process: u,U -> G,e,E

Cuts 3

Clr	Rest	Del	Size
Parameter	>	Min bound	< > Max bound <
t3		10	
y3		-5	5
m45		10	

F1-F2-Top-Bottom-GoTo-Find-Zoom-ErrMes

CompHEP version 4.5.0rc6

Process: p,p -> e,E,j1 (16 subprocesses)
(sub)Process: u,U -> G,e,E

Regularization 4

Clr	Rest	Del	Size
Momentum	>	Mass	< > Width <
45		0	1
45		MZ	wZ 2
13		0	2
14		0	0 2

F1-F2-Top-Bottom-GoTo-Find-Zoom-ErrMes

CompHEP version 4.5.0rc6

Process: p,p -> e,E,j1 (16 subprocesses)
(sub)Process: u,U -> G,e,E
Monte Carlo session: 1(begin)

Numerical Session

#IT	Cross section [pb]	Error %	nCall	chi**2

1	1.1705E+02	1.81E+00	90720	
2	1.1432E+02	1.16E+00	90720	
3	1.1540E+02	7.58E-01	90720	
4	1.1572E+02	1.03E+00	90720	
>	1.1539E+02	5.18E-01	362880	0.4
1	1.1613E+02	8.32E-01	90720	
2	1.1553E+02	6.48E-01	90720	
3	1.1679E+02	5.87E-01	90720	
4	1.1806E+02	9.86E-01	90720	
>	1.1644E+02	3.59E-01	362880	1
5	1.1465E+02	7.81E-01	90720	
6	1.1644E+02	8.01E-01	90720	
7	1.1518E+02	6.64E-01	90720	
8	1.1580E+02	5.92E-01	90720	
>	1.1596E+02	2.49E-01	725760	1

F1-Help F2-Man F4-Diagrams F6-Results F9-Quit

CompHEP version 4.5.0rc6

Process: p,p -> e,E,j1 (16 subprocesses)
(sub)Process: u,U -> G,e,E
Monte Carlo session: 3(begin)

Numerical Session

Start integration

Integration is over
Press any key

Batch system in CompHEP

Both symbolic & numerical parts have batch scripts:
Perl scripts: **symb_batch.pl** and **num_batch.pl**

Why the scripts are useful?

- Computations of many subprocesses – laborious task, can be significantly simplified especially for hadron colliders
- **Long/large-scale calculations:** GUI is not too handy
- **Support of parallel calculations:** very helpful for multi-CPU machines/computer clusters (pbs/lsf is available; grid in progress)
- **“Knowledge transfer”:** theorists/phenomenologists can prepare model/process.dat/batch.dat for further simulations by experimentalists

Symbolic parallel calculations

- Main idea:

- symbolic calculation of one diagram is an independent task. The only unified point is the final binary data file.
- Several calculation flows can be running at one time for several subsets of diagrams. The final point is to collect the binary data file
- Implemented in `./symb_batch.pl` with the option `-mp`
- 1st step! `./symb_batch.pl -help`
- Parallel calculations on one machine: `-mp N` means N symbolic calculations in parallel
- Batch system version (pbs/lsf) is being implemented
- Very easy to use!

Numerical parallel calculations

- Again, 1st step: `./num_batch.pl -help` (long and very detailed description). The script has lots of options (~30)!
- `n_comphep.exe` should be prepared! Main file is `batch.dat` in `/results` (based on `session.dat`). It can be edited by hand or via GUI and `./num_batch.pl -add` (customized subprocess added to `batch.dat`)
- Then `./num_batch.pl -run vegas` (cross section calculation for ALL subprocesses) and `./num_batch.pl -run max,evnt` (event generation)
- Parallel calculations available. Alone machine (useful for multi-CPU desktops): `-mp 3` (3 jobs are calculated simultaneously); computer clusters (with a batch system installed): `-lsf` and `-pbs`
- Many ways to present and monitor results and calculation process (see help)
- Very easy to use!

Event generation

- Events – phase space points, distributed according to $|M|^2$
- Monte-Carlo technique
- adaptive importance sampling method VEGAS
- For event generation: + stratified sampling
- Von Neumann (rejection) sampling:
 - If $g(x)$ – importance sampling function
 - $|M|^2 = f(x)$ – matrix element squared
 - Find $\omega_{\max}(x_0) = f(x_0)/g(x_0)$ and compare $\rho = \omega(x_i)/\omega_{\max} = \omega_i/\omega_{\max}$ and random number R
 - If $\rho > R$ – accept the point

Iterative Rejection sampling

- Usual efficiency in CompHEP – 0.1-1%.

What can we do with “waste”?


- Repeat the von Neumann procedure with the waste – rejected events!
- But importance sampling function is unknown...
- There is a way to calculate it

$$\omega_1 = \frac{N_{rej} \omega_0}{N_{tot} - N_{act} \omega_0 / I_{tot}}$$

- We can do several iterations, but should stop at some step.
- Stopping rule: $|I_0 - I_f| \geq \Delta I_0 + \Delta I_f$
- Real benefit: from 20% to 1000% of extra events **without** extra calculations of $|M|^2$

more details in 0807.2823

Modern Monte-Carlo Model



MC code landscape: diversity

General-purpose

Special

Hard Processes

A lot (CompHEP,
MadGraph, Grace)

Parton Showers

PYTHIA

Ariadne, NLLjet

Underlying Events

HERWIG

PHOJET, Jimmy

SHERPA

Hadronization

ISAJET

none

Decays

TAUOLA, EvtGen

Les Houches Agreements

There are many MC generators with their own advantages and application areas. Often we are forced to use several generators for reliable calculations:

Problems:

- Interfacing some MC codes (ME and SH generators): Les Houches Accord 1, Les Houches Event format
- Les Houches Accord 2: uniform interface to different PDF sets (LHAPDF package)
- Les Houches Accord 3: Interfacing SUSY codes to MC generators for parameters, spectrum, decays (SPA).
- BSM Les Houches Accord: fixing of parameter record for BSM
- Matching ME (LO/NLO) and SR(NL): CKKW, MC@NLO, Mrenna-Richardson, MLM, ...

LHEF, LHAPDF, SUSY LHA, BSM LHA

1. **LHA I** is implemented in CompHEP-Interfaces
2. **LHEF** - the format adopted by many developer groups (hep-ph/060917). Now CompHEP supports 3 event formats: cpyth-1, cpyth-2 (for experiments, where the formats are used), and LHEF with HepML header. There is a special option - Generator (LHEF format) - in the event menu in n_comphep
3. All modern PDFs are available via **LHAPDF**: CTEQ, MRST, Alekhin PDF, etc. Both options, LHAPDF and internal PDF, are available in CompHEP 4.5 with the same functionality in both regimes
4. **SUSY LHA** The SLHA interface is implemented in SUGRA and GMSB models of CompHEP. By default, the slhaScript file invokes SUSPECT
5. **BSM LHA** is still being implemented

CompHEP-interfaces package

The CompHEP-Interfaces includes two interfaces for PYTHIA and HERWIG, These interfaces allow us to use processes computed by CompHEP as external processes in PYTHIA/HERWIG

Main goal: provide ISR/FSR, hadronization (jet fragmentation), and decays as it is done in PYTHIA/HERWIG

- CompHEP generates unweighted events (event files)
- The command *mix_flows* mixes several event files in one event file
- Some governing parameters (Event file name, the number of events for generation) are kept in the file **INPARAM.DAT**
- A matching code for ME events and showers in PYTHIA are being developed in the package
- Simple routines for toy analysis are provided
- Program to translate data to ROOT file (record looks like LHA I)
- TAOLA interface available (be request)

HepML

- Unified XML format of MC event files metadata
 - to keep comprehensive information on event
 - to store generator input parameters and setup
 - an effort to fix a unified extensible way of MC events description
 - the LHEF standard permits XML code in event file headers
- Main purposes:
 - to unify MC event files description (parton and particle levels of MC simulation)
 - to facilitate passing information from Matrix Element generators to Shower generators
 - to simplify MC generators tuning and testing
- Contributors
 - CEDAR <http://www.cedar.ac.uk>
 - LCG MCDB <http://mcdb.cern.ch>
- Homepage <https://twiki.cern.ch/twiki/bin/view/Main/HepML>

CompHEP can
generate
HepML code!

more details in S. Belov's talk

Concluding Remarks

- CompHEP with the interface to PYTHIA/HERWIG is a powerful tool for a simulation of SM/BSM physics at hadron and lepton colliders
- CompHEP can calculate cross sections, build different distributions, and generate un-weighted events
- CompHEP is compatible with all modern “Monte-Carlo industry” standards (Les Houches Accords 1, 2, 3, LHE). This CompHEP-Interfaces can be easily used and included to experimental environments
- Parallel computations both in symbolic and numerical modules are implemented as part of batch scripts
- Advanced MC techniques for improving of generation efficiency applied
- In order to facilitate interfacing of different MC code and re-usage event samples CompHEP generates HepML code
- HepML is a new method to keep comprehensive information on events in event files

General information and references

- CompHEP collaboration: E. Boos, V. Bunichev, M. Dubinin, L. Dudko, V. Ilyin, A. Kryukov, V. Edneral, V. Savrin (Moscow State), A. Semenov (JINR, Dubna), A.S. (Moscow State and Oxford University)
- CompHEP homepage: <http://comphep.sinp.msu.ru>
- References:
 - CompHEP: E. Boos et al., Nucl.Inst.Meth. A534:250 (2004) [hep-ph/0403123]
 - LanHEP: A. Semenov, Nucl.Inst.Meth. A393:293 (1997) [hep-ph/0403123]; 0805.0555 (hep-ph)
 - CompHEP-Interfaces: A.Belyaev et al., hep-ph/0101232
 - MCDB/HepML: S.Belov et al. Comput.Phys.Comm. 178:222 [hep-ph/0703287]

Backup slides

MCDB – Monte-Carlo events Data Base

Team: S.Belov, L. Dudko, A.Ribon, A.S. (JINR, MSU, CERN, Oxford)

Motivation:

- Verified MC simulation of complicated processes requires sophisticated expertise and expert knowledge
- A physics group in a collaboration requests experts and/or MC generator authors to create MC samples for the particular process
- The same physics processes are investigated by various physics groups, the same MC samples can be used in different analyses

The main motivation – to make MC event samples, prepared by experts, available for various physics groups

MCDB tasks:

- The database has to be available via the Web and Grid
- Using CASTOR/GRID technologies to keep/upload/download MC samples
- 7. Simple and intuitive interface for events authors and end-users to find and manipulate event samples

more details in S. Belov's talk

MCDB: technical details

- Frontends: a Web site (mcdb.cern.ch) and API (via HepML) and Grid
- Backend: SQL for metadata and CASTOR for data
- Keep parton and particle level events with standard interface to the next level of simulation (PYTHIA/HERIWG, simu. software), based on LHA I
- Store detailed documentation for each set of event samples
- Provide communication between users and experts via MCDB web pages
- Direct programming interface of the collaboration software to LCG MCDB
- **Divided in two zones:**
 - **public area:** users can search for/browse the DB and download event files
 - **restricted area:** authors (experts) change MCDB content: upload and describe new event files, change the existed files and reply to user's comments

MCDB encourages end-users to cite event sample author's papers in case the events are used in physics analyses!

Paper: hep-ph/0703287