

α clusters in ultra-relativistic light-ion + Pb collisions

Maciej Rybczyński

Institute of Physics, Jan Kochanowski University, Kielce

Based on

W. Broniowski, E. R. Arriola, Phys. Rev. Lett. **112**, 112501 and

P. Bozek, W. Broniowski, E. R. Arriola, M. Rybczyński, Phys. Rev. **C90**, 064902

Outline

Two phenomena are related:

α clustering in light nuclei

and

harmonic flow in ultra-relativistic nuclear collisions

- ✓ New method of investigating many-particle nuclear correlations
- ✓ Another test of collective dynamics/harmonic flow

Some history

David Brink: *After Gamow's theory of α -decay it was natural to investigate a model in which nuclei are composed of α -particles. Gamow developed a rather detailed theory of properties in his book "Constitution of Nuclei" published in 1931 before the discovery of the neutron in 1932. He supposed that $4n$ -nuclei like ${}^8\text{Be}$, ${}^{12}\text{C}$, ${}^{16}\text{O}$... were composed of α -particles.*







Fig. 1. Alpha-particle configuration for some $4N$ nuclei.



α clusters in light nuclei



${}^9\text{Be}$



${}^{12}\text{C}$

ground

Hoyle 0^+



other excited, 2^+ ...

How can we detect the α clusters in the ground state?
What is their spatial arrangement?
Assessment of n-body correlations (one-body not enough)

Precursor: d-A by Bożek

The deuteron has an intrinsic dumbbell shape with very large deformation: $r_{\text{rms}} \approx 2$ fm

Initial entropy density
in a d-Pb collision with $N_W = 24$




Resulting large elliptic flow confirmed with the later RHIC analysis

Details:

P. Bożek, Phys. Rev. **C85**, 014911

From α clusters to flow in relativistic collisions

α clusters \rightarrow asymmetry of shape \rightarrow asymmetry of initial fireball \rightarrow
hydro or transport \rightarrow collective harmonic flow




nuclear elliptic geometry \rightarrow fireball elliptic geometry \rightarrow elliptic flow

What are the signatures, chances of detection?
(some blurring by fluctuations)
„Easy snap-shot but difficult development”

From α clusters to flow in relativistic collisions






α clusters \rightarrow asymmetry of shape \rightarrow asymmetry of initial fireball \rightarrow
hydro or transport \rightarrow collective harmonic flow



nuclear triangular geometry \rightarrow fireball triangular geometry \rightarrow triangular flow

What are the signatures, chances of detection?
(some blurring by fluctuations)
„Easy snap-shot but difficult development”






${}^7\text{Be}+{}^{208}\text{Pb}$ - single events



$v_{\text{NN}}=17$ GeV
most central ($b=0$ fm) collisions






All simulations with GLISSANDO 2

${}^9\text{Be}+{}^{208}\text{Pb}$ - single events



$v_{\text{NN}}=17$ GeV
most central ($b=0$ fm) collisions

$^{12}\text{C}+^{208}\text{Pb}$ - single events



$v_{\text{NN}}=17$ GeV
most central ($b=0$ fm) collisions

Eccentricity parameters

We need some quantitative measures of deformation (heavily used in heavy-ion analyses).
Eccentricity parameters:

$$\epsilon_n \exp(in\Phi_n) = \frac{\sum_j r_j^n \exp(in\phi_j)}{\sum_j r_j^n}$$

describe the shape of each event.

j labels the sources in the event, n =rank, Φ_n is the principal axis angle, $r = \sqrt{x^2 + y^2}$, $\tan(\phi) = \frac{y}{x}$

$n = 2$ – ellipticity, $n = 3$ – triangularity, . . .

Two components:

- ✓ intrinsic (from existent mean deformation of the fireball)
- ✓ from fluctuations

Geometry vs multiplicity correlations in $\text{Be} + {}^{208}\text{Pb}$ collisions

Two cases of angular orientation
cluster plane **parallel** or **perpendicular** to the transverse plane




higher multiplicity
higher ellipticity

lower multiplicity
lower (no) ellipticity

Geometry vs multiplicity correlations in $^{12}\text{C}+^{208}\text{Pb}$ collisions


Two cases of angular orientation
cluster plane **parallel** or **perpendicular** to the transverse plane



higher multiplicity
higher triangularity
lower ellipticity

lower multiplicity
lower triangularity
higher ellipticity

Ellipticity and triangularity vs multiplicity in $^{12}\text{C}+^{208}\text{Pb}$ collisions



Clusters: (qualitative signal!)

When $N_w \nearrow$ then $\langle\epsilon_3\rangle \nearrow$, $\langle\epsilon_2\rangle \searrow$, $\sigma(\epsilon_3)/\langle\epsilon_3\rangle \searrow$ and $\sigma(\epsilon_2)/\langle\epsilon_2\rangle \nearrow$

No clusters:

similar behavior for $n = 2$ and $n = 3$

Details:

W. Broniowski, E. R. Arriola, Phys. Rev. Lett. **112**, 112501

P. Bozek, W. Broniowski, E. R. Arriola, M. Rybczyński, Phys. Rev. **C90**, 064902

Cumulants, moments and its ratios

$$v_n = \kappa_n \epsilon_n$$

$$v_n\{m\} = \kappa_n \epsilon_n\{m\} \quad n = 2, 3, \dots \quad m = 2, 4, 6, \dots$$

$$\frac{v_n\{m\}}{v_n\{2\}} = \frac{\epsilon_n\{m\}}{\epsilon_n\{2\}}$$


$$\frac{\sigma(v_n)}{\langle v_n \rangle} = \frac{\sigma(\epsilon_n)}{\langle \epsilon_n \rangle}$$

$$\epsilon_n\{2\}^2 = \langle \epsilon_n^2 \rangle$$




$$\epsilon_n\{4\}^4 = 2\langle \epsilon_n^2 \rangle - \langle \epsilon_n^4 \rangle$$

$\sqrt{s_{NN}}=17$ GeV, central rapidity

${}^7\text{Be}+{}^{208}\text{Pb}$




${}^9\text{Be}+{}^{208}\text{Pb}$




$v_{s_{NN}}=17$ GeV

${}^7\text{Be}+{}^{208}\text{Pb}$




${}^9\text{Be}+{}^{208}\text{Pb}$






$\sqrt{s_{NN}}=17$ GeV, central rapidity

$^{12}\text{C}+^{208}\text{Pb}$



$^{16}\text{O}+^{208}\text{Pb}$



Summary

Nuclear structure from ultra-relativistic heavy ion collisions

Snapshots of the ground-state wave function

Spatial correlations in the ground state → harmonic flow


Possible data (NA61@SPS) in conjunction with a detailed knowledge of the evolution of the fireball would allow to place constraints on the α -cluster structure of the colliding nuclei.

Conversely, the knowledge of the clustered nuclear distributions helps to verify the fireball evolution models

Additional slides

Phenomenon of flow

Quark-gluon plasma is formed!




“Initial shape – final flow” transmutation detectable in the asymmetry of the momentum distribution of detected particles – follows from collectivity

Shape-flow transmutation

The eccentricity parameters are transformed (in all models based on collective dynamics) into asymmetry of the transverse-momentum flow.


Linear response:



[Božek 3+1 viscous hydro + THERMINATOR]

Our modeling


Generate nucleon positions with Monte Carlo, parameters (size of the cluster, distance between clusters) properly adjusted



All simulations with GLISSANDO 2


Our modeling (^{12}C)

Three α 's in a triangular arrangement, generate nucleon positions with Monte Carlo, parameters (size of the cluster, distance between clusters) properly adjusted



a)

dashed: data from charge form factor \rightarrow BEC,
solid - our BEC,
dotted - Jastrow (Buendia et al.)




b)

points - variational MC (Wiringa et al.),
line - our VMC


All simulations with GLISSANDO 2

Geometry of nucleus \rightarrow geometry of fireball

Triangular nucleus causes triangular "damage"!




intrinsic density of ^{12}C




geometry of the fireball

$v_{s_{NN}}=17$ GeV

$^{12}\text{C}+^{208}\text{Pb}$



$^{16}\text{O}+^{208}\text{Pb}$



$^2\text{H}+^{208}\text{Pb}$ @ $v_{s_{NN}}=17$ GeV

