

# Final state measurements in $t\bar{t}$ events and comparison to MC generators

L. Mijović

SM@LHC, Firenze, Italy, 21-Apr-2015



Discuss **ATLAS and CMS Run1 final state measurements in  $t\bar{t}$  events** for the following observables:

- **jet veto,**
- **jet multiplicities and  $p_T$ -spectra,**
- **$t\bar{t}$ +heavy flavour production.**


Comment also on the **MC generator modelling** of these quantities;

- $t\bar{t}$  MCs need to provide simulated samples suitable for:
  - precision  $t\bar{t}$  measurements,
  - measurements where  $t\bar{t}$  is an important **background** ( $t\bar{t}H$ , SUSY & other NP searches).
- $\Rightarrow$  good modelling of hard-process and additional jets related observables is required (frequently with the same sample).
- Stress **NLO+PS generators** and **newer generators** (e.g. C++ showers) expected to take over from the legacy ones during Run 2.


- $t\bar{t}$  dilepton events with 2  $b$ -jets
- additional jets ( $\sim$  not from Born-level ME)
- measure fraction of events with no additional jets with  $p_T > Q_0$

$$f_{\text{gap}}(Q_0) = n_{\text{gap}}(Q_0)/N_{t\bar{t}}$$

- High precision: many uncertainties cancel in the ratio.




b-jets  
leptons  
additional jet



- Probes the modelling of the additional jets; too large  $f_{\text{gap}} =$  too little activity.

ATLAS: EPJ C72 (2012) 2043, CMS: EPJ C74 (2014) 3014, CMS-PAS-TOP-12-041.





## CMS-PAS-TOP-12-041


- NLO+PS modelling ok figure:  $|\eta| < 0.8$  region that shows some data-MC difference (agreement better for inclusive  $|\eta|$ )
- CMS uncertainties small enough for data to prefer some NLO setups over other?
- Beware of low  $Q_0$  regions ( $Q_0 \ll m_t$ ). Soft interactions (e.g. hard parton-beam remnant interactions) can be of  $\sim O(5\%)$  (Phys Rev D 86 114038).

Trends observed to be correlated with other observables ( $n(\text{jets})$ , jet spectra,  $H_T$ ).

- Rivet implementation of ATLAS: EPJ C72 (2012) 2043
- Rivet = Robust Independent Validation of Experiment and Theory  
<http://rivet.hepforge.org/>
- ATL-PHYS-PUB-2015-002, ATL-PHYS-PUB-2014-005,  
ATL-PHYS-PUB-2015-007: new generator commissioning and tuning.
- With judicious choice of setups: data-MC agreement.



Measure  $n(\text{jets})$  distributions in  $t\bar{t}$  events, CMS: [CMS-PAS-TOP-12-041](#), EPJ C74 (2014) 3014, ATLAS: JHEP 01 (2015) 020.




Several theoretical approaches:

- **NLO+PS**, **(N)LO multi-leg+PS**, **LO+PS**.
- best to use in Run1? Depends on use-case, e.g.:
  - inclusive  $\sigma_{t\bar{t}}$ : sensitive to  $n(\text{jets})$  when using e.g.  $H_T$  or light jet  $b$ -tagging fake rates ,
  - SUSY** search in final state with high  $n(\text{jets})$ , where  $t\bar{t}$  is a large background,
  - high- $p_T$  NP signal simulation: which generator parameters to use?


Large  $n(\text{jets})$ : **multi-leg** best suited, **NLO+PS** not expected to work out of the box.

Example use-case: inclusive  $\sigma_{t\bar{t}}$ : sensitive to  $n(\text{jets})$  when using e.g.  $H_T$  or light jet  $b$ -tagging fake rates. High  $n(\text{jets})$ : NLO+PS MCs not expected to yield precise predictions with default setups. (CMS-PAS-TOP-12-041, ATL-PHYS-PUB-2015-002)



- MC@NLO+fortran HERWIG trend consistent with  $fgap$ .
- With judicious choice of MC setups: data-MC agreement can be achieved, also for newer generators (multi-leg NLO+PS provide improved predictions).

Measure jet  $p_T$ ,  $\eta$ , jet-jet  $\Delta R$  distributions in  $t\bar{t}$  events, CMS-PAS-TOP-12-041, ATLAS: JHEP01 (2015) 020.





ATLAS:

- single lepton decay channel,  $\geq 1$   $b$ -jet + extra  $p_T$ -cuts on the two leading jets,
- 1<sup>st</sup>-4<sup>th</sup> jet  $\sim$  proxy for hard process modelling (top,  $t\bar{t}$   $p_T$ ),
- 5<sup>th</sup> jet  $\sim$  proxy for additional jet modelling,
- Rivet implementation available.

CMS:

- additional jets kinematics in dilepton decay channel events with  $\geq 2$   $b$ -jets,
- probing additional jets modelling in the same phase-space as jet veto, but using different observables.

- ATLAS: JHEP01 (2015) 020, 7 TeV,  $l+l$  jets channel
- CMS: CMS-PAS-TOP-12-041, 8 TeV, dilepton channel



- ATLAS: MC@NLO+fortran HERWIG trend consistent with  $f_{\text{gap}}$ .
- With judicious choice of MC setups: data-MC agreement can be achieved, also for newer generators.

## $t\bar{t}+b\bar{b}$ production:

- presence of two very different scales:  $m_t$  and jet  $p_T$ .
- Thus large ren./fac. scale uncertainties from theory.
- Irreducible background to  $t\bar{t}H(H \rightarrow b\bar{b})$ .
- Measurements: test theory prediction and reduce  $t\bar{t}H$  measurement uncertainties.

$t\bar{t} + jj$  and  $t\bar{t} + c\bar{c}$ : **source of reducible background due to  $b$ -jet mis-tagging.**

## Experimental measurements:

- so far in  $t\bar{t}$  dilepton decay channel
- ATLAS: Phys. Rev. D 89, 072012 (2014):  
 $\sigma(t\bar{t}+HF \text{ jet})/\sigma(t\bar{t}+\text{light jet})$ , where HF= $b, c$
- CMS: CMS-PAS-TOP-12-024 (7 TeV), arXiv:1411.5621 (8 TeV):
  - ratio:  $\sigma(t\bar{t}+b\bar{b})/\sigma(t\bar{t}+jj)$ ,
  - as well as  $\sigma(t\bar{t}+b\bar{b})$ ,  $\sigma(t\bar{t}+jj)$  values.



CMS-PAS-TOP-12-024 (7 TeV), [arXiv:1411.5621](https://arxiv.org/abs/1411.5621) (8 TeV)

Results reported in two different regions of the phase space:

- **Visible (aka fiducial) phase space:**
  - leptons, jets from  $t\bar{t}+b\bar{b}$ : required to be within experimentally accessible region;
  - $p_T > 20$  GeV,  $|\eta| < 2.4$  (2.5 for jets).
- **Full phase space:**
  - purpose: comparison to parton-level theoretical calculations.
  - ( Though NLO+PS predictions now also available. )
  - Perform jet reconstruction on partons (gluons, quarks lighter than top) and  $\tau \rightarrow$  hadrons.
  - extra jet  $p_T > 20$  GeV and  $p_T > 40$  GeV results provided.

Measurement strategy:

- **combined secondary vertex (CSV)  $b$ -tagging algorithm:**
  - (SV+lifetime info of tracks)  $\rightarrow b$ -tagging discriminator
- **data fit of CSV discriminator of 3rd and 4th highest CSV jets to obtain  $\sigma(t\bar{t}+b\bar{b})/\sigma(t\bar{t}+jj)$  and the absolute normalization.**



Phase Space (PS)	$\sigma_{t\bar{t}b\bar{b}}$ [pb]	$\sigma_{t\bar{t}jj}$ [pb]	$\sigma_{t\bar{t}b\bar{b}}/\sigma_{t\bar{t}jj}$
Visible PS (particle)			
Jet $p_T > 20$ GeV/c	$0.029 \pm 0.003 \pm 0.008$	$1.28 \pm 0.03 \pm 0.15$	$0.022 \pm 0.003 \pm 0.005$
Full PS (parton)			
Jet $p_T > 20$ GeV/c	$1.11 \pm 0.11 \pm 0.31$	$52.1 \pm 1.0 \pm 6.8$	$0.021 \pm 0.003 \pm 0.005$
Jet $p_T > 40$ GeV/c	$0.36 \pm 0.08 \pm 0.10$	$16.1 \pm 0.7 \pm 2.1$	$0.022 \pm 0.004 \pm 0.005$
NLO calculation			
Jet $p_T > 40$ GeV/c	$0.23 \pm 0.05$	$21.0 \pm 2.9$	$0.011 \pm 0.003$

Dominant uncertainties:  $b$ -tagging efficiency and MC modelling.

ATLAS and CMS have performed a number of final state measurements in  $t\bar{t}$  events during Run1.

- These offer tests of pQCD modelling (and beyond, e.g. jet veto),
- and enable us to improve the performance of the MC generators.
- **Rivet implementation** of many measurements available and used for tuning and Run 2 generators commissioning.

MC Modelling:



- **good modelling in MC-s: jet veto, jet multiplicities and  $p_T$ -spectra;**
  - achieved with measurements and subsequent tuning of the sensitive parameters.
- For some distributions (e.g. top  $p_T$ ) modelling issues more tricky;
  - in L. Scodellaro's talk.

$t\bar{t}$ +HF measurements: test state-of-the-art theory.



- In the future we will hopefully be able to reduce measurement uncertainties (currently  $\sim 30\%$ ),
- and provide **Rivet implementation** of the fiducial measurements.

# Extra



- **Rivet implementation** of ATLAS: JHEP 01 (2015) 020
- **ATL-PHYS-PUB-2015-002, ATL-PHYS-PUB-2015-007**: new generator commissioning and tuning
- **Using ME and PS handles**: can reach data-MC agreement in *newer generators*.



- **Rivet implementation** of ATLAS: JHEP 01 (2015) 020
- **ATL-PHYS-PUB-2015-002, ATL-PHYS-PUB-2015-007**: new generator commissioning and tuning
- **Using ME and PS handles**: can reach data-MC agreement in *newer generators*.



- **Rivet implementation** of ATLAS: JHEP 01 (2015) 020
- **ATL-PHYS-PUB-2015-002, ATL-PHYS-PUB-2015-007**: new generator commissioning and tuning
- **Using ME and PS handles**: can reach data-MC agreement in *newer generators*.



arXiv:1411.5621

**truth-level objects (fiducial measurement):**

- leptons:  $pt > 20$ ,  $\eta < 2.4$ . From W from top decays.  $\tau \rightarrow e, \mu$  = included
- jets:  $pt > 20$ ,  $\eta < 2.5$ ,  $dR(j,j) > 0.5$ , remove jet if  $dR(j,l) < 0.5$ . From all final state particles, excluding  $\nu$ , antikt, 0.5.
- truth b-jet: B-hadron = one of the constituents. c-jets: similar with lower priority

**ttbb,cc,LF definition:**

- ttbb(cc): 2 extra jets contain B(C)-hadron in the constituents
- ttbj: only 1 extra jets contains B-hadron in the constituents
- c-tagging done only if b-tagging fails
- ttLF: both jets from light flavour or light+c
- tt+other: the background contributions from single top production and from tt events that fail the visible phase-space cuts.

**Fiducial phase-space: all charged particles (but not  $\nu$ ) from ttjj in the fiducial region defined above.**

arXiv:1411.5621

**measurement phase-space:**

- leptons:  $pt > 20$ ,  $\eta < 2.4$  (same as truth-level)
- leptons: relative isolation (not at truth-level)
- jets:  $pt > 30$ ,  $\eta < 2.5$  (truth-level:  $pt > 20$ )
- b-tagging: CSV algorithm,  $\epsilon_b \sim 45\%$   $\epsilon_{light} \sim 0.1\%$
- extra event-level cuts (not at truth-level): os leptons,  $m_{ll} > 12$  GeV  
ee,emu:  $|m_{ll} - m_Z| < 15$  GeV, ee,mumu:  $MET > 30$  GeV


arXiv:1411.5621

**full phase-space measurement:**

- jet reconstruction is performed on the partons (gluons, as well as quarks lighter than top) before hadronization +  $\tau \rightarrow$  hadrons
- showering done with Pythia6; thus parton 4-vectors are after all boosts and rotations from PS and hadronization
- jet reconstruction algorithm: same as for the fiducial measurement
- b jets:  $dR=0.5$  truth-matching of parton-jets to b-quarks
- results are quoted for two different jet  $p_T$  thresholds of  $p_T > 20$  and  $> 40$  GeV for extra jets (not from top).

arXiv:1411.5621

Normalized distribution of the b jet discriminator:



fit:

- simultaneously to  $ttb\bar{b}$  and  $ttj\bar{j}$  CSV discriminator distribution
- contains two free parameters, an overall normalization and the ratio of the number of  $ttb\bar{b}$  events to  $ttj\bar{j}$  events.
- $ttcc$  and  $ttLF$  contributions are combined, and the ratio of the  $ttb\bar{b}$  to  $ttb_j$  contributions is constrained using the predictions from the MC simulation.

arXiv:1411.5621

Source	$\sigma_{t\bar{t}b\bar{b}}$ (%)	$\sigma_{t\bar{t}jj}$ (%)	$\frac{\sigma_{t\bar{t}b\bar{b}}}{\sigma_{t\bar{t}jj}}$ (%)
Pileup	1.0	1.0	1.0
JES & JER	11	8.0	5.0
b tag (b quark flavour)	15	<0.1	15
b tag (c quark flavour)	4.0	<0.1	4.0
b tag (light flavour)	7.0	<0.1	7.0
Ratio of $t\bar{t}b\bar{b}$ and $t\bar{t}bj$	9.0	<0.1	9.0
Bkgnd modelling	1.0	1.0	1.0
$t\bar{t}c\bar{c}$ fraction in the fit	4.2	0.2	4.0
Lepton identification	4.0	4.0	—
MC generator	3.0	3.0	3.0
$Q^2$ scale	8.0	3.0	6.0
PS matching	12	5.0	3.0
PDF	4.0	4.0	<0.1
Eff. ( $t\bar{t}c\bar{c}$ fraction)	—	1.6	1.6
Luminosity	2.6	2.6	—
Total uncertainty	28	12	22