



Contribution ID: 113

Type: not specified

Neutral pion form factor measurement by the NA62 experiment

Monday, 29 August 2016 19:10 (20 minutes)

The NA62 experiment at CERN collected a large sample of charged kaon decays with a highly efficient trigger for decays into electrons in 2007. The kaon beam represents a source of tagged neutral pion decays in vacuum. A measurement of the electromagnetic transition form factor slope of the neutral pion in the time-like region from ~ 1 million fully reconstructed π^0 Dalitz decay is presented. The limits on dark photon production in π^0 decays from the earlier kaon experiment at CERN, NA48/2, are also reported.

Summary

The NA62 experiment at CERN collected a large sample of charged kaon decays with a highly efficient trigger for decays into electrons in 2007. The kaon beam represents a source of tagged neutral pion decays in vacuum. A measurement of the electromagnetic transition form factor slope of the neutral pion in the time-like region from ~ 1 million fully reconstructed π^0 Dalitz decay is presented. The limits on dark photon production in π^0 decays from the earlier kaon experiment at CERN, NA48/2, are also reported.

Primary author: PEPE, Monica (INFN Perugia (IT))

Presenter: PEPE, Monica (INFN Perugia (IT))

Session Classification: Section B

Track Classification: Section B: Light Quarks