## XIIth Quark Confinement and the Hadron Spectrum



Contribution ID: 308

Type: not specified

## Overview of experimental results on collective flow with identified particles at RHIC and the LHC

Monday, 29 August 2016 18:00 (30 minutes)

Anisotropic flow studies play a crucial role in improving our understanding of the behaviour and the nature of matter created in collisions of heavy ions. The different flow harmonics  $(v_n)$  harmonics for identified particles can be used to constrain the initial conditions and the value of shear viscosity over entropy density ratio. These studies allow also to reveal the role of the hadronic rescattering phase in the development of flow. In this talk I review the results from measurements of elliptic  $(v_2)$ , triangular  $(v_3)$ , quantrangular  $(v_4)$  and pentagonal  $(v_5)$  of identified particles from the RHIC and LHC heavy-ion physics programs.

## **Summary**

Anisotropic flow studies play a crucial role in improving our understanding of the behaviour and the nature of matter created in collisions of heavy ions. The different flow harmonics  $(v_n)$  harmonics for identified particles can be used to constrain the initial conditions and the value of shear viscosity over entropy density ratio. These studies allow also to reveal the role of the hadronic rescattering phase in the development of flow. In this talk I review the results from measurements of elliptic  $(v_2)$ , triangular  $(v_3)$ , quantrangular  $(v_4)$  and pentagonal  $(v_5)$  of identified particles from the RHIC and LHC heavy-ion physics programs.

Primary author: CHRISTAKOGLOU, Panos (Nikhef National institute for subatomic physics (NL))Presenter: CHRISTAKOGLOU, Panos (Nikhef National institute for subatomic physics (NL))Session Classification: Section D

Track Classification: Section D: Deconfinement