



Contribution ID: 68

Type: not specified

Approximate degeneracy of heavy-light mesons with the same L

Thursday 1 September 2016 15:30 (20 minutes)

In this work, we point out that there exists an approximate degeneracy among heavy-light systems with the same L .

This is supported by an experimental fact which can be seen from the observed data. This approximate symmetry explains why the GI model obtains results similar to those of the heavy-light systems which are fitted well with experimental data. This is because the GI model has this symmetry from the beginning which is broken by the spin-orbit interactions. Numerical results of the GI model together with those of other models respecting heavy quark symmetry are compared with the experimental data of the D mesons and they well give similar results to each other.

We analytically show that expectation values of $[H_0, \vec{L}^2]$ give us at most of the order of $1/m_Q$ for 0^- and 1^- states and the similar arguments will give us the same conclusion for other higher states in our model which respects heavy quark symmetry. Note that this order of magnitude, $1/m_Q$, is the same as those which break degeneracy of a spin doublet of heavy-light systems. It is shown that there is a rotational symmetry in the limit of $m_Q \rightarrow \infty$ and nonrelativistic limit of heavy-quark symmetry.

Simple application of our idea to other states can be given by baryons QQq like Ξ_{cc}^+ , multi-quark states in which one light quark is included like $QQ\bar{Q}q$, and probably other states in which a couple of light quarks can be regarded as a brown quark. A good example is given by a spectrum of Λ_c which gives us $\Lambda_c(2286)$ with $L = 0$, $\Lambda_c^+(2595)$ and $\Lambda_c^+(2625)$ with $L = 1$, and $\Lambda_c^+(2880)$ and $\Lambda_c^+(2940)$ with $L = 2$, where a spin multiplet is given by member/members with the same L . L is defined by an angular momentum between a heavy quark c and two light quarks (ud). One can easily see that gaps between different spin multiplets are nearly equal to $\Lambda_{QCD} \sim 300$ MeV, which coincides with the observation of heavy-light mesons.

This property may be explained by adopting a string picture of a heavy-light system, i.e., having a fixed end and a free end and connecting both ends with a string.

Summary

Careful observation of the experimental spectra of heavy-light mesons tells us that heavy-light mesons with the same angular momentum L are almost degenerate. The estimate is given how much this degeneracy is broken in our relativistic potential model, and it is analytically shown that expectation values of a commutator between the lowest order Hamiltonian and \vec{L}^2 are of the order of $1/m_Q$ with a heavy quark mass m_Q . It turns out that nonrelativistic approximation of heavy quark system has a rotational symmetry and hence degeneracy among states with the same L . This property may be explained by adopting a string picture of a heavy-light system, i.e., having a fixed end and a free end and connecting both ends with a string. This feature can be tested by measuring higher orbitally and radially excited heavy-light meson spectra for $D/D_s/B/B_s$ in LHCb and forthcoming BelleII.

Author: MATSUKI, Takayuki (Tokyo Kasei University)

Co-authors: Dr L^U, Qi-Fang (Institute of High Energy Physics, CAS, Beijing 100049, People's Republic of China); Dr MORI, Toshiyuki (Faculty of Human Development, Kobe University, 3-11 Tsurukabuto, Nada, Kobe 657-8501, Japan); Dr DONG, Yubing (Institute of High Energy Physics, CAS, Beijing 100049, People's Republic of China)

Presenter: MATSUKI, Takayuki (Tokyo Kasei University)

Session Classification: Section C

Track Classification: Section C: Heavy Quarks