



Contribution ID: 387

Type: **not specified**

## H-dibaryon in Holographic QCD

*Tuesday 30 August 2016 21:20 (20 minutes)*

We investigate the H-dibaryon ( $uuddss$ ) in holographic QCD [1, 2]. Holographic QCD is derived from a QCD-equivalent D-brane system in the superstring theory via the gauge/gravity correspondence. In holographic QCD, all baryons appear as topological chiral solitons of Nambu-Goldstone bosons and (axial) vector mesons [1, 2]. In this framework, the H-dibaryon can be described as an  $SO(3)$ -type hedgehog state [3]. In this paper, we present the formalism of the H-dibaryon in holographic QCD, and investigate its properties.

[1] T. Sakai and S. Sugimoto, Prog. Theor. Phys. 113 (2005) 843; 114 (2005) 1083.

[2] K. Nawa, H. Suganuma and T. Kojo, Phys. Rev. D75 (2007) 086003.

[3] A.P. Balachandran et al., Phys. Rev. Lett. 52 (1984) 887.

### Summary

We investigate the H-dibaryon ( $uuddss$ ) in holographic QCD, which is derived from a QCD-equivalent D-brane system via the gauge/gravity correspondence.

**Primary author:** SUGANUMA, Hideo (Kyoto University)

**Co-authors:** MATSUMOTO, Kohei (Kyoto University); NAKAGAWA, Yuya (Kyoto University)

**Presenter:** SUGANUMA, Hideo (Kyoto University)

**Session Classification:** Poster Session and Wine Tasting

**Track Classification:** Poster session