Irregularities at chemical freeze-out of hadrons as an evidence of quark-gluon plasma formation

V.V. Sagun

V.V. Sagun

K.A. Bugaev, A.I. Ivanytskyi, D.R. Oliinychenko

Thessaloniki, August 28 - September 4, 2016

Irregularities at chemical freeze-out of hadrons as an evider

Hadron resonance gas model (HRGM)

- Basic assumption thermal/chemical equilibrium \Rightarrow parameters: T, $\mu_{\rm B}$, $\mu_{\rm I3}$ P. Braun-Munzinger et al., Phys. Lett. B 344, 43, (1995) J. Cleymans et al., Z. Phys. C 74, 319 (1997)
- HRGM accounts for all hadrons from PDG tables with masses up to 3.2 GeV K.A. Bugaev et al., Eur. Phys. J. A 49, 30 (2013)
- Hadronic gas mixture with multicomponent hard-core repulsion ⇒ equation of state of the Van der Wals type Corrections:
 - Strange particle nonequilibrium
 - Decaying of resonances
 - Width correction

Fit parameters: $T, \mu_B, \mu_{I_3}, \gamma_s$

 R_{pions} , R_{kaons} , R_{mesons} , $R_{baryons}$, R_{lambda} - fixed hard-core radii. μ_S - is found from the net zero strangeness condition. K. A. Bugaev et al., EPJ A 49, 30-1-8 (2013); K. A. Bugaev et al., EPL 104, 22002, p.1 - 6 (2013);

1/11

Strangeness Horn and Λ Horn With new radii and γ_s fit



Total fit of 121 hadron ratios is the best of existing!

$$\chi^2/dof = 63.978/65 \approx 0.98$$

Jump of ChFO Pressure at AGS Energies

• Temperature T_{CFO} as a function of collision energy \sqrt{s} is rather non smooth



• Significant jump of pressure ($\simeq 6$ times) and energy density ($\simeq 5$ times)



K.A. Bugaev et al., Phys. Part. Nucl. Lett. 12(2015) [arXiv:1405.3575]; Ukr. J. Phys. 60 (2015) [arXiv:1312.4367]

Jump of Entropy Density and Trace Anomaly Peak



4/11

Baryonic density as functions of collision energy at CFO



Sharp peak of the baryonic charge density at $\sqrt{s_{NN}} = 4.9$ GeV

Irregularities in hadron production





In 1982 J. Rafelski and B. Müller predicted that enhancement of strangeness production is a signal of deconfinement

total to thermal particle yields asymmetry indicating a significant role of the particle decays at CFO

freeze-out

Phys. Rev. Lett. 48(1982)

Narrow collision energy range $\sqrt{s_{NN}} = 4.3-4.9$ GeV

Conclusions

- With our HRGM the high quality fit is achieved for 121 hadron ratios measured at 14 values of the center of mass energy $\sqrt{s_{NN}}$ at the AGS, SPS and RHIC with the accuracy $\chi^2/dof = 63.978/65 \simeq 0.98;$
- high quality description of the CFO data allowed us to find few novel irregularities in the collision energy range $\sqrt{s_{NN}} = 4.3 \cdot 4.9$ GeV (pressure, energy density jumps and $\frac{\Lambda}{n}$ yield ratio);
- in addition, we found a sharp peak of the trace anomaly $\delta = \frac{e-3p}{T^4}$ and baryonic charge density at $\sqrt{s_{NN}} = 4.9 \text{ GeV}$;
- we found two sets of strongly correlated quasi-plateaus in entropy per baryon and the pion number (thermal and total) per baryon at CFO which were predicted based on shock adiabat model as a signal of the quark-gluon-hadron mixed phase formation;

• these irregularities are also accompanied by the total to thermal particle yields asymmetry, i.e. $\frac{K_{tot}^+ - K_{th}^+}{K_{tot}^+ + K_{th}^+}, \frac{A_{tot} - A_{th}}{A_{tot} + A_{th}}, \frac{\Xi_{tot} - \Xi_{th}}{\Xi_{tot} + \Xi_{th}}, \text{ indicating a significant role of the particle decays at chemical freeze-out:}$

• we conclude that a dramatic change in the system properties seen in the narrow collision energy range $\sqrt{s_{NN}} = 4.3 - 4.9$ GeV opens entirely new possibilities for experimental studies on FAIR and NICA.



Hadron Resonance Gas Model

One component gas:
$$p = p^{id.gas} \cdot \exp\left(-\frac{pV}{T}\right)$$

Multicomponent case: $p = \sum_{i} p_{i} = \sum_{i} T \phi_{i} \exp\left[\frac{\mu_{i}-2\sum_{j} p_{j}V_{ji}+\sum_{j} p_{j}V_{ji}p_{i}/p}{T}\right]$
All hadrons are in full chemical equilibrium

The number of particles of *i*-th sort:

$$N_i = \phi_i(T, m_i, g_i) e^{\frac{\mu_i}{T}} \equiv \frac{g_i V}{(2\pi)^3} \int exp\left(\frac{-\sqrt{k^2 + m_i^2 + \mu_i}}{T}\right) d^3k$$

hard-core repulsion of the Van der Waals type

$$\mu_i = \mu_B B_i + \mu_S S_i + \mu_{I_3} I_{3_i}, \ i = 1..s$$

 g_i - degeneracy factor ϕ_i -thermal particle density $V_{ij} = \frac{2\pi}{3}(R_i + R_j)^3$ - excluded volume



Zinovjev G. M., Eur. Phys. J. A 49 (2013) 30-1-8.

Hadron Resonance Gas Model corrections

Strangeness corrections:

$$\phi_i(T) \rightarrow \phi_i(T) \gamma_s^{s_i}$$

 s_i — number of strange valence quarks and anti-quarks.

J. Rafelski, Phys. Lett. B 62, 333 (1991);

• Resonance decay:

$$n^{fin}(X) = \sum_{Y} BR(Y \to X) n^{th}(Y),$$

where $BR(X \rightarrow X) = 1$, BR=BRANCHING RATIO (taken from PDG); • Width correction:

$$\int exp\left(\frac{-\sqrt{k^2+m_i^2}}{T}\right) d^3k \rightarrow \frac{\int_{M_0}^{\infty} \frac{dx_i}{(x-m_i)^2+\Gamma^2/4} \int exp\left(\frac{-\sqrt{k^2+x^2}}{T}\right) d^3k}{\int_{M_0}^{\infty} \frac{dx_i}{(x-m_i)^2+\Gamma^2/4}},$$

Breit-Wigner distribution having a threshold M_0 ,

m - resonance mass, Γ - resonance width.

Ratios:

 $R_{ij} = \frac{N_i}{N_j} = \frac{
ho_i}{
ho_j} \implies \text{volume is excluded}$

Fit parameters: T, μ_B , μ_{I_3} , γ_s R_{pions} , R_{kaons} , R_{mesons} , $R_{baryons}$, R_{lambda} - fixed hard-core radii. μ_S - is found from the net zero strangeness condition. K. A. Bugaev et al., EPJ A 49, 30-1-8 (2013); K. A. Bugaev et al., EPL 104, 22002, p.1 - 6 (2013) 10/11

Correlated Quasi-Plateaus

Mixed phase has anomalous thermodynamic properties => plateau in collision energy dependence of entropy per baryon!

K.A. Bugaev, M.I. Gorenstein, B. Kampher, V.I. Zhdanov, Phys. Rev. D 40, 9, (1989) K.A. Bugaev, M.I. Gorenstein, D.H. Rischke, Phys. Lett. B 255, 1, 18 (1991)

Since the main part of the system entropy is defined by thermal pions => thermal pions/baryon should have a plateau too!

Also the total number of pions per baryons should have a (quasi)plateau!

Entropy per baryon has wide plateaus due to large errors

Quasi-plateau in total pions per baryon ?

Thermal pions demonstrate 2 plateaus



11/11