Quark Matter 2015 - XXV International Conference on Ultrarelativistic Nucleus-Nucleus Collisions



Contribution ID: 719

Type: Poster

## QGP properties from azimuthal-angular dependence of charged-pion interferometry

Tuesday 29 September 2015 16:30 (2 hours)

Recently the results of HBT measurements of charged pions with respect to the second and third order event plane are presented by PHENIX [1]. They extract  $\epsilon_2$  and  $\epsilon_3$  from the HBT measurements which contain information about not only the source shape at freezeout but also the space-time evolution of QGP matter. They show the relation between initial  $\epsilon_{2,3}$  which are obtained from a Glauber model and final  $\epsilon_{2,3}$  which are extracted from the HBT radii. They find that the final  $\epsilon_2$  from the HBT radii is finite and smaller than the initial  $\epsilon_2$ . On the other hand, the final  $\epsilon_3$  is significantly reduced and potential reversed by the end in spite of existence of finite initial  $\epsilon_3$ . The interesting different response of  $\epsilon_2$  and  $\epsilon_3$  during space-time evolution gives us a clue to understand the detailed QGP properties.

For analyses of such high statistics experimental results, we develop a state of the art numerical scheme of causal viscous hydrodynamics for relativistic heavy ion collisions, which has a shock-wave capturing scheme and less numerical dissipation [2]. Furthermore, using the hydrodynamic algorithm, we construct a hybrid model of hydrodynamic model plus UrQMD to include the realistic freezeout processes. Using the model we investigate the time evolution of spatial anisotropies  $\epsilon_n$ . We find that the sign of  $\epsilon_3$  changes from positive to negative during the space-time evolution, which suggests a solution of the vanishing final  $\epsilon_3$  from the HBT radii by PHENIX. From detailed analyses, we discuss the initial conditions of hydrodynamic model and the detailed QGP properties such as transport coefficients.

[1] A.Adare et al. [PHENIX collaboration], Phys. Rev. Lett. 222301 (2014).

[2] Y. Akamatsu, S. Inutsuka, C. Nonaka, M. Takamoto, J. Comp. Phys. (2014), pp. 34-54; K.Okamoto, Y.Akamatsu, C.Nonaka, in preparation.

## On behalf of collaboration:

NONE

Primary author: NONAKA, Chiho (Nagoya University)

Co-authors: OKAMOTO, Kazuhisa (Nagoya University); AKAMATSU, Yukinao (Nagoya University)

Presenter: NONAKA, Chiho (Nagoya University)

Session Classification: Poster Session

Track Classification: Collective Dynamics