



Contribution ID: 458

Type: **Contributed talk**

Results from (anti-)(hyper-)nuclei production and searches for exotic bound states with ALICE at the LHC

Monday, 28 September 2015 12:35 (20 minutes)

The high collision energies reached at the LHC enable significant production rates of light (anti-)(hyper-)nuclei in proton-proton, proton-lead and, in particular, lead-lead collisions. The excellent particle identification capabilities of the ALICE apparatus, based on the specific energy loss in the time projection chamber and the velocity information from the time-of-flight detector, allow for the detection of these (anti-)nuclei.

Furthermore, the high tracking resolution provided by the inner tracking system enables the separation of primary nuclei from those coming from the decay of heavier systems.

This allows for the reconstruction of decays such as the hypertriton mesonic weak decay (${}^3_{\Lambda}\text{H} \rightarrow {}^3\text{He} + \pi^{-}$), the decay of a hypothetical bound state of a Λ with a neutron into a deuteron and pion or the H-dibaryon decaying into a Λ , a proton and a π^{-} .

Results on the production of stable nuclei and anti-nuclei in Pb-Pb and lighter collision systems will be presented. Hypernuclei production rates in Pb-Pb will also be shown, together with upper limits estimated on the production of hypothetical exotica candidates. The results will be compared with predictions for the production in thermal (statistical) and coalescence models.

On behalf of collaboration:

ALICE

Primary author: SHARMA, Natasha (University of Tennessee (US))

Presenter: SHARMA, Natasha (University of Tennessee (US))

Session Classification: Open Heavy Flavors and Strangeness I

Track Classification: Open Heavy Flavors and Strangeness