



Contribution ID: 202

Type: **Contributed talk**

## Heavy-light flavor correlations and the QCD phase boundary

*Monday 28 September 2015 18:40 (20 minutes)*

Modifications in magnitude of fluctuations for different observables are an excellent probe of a phase transition or its remnant.

In heavy-ion collision, fluctuations related to conserved charges carried by light and strange quarks play an important role to identify the QCD chiral crossover and deconfinement properties.

Recent Lattice QCD simulations have revealed that the charmed mesons are deconfined together with light-flavor mesons in the temperature range where the chiral symmetry is partially restored. This result strongly suggests that the light-flavor dynamics interferes non-trivially with the heavy flavors.

We discuss correlations between the light and heavy-light flavored mesons at finite temperature within a chiral effective theory implementing heavy quark symmetry. We show that the heavy quark dynamics is tied to the light flavor physics, and the thermodynamics is strongly dragged by the chiral crossover dominated by the non-strange flavors. Consequently, the fluctuations carried by the states with strangeness can be used to characterize the onset of the chiral symmetry restoration.

References:

- [1] C. Sasaki, Phys. Rev. D 90, no. 11, 114007 (2014).
- [2] C. Sasaki and K. Redlich, Phys. Rev. D 91, no. 7, 074021 (2015).

### **On behalf of collaboration:**

NONE

**Author:** Dr SASAKI, Chihiro (FIAS & Uni of Wroclaw)

**Co-author:** Prof. REDLICH, Krzysztof (University of Wroclaw)

**Presenter:** Dr SASAKI, Chihiro (FIAS & Uni of Wroclaw)

**Session Classification:** Correlations and Fluctuations II

**Track Classification:** Correlations and Fluctuations