Quark Matter 2015 - XXV International Conference on Ultrarelativistic Nucleus-Nucleus Collisions



Contribution ID: 362

Type: Poster

Bottomonia suppression in 2.76 TeV Pb-Pb collisions

Tuesday, 29 September 2015 16:30 (2 hours)

We compute the QGP suppression of $\Upsilon(1s)$, $\Upsilon(2s)$, and $\Upsilon(3s)$, χ_{b1} and χ_{b2} states in $\sqrt{s_{NN}} = 2.76$ TeV Pb-Pb collisions. Using the suppression of each of these states, we estimate the inclusive R_{AA} for the $\Upsilon(1s)$ and $\Upsilon(2s)$ states as a function of N_{part} , y, and p_T including the effect of excited state feed down. We find that our model provides a reasonable description of recently obtained CMS results for the N_{part} , y-, and p_T -dependence of R_{AA} for both the $\Upsilon(1s)$ and $\Upsilon(2s)$. Comparing to our previous results, we find a flatter rapidity dependence, thereby reducing some of the tension between our model and ALICE forward-rapidity results for $\Upsilon(1s)$ suppression.

On behalf of collaboration:

NONE

Primary author: STRICKLAND, Michael (Kent State University)

Co-authors: Mr KROUPPA, Brandon (Kent State University); Dr RYBLEWSKI, Radoslaw (Institute of Nuclear Physics PAN)

Presenter: STRICKLAND, Michael (Kent State University)

Session Classification: Poster Session

Track Classification: Quarkonia