



Contribution ID: 500

Type: **Contributed talk**

Measurement of two-particle pseudorapidity correlations in proton-lead and lead-lead collisions with the ATLAS detector

Tuesday, 29 September 2015 15:20 (20 minutes)

Two-particle pseudorapidity correlations are measured using charged particles from $\sqrt{s_{NN}}=2.76$ TeV Pb+Pb collisions by the ATLAS experiment at the LHC. The correlation function $C_N(\eta_1, \eta_2)$ is measured for different centrality intervals for $|\eta_1, \eta_2| < 2.4$ and transverse momentum $p_T > 0.5$ GeV. An enhancement is observed along $\eta_1 - \eta_2 \approx 0$ and a suppression is observed at large $\eta_1 - \eta_2$ values. The correlation function is expanded in Legendre polynomials, and root-mean-square values of the Legendre coefficients at the single particle level are measured. Significant values are observed for a_n , which decrease quickly for larger n . The leading coefficient a_1 is compared to that estimated from a fit to $C_N(\eta_1)$ for different η_1 slices, as well as to the asymmetry of the number of participating nucleons between the two colliding nuclei $A_{Npart} = (N_{partF} - N_{partB}) / (N_{partF} + N_{partB})$. The centrality dependence of a_1 shows a very similar shape as A_{Npart} in mid-central collisions (20-50%), but shows faster increase in more central and more peripheral collisions. The latter behavior suggests additional forward-backward multiplicity fluctuations that may arise from fluctuations at the sub-nucleonic level. The status of a similar measurement in proton-lead collisions will be reported. The implications of these measurements for constraining the early time dynamics of high-energy nuclear collisions are discussed.

On behalf of collaboration:

ATLAS

Primary author: JIA, Jianguo (State University of New York (US))

Presenter: JIA, Jianguo (State University of New York (US))

Session Classification: Correlations and Fluctuations IV

Track Classification: Correlations and Fluctuations