



Contribution ID: 218

Type: **Poster**

Systematic study of the parton energy loss from $p(d, {}^3\text{He})+A$ to A+A collisions using high p_T hadrons measured by the PHENIX experiment at RHIC

Tuesday 29 September 2015 16:30 (2 hours)

The energy loss of the hard scattered partons in the hot dense medium produced in relativistic heavy ion collisions has been of interest for investigating the properties of the medium.

Recently, the second-order azimuthal anisotropy of particle emission has been observed at a low p_T (1-3 GeV/c for the RHIC case) in long-range two-particle correlations in small systems such as $d+\text{Au}$ collisions at RHIC and $p+\text{Pb}$ collisions at LHC. At higher p_T , on the other hand, the strength of the anisotropy tends to become weaker, which is qualitatively similar to what was observed in A+A collisions. In A+A collisions, the anisotropy at higher p_T has been understood as the consequence of the path-length dependent energy loss of hard scattered partons. It is of a great interest whether or not the similar energy loss phenomenon is observed in the small systems ($p+A$, $d+A$).

We present a systematic study of the azimuthal anisotropy of particle emission from $p(d, {}^3\text{He})+A$ to A+A collisions using high p_T hadrons measured by the PHENIX experiment at RHIC, and discuss the systematics of the parton energy loss across the systems.

On behalf of collaboration:

PHENIX

Primary author: Dr SAKAGUCHI, Takao (BNL)

Presenter: Dr SAKAGUCHI, Takao (BNL)

Session Classification: Poster Session

Track Classification: QGP in Small Systems