

Contribution ID: 436

Type: poster

J/ψ production in U+U collisions at the STAR experiment

Quark-gluon plasma (QGP), a novel state of deconfined nuclear matter, has been extensively studied in highenergy heavy-ion collisions

at the Relativistic Heavy Ion Collider (RHIC). Suppression of heavy quarkonium production (e.g. J/ψ , Υ) in heavy-ion

collisions compared to proton-proton collisions due to the color screeening of the quark-antiquark potential is expected

to be an indicator of the partonic phase. However, there are also other effects that may influence the expected suppression

pattern of heavy quarkonia (e.g. secondary production in the QGP, cold-nuclear-matter effects).

To understand those different contributions we need to study production of heavy quarkonia in various colliding systems.

At RHIC, effects of the hot medium on heavy quarkonia have been studied in Au+Au and U+U collisions. Since U nuclei are larger than Au nuclei and are non-spherical, it is expected that in the most

central U+U collisions the energy density of the created medium is higher than in Au+Au collisions.

Thus they allow for further testing of the color screening hypothesis.

In this poster we will present preliminary results on nuclear modification factor of J/ψ production reconstructed at midrapidity via di-electron

decay channel in minimium-bias U+U collisions at $\sqrt{s_{\rm NN}} = 193$ GeV at the STAR experiment and current status of analysis of J/ψ

production in central U+U collisions.

Author: FODOROVA, Jana (Czech Technical University in Prague)

Presenter: FODOROVA, Jana (Czech Technical University in Prague)

Track Classification: Heavy Ion Physics