

MMHT2014 PDFs - Published sets and Updates

Robert Thorne

July 24th, 2015



University College London

In collaboration with Lucian Harland-Lang, Patrick Motylinski and Alan Martin



and thanks to Ben Watt, Graeme Watt and James Stirling

EPS 2015 – July 2015

Updates in PDF Fits – the **MMHT 2014** PDFs.

I will present results on the update in PDFs within the general **MSTW** framework due to some theory improvements and a variety of new data sets, including most of the up-to-date **LHC** data. The release of a new set of **MMHT** PDFs (<http://arxiv.org/abs/arXiv:1412.3989> – now accepted for publication in **EPJ C**) is summarised, and some subsequent results and future plans are discussed.

MMHT 2014 PDFs




Available in **LHAPDF5** and **LHAPDF6**.

Also at <http://www.hep.ucl.ac.uk/mmht> where there is stand-alone Fortran code, a C++ wrapper and Mathematica implementations as well as grids in **LHAPDF5** and **LHAPDF6** format.

MMHT2014 – Changes in theoretical treatment or procedures.

Continue to use extended parameterisation with Chebyshev polynomials, and freedom in deuteron nuclear corrections, as in (Eur.Phys.J. C73 (2013) 2318) – change in $u_V - d_V$ distribution.



Result for fitted deuteron correction, 4 parameters left free. Good comparison to CJ12mid model (Accardi). Uncertainty of about 0.5–1%. Feeds into PDF uncertainty.

Now use “optimal” GM-VFNS choice (Phys.Rev. D86 (2012) 074017) which is smoother near to heavy flavour transition points (more so at NLO).

Errors multiplicative not additive. Using χ^2 definition

$$\chi^2 = \sum_{i=1}^{N_{pts}} \left(\frac{D_i + \sum_{k=1}^{N_{corr}} r_k \sigma_{k,i}^{corr} - T_i}{\sigma_i^{uncorr}} \right)^2 + \sum_{k=1}^{N_{corr}} r_k^2,$$

where $\sigma_{k,i}^{corr} = \beta_{k,i}^{corr} T_i$ and $\beta_{k,i}^{corr}$ are the percentage error. Additive would use $\sigma_{k,i}^{corr} = \beta_{k,i}^{corr} D_i$. Previously did this for all but normalisation uncertainty.

Strange branching ratio. Now avoid those determined by fits to dimuon data relying on PDF input. Also apply error which feeds into PDFs. Use $B_\mu = 0.092 \pm 10\%$ from [hep-ex/9708014](#). Fits prefer $B_\mu = 0.085 - 0.091 \pm 15\%$.

Have been using [de Florian, Sassot](#) nuclear corrections. Update to more recent version, [de Florian, Sassot, Stratmann, Zurita, Phys.Rev. D85 \(2012\) 074028](#).

Changes in data sets.

Replacement of HERA run I neutral and charged current data from HERA and ZEUS with combined data set with full treatment of correlated errors. Fit to data very good. Slightly better fit at NNLO.



Inclusion of HERA combined data on $F_2^c(x, Q^2)$. Fit quality $\sim 60-80$ for 52 points.

Inclusion of all direct published HERA $F_L(x, Q^2)$ measurements. Undershoot data a little at lower Q^2 , but χ^2 not much more than one per point.

No inclusion of separate run II H1 and ZEUS data yet. Wait for Run II combination. (See later).

Inclusion of the CDF W -asymmetry data, the D0 electron asymmetry data $p_T > 25\text{GeV}$ based on 0.75 fb^{-1} and new D0 muon asymmetry data for $p_T > 25\text{GeV}$ based on 7.3 fb^{-1} .

Fit quality at NLO and NNLO for $F_L(x, Q^2)$ data



LHC data on W,Z


Now using APPLGrid – MCFM and DYNNLO/FEWZ include the ATLAS W,Z rapidity data directly in the fit. → slight change is in the strange quark.

$W^+ - W^-$ asymmetry no longer an issue at all both for ATLAS and CMS asymmetry data. Slightly better at NLO.

Include LHCb data on W^+, W^- , and $Z \rightarrow e^+e^-$. Both predicted/fit well at NLO.

Include CMS data on $Z \rightarrow e^+e^-$, and ATLAS high mass Drell-Yan data. Again both predicted/fit well.



Fit CMS double differential Drell Yan data extending to low mass. NNLO fits enormously better than NLO at lowest mass $\sim 20 - 45\text{GeV}$.



W asymmetry data now fit very well, though a little better at NLO than at NNLO.

CMS Drell Yan data.

Fit very poor at **NLO** in lowest mass bins (where it is effectively **LO**), even when data highly weighted.





Enormously improved fit quality at **NNLO** due to improvement in cross-sections.

Sensitivity to strange fraction in quarks, but differs at **NLO** and **NNLO** and weak compared to direct constraint from di-muon data.

LHC data on $t\bar{t}$

Include data on $\sigma_{t\bar{t}}$ from **Tevatron** (combined cross section measurement from **D0** and **CDF**), and all published data from **ATLAS** and **CMS** for **7TeV** and one point at **8TeV**. Use $m_t = 172.5$ GeV (value used in **Tevatron** combination) with an error of **1 GeV**, with χ^2 penalty applied. Predictions and fit good, with **NLO** preferring masses slightly below $m_t = 172.5$ GeV and **NNLO** masses slightly above.



LHC data on jets

At **NLO** include **CMS** data and **ATLAS 7 TeV + 2.76 TeV** data. **ATLAS** $\chi^2 = 107/116$ and **CMS** $\chi^2 = 143/133$ before included directly.

Enormous project of full **NNLO** calculation (**Gehrmann-de-Ridder et al.**) nearing completion. Some indications of full form.

As default at **NNLO** still fit **Tevatron** data which are relatively near to threshold. However, omit **LHC** data. Investigate inclusion of **K**-factor.

data set	N_{pts}	MMSTWW	MMHT14 (no LHC)	MMHT14 (with LHC)
NLO				
ATLAS (2.76+7 TeV)	116	107	107	106
CMS (7 TeV)	133	140	143	138
NNLO small				
ATLAS (2.76+7 TeV)	116	(107)	(123)	(122) 115
CMS (7 TeV)	133	(142)	(137)	(138) 137
NNLO large K-factor				
ATLAS (2.76+7 TeV)	116	(117)	(132)	(132) 126
CMS (7 TeV)	133	(145)	(137)	(139) 139

MMHT2014 PDFs compared to MSTW2008 PDFs.


Use same “dynamic tolerance” prescription to determine eigenvectors.

Typical tolerance $T = \Delta\chi^2 \sim 10$.

We now have 25 eigenvector pairs, rather than the 20 in MSTW or even the 23 in MMSTWW.





Eigenvector sets made available for $\alpha_S(M_Z^2) = 0.135$ (LO), $\alpha_S(M_Z^2) = 0.118, 0.120$ (NLO) and $\alpha_S(M_Z^2) = 0.118$ (NNLO)

Comparison of PDFs at NLO



Change in NLO PDFs from all updates, including LHC data updates.

Comparison of PDFs at NNLO




Change in NNLO PDFs from all updates. Gluon uncertainty at high- x slightly greater than at NLO. At NNLO final extracted $\alpha_S(M_Z^2) = 0.11722$, but PDFs presented for $\alpha_S(M_Z^2) = 0.1180$.

$\alpha_S(m_Z^2)$ and PDF sets.

$\alpha_S(m_Z^2)$ coming out similar to 2008 fit. Still a NLO/NNLO difference. Both fairly compatible with global average, i.e.

NLO – $\alpha_S(m_Z^2) = 0.1201$, NNLO – $\alpha_S(m_Z^2) = 0.1172$.



$\alpha_S(m_Z^2)^{\text{world}} = 0.1186 \pm 0.0006$. Decide to present MMHT2014 PDFs with eigenvectors at round value of $\alpha_S(m_Z^2) = 0.118$ at NNLO and at NLO also at $\alpha_S(m_Z^2) = 0.120$.





Make central sets available for $\alpha_S(m_Z^2) = 0.108 - 0.128$ in steps of 0.001.

Individual Data Sets – Most constraining Data Sets.

NLO




NNLO




Also constraining lower limit – NLO

NuTeV νN $x F_3$: $\chi_{n,0}^2 = 36$ for 42 pts.




SLAC μd F_2 : $\chi_{n,0}^2 = 29.4$ for 38 pts.




Also constraining upper limit – NNLO

BCDMS μp F_2 : $\chi_{n,0}^2 = 173$ for 163 pts.





SLAC ep F_2 : $\chi_{n,0}^2 = 30.4$ for 37 pts.



Obtain $\alpha_S(m_Z^2)^{\text{NLO}} = 0.1201 \pm 0.0015$ and $\alpha_S(m_Z^2)^{\text{NNLO}} = 0.1172 \pm 0.0013$.


New data sets for fit – $W + c$ differential distributions.

	GeV	data	MSTW2008	MMHT2014
$\sigma(W + c)$	$p_T^{\text{lep}} > 25$	$107.7 \pm 3.3(\text{stat.}) \pm 6.9(\text{sys.})$	102.8 ± 1.7	110.2 ± 8.1
$\sigma(W + c)$	$p_T^{\text{lep}} > 35$	$84.1 \pm 2.0(\text{stat.}) \pm 4.9(\text{sys.})$	80.4 ± 1.4	86.5 ± 6.5
R_c^\pm	$p_T^{\text{lep}} > 25$	$0.954 \pm 0.025(\text{stat.}) \pm 0.004(\text{sys.})$	0.937 ± 0.029	0.924 ± 0.026
R_c^\pm	$p_T^{\text{lep}} > 35$	$0.938 \pm 0.019(\text{stat.}) \pm 0.006(\text{sys.})$	0.932 ± 0.030	0.904 ± 0.027



MSTW2008 a bit low (especially for **ATLAS**), but **MMHT2014** seems fine particularly for **CMS** (shown). Data will add some constraint.

New data on high rapidity W production **LHCb** at **7 TeV**.



Generally perfectly good agreement using **NNLO**.

PDFs and Heavy Quarks

As before we will make the same PDFs sets (i.e. exactly the same input at $Q_0^2 = 1 \text{ GeV}^2$) available for three flavour and four flavour fixed-flavour number schemes (FFNS).

As default will also fix the number of flavours in α_S , but will probably also provide analogous sets with variable flavour α_S as there were some requests for MSTW2008.

Will also make available sets with fits done for m_c and m_b (defined in pole scheme) varying from default values of $m_c = 1.40 \text{ GeV}$ and $m_b = 4.75 \text{ GeV}$ in steps of 0.05 GeV and 0.25 GeV respectively.

Probably not as wide a range as last time – $m_c = 1.05 - 1.75 \text{ GeV}$ and $m_b = 4.00 - 5.50 \text{ GeV}$.

m_b constrained to fairly close to $m_b = 4.75 \text{ GeV}$ from direct $F_2^{\bar{b}b}(x, Q^2)$ data from HERA and m_c also constrained far better than previous range from various sources.

Dependence on m_c at NLO in fits at fixed $\alpha_s(M_Z^2) = 0.120$.

m_c (GeV)	χ_{global}^2 2996 pts	$\chi_{F_2^c}^2$ 52 pts	$\alpha_s(M_Z^2)$
1.15	3242	76	0.120
1.2	3239	74	0.120
1.25	3240	72	0.120
1.3	3245	71	0.120
1.35	3254	71	0.120
1.4	3267	71	0.120
1.45	3283	73	0.120
1.5	3303	76	0.120
1.55	3327	80	0.120

Similar variation with m_c for varying $\alpha_s(M_Z^2) = 0.120$. For $0.13 \text{ GeV} < m_c < 1.5 \text{ GeV}$ difference compared to free coupling negligible.

Preference for $m_c \sim 1.20 \text{ GeV}$, or marginally higher.

Slight tension between global fit and charm data.

Dependence on m_c at NNLO in fits at fixed $\alpha_s(M_Z^2) = 0.118$.

m_c (GeV)	χ_{global}^2 2663 pts	$\chi_{F_2^c}^2$ 52 pts	$\alpha_s(M_Z^2)$
1.15	2712	79	0.118
1.2	2706	77	0.118
1.25	2705	76	0.118
1.3	2706	77	0.118
1.35	2711	79	0.118
1.4	2720	83	0.118
1.45	2731	88	0.118
1.5	2750	96	0.118
1.55	2770	106	0.118

Similar variation with m_c for varying $\alpha_s(M_Z^2)$.

Less tension between global fit and charm.

Again preference for $m_c \sim 1.25\text{GeV}$.

HERA II Combined data

Recently released in [arXiv:1506.06042](https://arxiv.org/abs/1506.06042).

Using $Q_{\min}^2 = 2\text{GeV}^2$ then there are 1185 data points with 162 correlated systematics, 7 procedural uncertainties and luminosity uncertainty.

Separated into 7 subsets, depending on whether e^+ or e^- , neutral or charged current and on E_p .

Compared to 621 data points, separated into 5 subsets, with generally larger uncertainties from HERA I (but fewer systematics) combined data used previously.

Prediction with MMHT2014 PDFs already fairly good.

NLO – $\chi^2 = 1602.1/1185 = 1.35$ per point

NNLO – $\chi^2 = 1497.5/1185 = 1.26$ per point

(HERAPDF2.0 get ~ 1.20 with $Q_{\min}^2 = 2 \text{ GeV}^2$ at NLO and NNLO).

Under refitting in global fit

NLO – $\chi^2 = 1533.8/1185 = 1.29$ per point, with deterioration $\Delta\chi^2 = 20$ in other data.

NNLO – $\chi^2 = 1457.7/1185 = 1.23$ per point, with deterioration $\Delta\chi^2 = 8$ in other data.


Also trying fitting only **HERA II** data, with 4 parameters fixed to avoid particularly unusual PDFs.

NLO – $\chi^2 = 1416.0/1185 = 1.19$ per point

NNLO – $\chi^2 = 1380.6/1185 = 1.17$ per point


NNLO definitely better than **NLO**.

Charged current χ^2 over 20 units better in **HERA II** only fit, and over 10 units better at **NNLO**.



Look at **NLO** compared to **NNLO** with different Q_{\min}^2 without refitting.

NNLO clearly superior, but less obvious in fit to only **HERA II** data.




Also look at effect of changing the Q^2 cut, on only HERA II data, at both NLO and NNLO (note – definition of χ^2 for HERAPDF2.0 not identical).





Improvement in χ^2 with Q_{\min}^2 largely achieved without refitting.

Less improvement than for HERAPDF2.0 particularly in global fit and at NNLO.

Quite large fluctuations in theory/data at low Q^2 rather than obvious systematic issue.



HERA II modified PDFs very well within MMHT2014 uncertainties. PDFs from HERA II data only fit in some ways similar to HERAPDF2.0. Predictions for e.g. $gg \rightarrow H$ change by $< 0.2\%$ for full range of LHC energies.



Uncertainties (preliminary) quite similar to **MMHT2014**.

Most obvious improvement in gluon for $x \sim 0.001$.

Conclusions

MMHT2014 PDFs recently released. Improvement in parameterisation, heavy flavour treatments, nuclear corrections, and branching ratio for dimuon data.

Inclusion of up-to-date **HERA**, **Tevatron** and relevant published **LHC** data. The fit is always good (except low mass **Drell-Yan** at **NLO**).

Few dramatic effects on PDFs. In general predictions remain very close to those with **MSTW2008** PDFs.

25 eigenvector sets available with $\alpha_S(M_Z^2) = 0.135$ at **LO**, with $\alpha_S(M_Z^2) = 0.118, 120$ at **NLO** and with $\alpha_S(M_Z^2) = 0.118$ at **NNLO**.



$\alpha_S(m_Z^2)^{\text{NLO}} = 0.1201 \pm 0.0015$ and $\alpha_S(m_Z^2)^{\text{NNLO}} = 0.1172 \pm 0.0013$ from study of $\alpha_S(M_Z^2)$ dependence. Sets for $0.108 - 0.128$ produced at **NLO** and **NNLO**.

Sets in different flavour schemes and different quark masses very soon.

New **HERA II** combined data studied. Fit quality good – better at **NNLO**. No very significant changes in PDFs or predictions. Effect of Q_{\min}^2 in data not obviously sign of problems at low x, Q^2 .

Back -up

Fit quality at NLO and NNLO for D0 asymmetry data



MMHT2014 PDFs compared to MSTW2008 PDFs.

Use same “dynamic tolerance” prescription to determine eigenvectors.

Typical tolerance $T = \Delta\chi^2 \sim 10$.

We now have **25** eigenvector pairs, rather than the **20** in **MSTW** or even the **23** in **MMSTWW**.

Eigenvector sets made available for $\alpha_S(M_Z^2) = 0.135$ (**LO**), $\alpha_S(M_Z^2) = 0.118, 0.120$ (**NLO**) and $\alpha_S(M_Z^2) = 0.118$ (**NNLO**)

In addition the central sets are available at

LO $\alpha_S(M_Z^2) = 0.134, 0.135, 0.136$

NLO $\alpha_S(M_Z^2) = 0.117, 0.118, 0.119, 0.120, 0.121$

NNLO $\alpha_S(M_Z^2) = 0.117, 0.118, 0.119$

This allows the **PDF** + α_S uncertainty to be calculated, if using the prescription of adding in quadrature.

NLO

HERA structure – 6 eigenvector directions.

fixed target data DIS data – 13 eigenvector directions

LHC data – 4 eigenvector directions

Tevatron data – 9 eigenvector directions

Dimuon data – 8 eigenvector directions

E866 Drell Yan data – 10 eigenvector directions.

NNLO

HERA structure – 11 eigenvector directions.

fixed target data DIS data – 10 eigenvector directions

LHC data – 8 eigenvector directions


Tevatron data – 6 eigenvector directions

Dimuon data – 9 eigenvector directions


E866 Drell Yan data – 6 eigenvector directions.

data set	N_{pts}	MMSTWW	MMHT14 (no LHC)	MMHT14 (LHC)
NLO				
ATLAS W^+, W^-, Z	30	47	44	38
CMS W asym $p_T > 35$ GeV	11	9	16	7
CMS asym $p_T > 25, 30$ GeV	24	9	17	8
LHCb $Z \rightarrow e^+e^-$	9	13	13	13
LHCb W asym $p_T > 20$ GeV	10	12	14	12
CMS $Z \rightarrow e^+e^-$	35	21	22	19
ATLAS high-mass Drell-Yan	13	20	20	21
CMS double diff. Drell-Yan	132	385	396	372
NNLO				
ATLAS W^+, W^-, Z	30	72	53	39
CMS W asym $p_T > 35$ GeV	11	18	15	8
CMS asym $p_T > 25, 30$ GeV	24	18	17	9
LHCb $Z \rightarrow e^+e^-$	9	23	22	21
LHCb W asym $p_T > 20$ GeV	10	24	21	18
CMS $Z \rightarrow e^+e^-$	35	30	24	22
ATLAS high-mass Drell-Yan	13	18	16	17
CMS double diff. Drell Yan	132	159	151	150


Fit quality at NLO and NNLO for ATLAS W, Z rapidity data




Fit quality at NLO and NNLO for LHCb W, Z rapidity data



Fit to CMS inclusive jet data.





Fit to CMS inclusive jet data at NNLO with large K -factor.





Individual Data Sets

HERA data in global fit prefers higher $\alpha_S(M_Z^2)$.


NLO



NLO



Global χ^2 depends on m_t but minimises at very similar $\alpha_S(M_Z^2)$ for a rather wide range.



However, fit quality to $\sigma_{t\bar{t}}$ data alone very sensitive to m_t and $\alpha_S(M_Z^2)$ interplay.



Values determined by free best fit using $m_t = 172.5 \text{ GeV} \pm 1 \text{ GeV}$ are $m_t(\text{NLO, NNLO}) = 171.7, 174.2 \text{ GeV}$, as opposed to world average of $m_t = 173.34 \pm 0.76 \text{ GeV}$.

Be conservative on $\alpha_S(M_Z^2)$ constraints direct from $\sigma_{t\bar{t}}$, but similar constraints from other sets.

New data sets for fit – $t\bar{t}$ differential distributions.

Variety of data sets not in PDF determination as they did not meet cut-off date and/or missing NNLO corrections.


For example, differential $t\bar{t}$ production (show CMS below). $y_{t\bar{t}}$ distribution at NLO very good, p_t distribution off in shape ($m_{t\bar{t}}$ somewhere in between).



Interesting to see NNLO corrections.


Differential Data

As it improves differential top production data will help constrain the gluon.



However, here potentially inclusion of **NNLO** is very important as available approximation using threshold resummation (Guzzi, Lipka, Moch) implies. Softer PDF currently preferred at **NLO**, contrary to requirement of inclusive cross-section, may be misleading.

New data sets for fit – $W + c$ differential distributions.




Also fine for normalised distribution.

Only at **NLO**, but possibly less sensitivity to higher order in normalised distribution.


New data on high rapidity W production LHCb at 7 TeV.

The same data is also available as a ratio or asymmetry, with a reduction in systematic uncertainty (particularly luminosity).



Again clearly good agreement. PDF uncertainties still a little smaller than data constraints – except maybe highest rapidity (high- x PDF).

Comparison of PDFs at NNLO when LHC jet data included



NNLO PDFs with LHC jet data included in fit with both K -factor choices compared to standard NNLO PDFs.

NNLO total cross sections

	MSTW08	MMHT14 no LHC	MMHT14
W Tevatron (1.96 TeV)	$2.746^{+0.049}_{-0.042}$	2.803	$2.782^{+0.056}_{-0.056}$
Z Tevatron (1.96 TeV)	$0.2507^{+0.0048}_{-0.0041}$	0.2574	$0.2559^{+0.0052}_{-0.0046}$
W^+ LHC (7 TeV)	$6.159^{+0.111}_{-0.099}$	6.214	$6.197^{+0.103}_{-0.092}$
W^- LHC (7 TeV)	$4.310^{+0.078}_{-0.069}$	4.355	$4.306^{+0.067}_{-0.076}$
Z LHC (7 TeV)	$0.9586^{+0.020}_{-0.014}$	0.9695	$0.9638^{+0.014}_{-0.013}$
W^+ LHC (14 TeV)	$12.39^{+0.22}_{-0.21}$	12.49	$12.48^{+0.22}_{-0.18}$
W^- LHC (14 TeV)	$9.33^{+0.16}_{-0.16}$	9.39	$9.32^{+0.15}_{-0.14}$
Z LHC (14 TeV)	$2.051^{+0.035}_{-0.033}$	2.069	$2.065^{+0.035}_{-0.030}$
Higgs Tevatron	$0.853^{+0.028}_{-0.029}$	0.877	$0.874^{+0.024}_{-0.030}$
Higgs LHC (7 TeV)	$14.40^{+0.17}_{-0.23}$	14.54	$14.56^{+0.21}_{-0.29}$
Higgs LHC (14 TeV)	$47.50^{+0.47}_{-0.74}$	47.61	$47.69^{+0.63}_{-0.88}$
$t\bar{t}$ Tevatron	$7.19^{+0.17}_{-0.12}$	7.54	$7.51^{+0.21}_{-0.20}$
$t\bar{t}$ LHC (7 TeV)	$171.1^{+4.7}_{-4.8}$	176.5	$175.9^{+3.9}_{-5.5}$
$t\bar{t}$ LHC (14 TeV)	953.3^{+16}_{-18}	969.0	969.9^{+16}_{-20}

Few changes greater than one sigma (PDF uncertainty only).

Dependence on m_c (pole mass) at NLO in fits.

m_c (GeV)	χ_{global}^2 2996 pts	$\chi_{F_2^c}^2$ 52 pts	$\alpha_s(M_Z^2)$
1.15	3239	75	0.1190
1.2	3237	73	0.1192
1.25	3239	71	0.1194
1.3	3245	70	0.1195
1.35	3254	70	0.1196
1.4	3268	71	0.1198
1.45	3283	73	0.1200
1.5	3303	76	0.1201
1.55	3327	81	0.1202

Some correlation between m_c and $\alpha_s(M_Z^2)$.

Preference for $m_c \sim 1.20\text{GeV}$.

NMC data prefer lower m_c – quicker threshold evolution respectively.

Slight tension between global fit and charm data.


Dependence on m_c at NNLO in m_c fits.

m_c (GeV)	χ_{global}^2 2663 pts	$\chi_{F_2^c}^2$ 52 pts	$\alpha_s(M_Z^2)$
1.15	2703	78	0.1164
1.2	2699	76	0.1166
1.25	2698	75	0.1167
1.3	2701	76	0.1169
1.35	2707	78	0.1171
1.4	2717	82	0.1172
1.45	2729	88	0.1173
1.5	2749	96	0.1173
1.55	2769	105	0.1175

Slightly less correlation between m_c and $\alpha_s(M_Z^2)$.


Less variation in fit quality and much less tension.

Preference for $m_c \sim 1.25\text{GeV}$.



Higher luminosity LHCb $Z \rightarrow \mu^+ \mu^-$ data.

Dijets



Using reweighting exercise for **CMS** dijets results in a rather modified shape of gluon.

Not as high rapidity as other sets – dependence on renormalisation/factorisation scales not so severe.


Reflection of different shape of higher order corrections?


Different conclusions for fits to **D0** and **ATLAS** dijet data, though they are not not necessarily incompatible.

Similar to changes required by **LHC** inclusive jet data.

Different range of rapidity spanned. Need to use scale other than p_T to get good fits. $\mu = 2p_T$ best for **ATLAS** and $\mu = M_{JJ}$ best for **D0**.

For **ATLAS** rapidity dependent scale choices give results more like that for **CMS**, but with a worse fit and lower value of N_{eff} .





At high rapidity calculations unstable for scales equal to relatively low multiples of p_T .