


New Physics at *Belle II*

Tobias Hurth

Johannes Gutenberg University Mainz



Conveners:

Florian Bernlochner (Bonn)
Ryosuke Itoh (KEK)
Vittorio Lubicz (Rome)
Jernej Kamenik (IJS Ljubljana)
Emi Kou (LAL-Orsay)
Ulrich Nierste (KIT)
Luca Silvestrini (Rome)

LHCb anomalies and Superiso

Local Organizing Committee:

Monika Blanke	Thomas Deppisch
Pablo Goldenzweig	Martin Heck
Thomas Kuhr	Ulrich Nierste
Andreas Pargner	Stefan Schacht
Martina Schorn	Paul Tremper

Belle II Theory Interface Platform — Working Group 9

February 23rd-25th 2015

Karlsruhe Institute of Technology (KIT) — Germany

<https://indico.cern.ch/event/357770/>

blueyonder



Plan of the Talk

- Some remarks about Superiso (Nazila Mahmoudi)
- Tension with SM in semileptonic exclusive penguin decays in LHCb data
- Signs for lepton non universality
- Correlations between charged lepton and neutrino modes

Superiso

Nazila Mahmoudi

SuperIso

- public C program
- dedicated to the **flavour physics** observable calculations
- based on the most precise calculations publicly available
- various models implemented
- interfaced to several spectrum calculators
- modular program with a well-defined structure
- complete reference manual available (~160 pages)
- latest release: v3.4 (Oct. 2014)

<http://superiso.in2p3.fr>

F. Mahmoudi, Comput. Phys. Commun. 178 (2008) 745
F. Mahmoudi, Comput. Phys. Commun. 180 (2009) 1579
F. Mahmoudi, Comput. Phys. Commun. 180 (2009) 1718

Models

- Standard Model
- General Two Higgs Doublet Model, and types I-IV
Interface available with *2HDMC*
- Supersymmetry: MSSM, NMSSM, BMSSM
Interfaces available with *Softsusy*, *Isajet*, *Spheno*, *Suspect* and *NMSSMTools*
- Generic models through Flavour Les Houches Accord (FLHA) files

Observables

- **Radiative penguin decays**

- inclusive branching ratios of $B \rightarrow X_{s,d}\gamma$ (NNLO)
- isospin asymmetry of $B \rightarrow K^*\gamma$

- **Electroweak penguin decays**

- branching ratios of $B_{s,d} \rightarrow \ell^+\ell^-$ (NNLO QCD + NLO EW)
- observables related to $B \rightarrow X_s\ell^+\ell^-$: BR, A_{FB} , zero crossing $q_0(A_{\text{FB}})$
- observables related to $B \rightarrow K^*\ell^+\ell^-$: BR, A_{FB} , $q_0(A_{\text{FB}})$, F_L , A_T^1 , $P_i^{(\prime)}$, ...
- observables related to $B \rightarrow K\ell^+\ell^-$: BR, A_{FB} , F_H , R_K , ...

- **Neutrino modes**

- branching ratio of $B \rightarrow \tau\nu$
- branching ratio of $B \rightarrow D\tau\nu$
- branching ratios of $D_{(s)} \rightarrow \ell\nu$
- branching ratio of $K \rightarrow \mu\nu$

- **Others**

- LEP and Tevatron direct search limits
- Electroweak precision tests: oblique parameters (S, T, U)
- Muon anomalous magnetic moment
- Relic density \rightarrow **SuperIso Relic**

Structure of the code

- Starting from an SLHA file automatically generated, or user-provided
- Reading of the SLHA file and filling of an internal parameter C-structure
- Calculation of the tree level observables
- Calculation of the Wilson coefficients
- Calculation of all the flavour observables
- Results displayed on screen and in FLHA output files

- Calculation of the other observables


- Possibility to use FLHA input files

Perspectives

- Addition of new observables:
 - observables related to $B_s \rightarrow \phi \ell^+ \ell^-$
 - branching ratio of $B \rightarrow D^* \tau \nu$
 - branching ratios of $B \rightarrow K^{(*)} \nu \nu$
 - CP violation observables
 - ...
- Addition of more new physics scenarios (Vector like quark models, ...)
- Towards a generic new physics model implementation
 - starting from a given new physics Lagrangian
 - automatic generation of the Wilson coefficients
 - automatic calculation of the observables

Radiative and semileptonic penguin decays

Separation of new physics and hadronic effects



Operator product expansion: Factorization of **short-** and **long-distance** physics

- Electroweak effective Hamiltonian: $H_{eff} = -\frac{4G_F}{\sqrt{2}} \sum C_i(\mu, M_{heavy}) \mathcal{O}_i(\mu)$
- $\mu^2 \approx M_{New}^2 \gg M_W^2$: 'new physics' effects: $C_i^{SM}(M_W) + C_i^{New}(M_W)$

How to compute the hadronic matrix elements $\mathcal{O}_i(\mu = m_b)$?


How to compute the hadronic matrix elements $\mathcal{O}_i(\mu = m_b)$?

Inclusive modes $B \rightarrow X_s \gamma$ or $B \rightarrow X_s l^+ l^-$

- Heavy mass expansion for inclusive modes:

$$\Gamma(\bar{B} \rightarrow X_s \gamma) \xrightarrow{m_b \rightarrow \infty} \Gamma(b \rightarrow X_s^{\text{parton}} \gamma), \quad \Delta^{\text{nonpert.}} \sim \Lambda_{QCD}^2 / m_b^2$$

No linear term Λ_{QCD}/m_b (perturbative contributions dominant)



Inclusive modes $B \rightarrow X_s \gamma$ or $B \rightarrow X_s \ell^+ \ell^-$

- Heavy mass expansion for inclusive modes:

$$\Gamma(\bar{B} \rightarrow X_s \gamma) \xrightarrow{m_b \rightarrow \infty} \Gamma(b \rightarrow X_s^{\text{parton}} \gamma), \quad \Delta^{\text{nonpert.}} \sim \Lambda_{QCD}^2 / m_b^2$$

No linear term Λ_{QCD}/m_b (perturbative contributions dominant)

An old story:

- If one goes beyond the leading operator ($\mathcal{O}_7, \mathcal{O}_9$):
breakdown of local expansion

A dedicated analysis:

naive estimate of non-local matrix elements leads to 5% uncertainty.

[Benzke, Lee, Neubert, Paz, arXiv:1003.5012](#)





Latest improvements of inclusive $\bar{B} \rightarrow X_s \ell^+ \ell^-$

Beyond existing NNLL QCD precision electromagnetic corrections

were calculated: Huber, Hurth, Lunghi, Nucl. Phys. B802(2008)40 and work in progress

Corrections to matrix elements lead to large collinear $\text{Log}(m_b/m_\ell)$

$$\delta\text{BR}(B \rightarrow X_s \mu^+ \mu^-) = \begin{cases} (+2.0\%) & \text{low } q^2 \\ (-6.8\%) & \text{high } q^2 \end{cases} \quad \delta\text{BR}(B \rightarrow X_s e^+ e^-) = \begin{cases} (+5.2\%) & \text{low } q^2 \\ (-17.6\%) & \text{high } q^2 \end{cases}$$




⋮

Until very recently:

'Latest' Babar and Belle measurements of inclusive $\mathcal{B}(b \rightarrow sll)$

Belle hep-ex/0503044 (!!!) (based $152 \times 10^6 B\bar{B}$ events)

Babar hep-ex/0404006 (!!!) (based $89 \times 10^6 B\bar{B}$ events)



Two new analyses from the B factories:

New Babar analysis on dilepton spectrum arXiv:1312.3664

New Belle analysis on AFB arXiv:1402.7134

Forthcoming theory analysis including all three independent angular observables ($z = \cos\theta$)



Huber, Hurth, Lunghi

- Observables

$$\frac{d^2\Gamma}{dq^2 dz} = \frac{3}{8} [(1 + z^2) H_T(q^2) + 2z H_A(q^2) + 2(1 - z^2) H_L(q^2)]$$

$$\frac{d\Gamma}{dq^2} = H_T(q^2) + H_L(q^2)$$

$$\frac{dA_{\text{FB}}}{dq^2} = 3/4 H_A(q^2)$$



Low- q^2 region: $1 \text{ GeV}^2 < q^2 < 6 \text{ GeV}^2$

High- q^2 region: $q^2 > 14.4 \text{ GeV}^2$

How to compute the hadronic matrix elements $\mathcal{O}(m_b)$?

Exclusive modes $B \rightarrow K^* \gamma$ or $B \rightarrow K^* \ell^+ \ell^-$

Naive approach:

Parametrize the hadronic matrix elements in terms of form factors

Exclusive modes $B \rightarrow K^* \gamma$ or $B \rightarrow K^* \ell^+ \ell^-$

QCD-improved factorization: BBNS 1999

$$\mathcal{T}_a^{(i)} = C_a^{(i)} \xi_a + \phi_B \otimes T_a^{(i)} \otimes \phi_{a,K^*} + O(\Lambda/m_b)$$

Existence of 'non-factorizable' strong interaction effects
which do *not* correspond to form factors

Exclusive modes $B \rightarrow K^* \gamma$ or $B \rightarrow K^* \ell^+ \ell^-$

QCD-improved factorization: BBNS 1999

$$T_a^{(i)} = C_a^{(i)} \xi_a + \phi_B \otimes T_a^{(i)} \otimes \phi_{a,K^*} + O(\Lambda/m_b)$$

- Separation of **perturbative hard kernels** from **process-independent nonperturbative** functions like form factors
- **Relations between formfactors** in large-energy limit Charles et al. 1998
- **Limitation: insufficient information on power-suppressed Λ/m_b terms** (breakdown of factorization: 'endpoint divergences')

Phenomenologically highly relevant issue

general strategy of LHCb to look at ratios of exclusive modes

Egede, Hurth, Matias, Ramon, Reece, arXiv:0807.2589, arXiv:1005.0571


see also Altmannshofer et al., arXiv:0811.1214; Bobeth et al., arXiv:0805.2525

Kinematics

- Assuming the \bar{K}^* to be on the mass shell, the decay $\bar{B}^0 \rightarrow \bar{K}^{*0}(\rightarrow K^- \pi^+) \ell^+ \ell^-$ described by the lepton-pair invariant mass, s , and the three angles θ_l , θ_K , ϕ .

After summing over the spins of the final particles:

$$\frac{d^4\Gamma}{dq^2 d\cos\theta_l d\cos\theta_K d\phi} = \frac{9}{32\pi} J(q^2, \theta_l, \theta_K, \phi)$$



$$\begin{aligned}
 J(q^2, \theta_l, \theta_K, \phi) = & \\
 = & J_{1s} \sin^2 \theta_K + J_{1c} \cos^2 \theta_K + (J_{2s} \sin^2 \theta_K + J_{2c} \cos^2 \theta_K) \cos 2\theta_l + J_3 \sin^2 \theta_K \sin^2 \theta_l \cos 2\phi \\
 & + J_4 \sin 2\theta_K \sin 2\theta_l \cos \phi + J_5 \sin 2\theta_K \sin \theta_l \cos \phi + (J_{6s} \sin^2 \theta_K + J_{6c} \cos^2 \theta_K) \cos \theta_l \\
 & + J_7 \sin 2\theta_K \sin \theta_l \sin \phi + J_8 \sin 2\theta_K \sin 2\theta_l \sin \phi + J_9 \sin^2 \theta_K \sin^2 \theta_l \sin 2\phi
 \end{aligned}$$


$$J_i = J_i(A_{\perp L/R}, A_{\parallel L/R}, A_{0L/R}) \quad A_{\perp, \parallel} = (H_{+1} \mp H_{-1})/\sqrt{2}, \quad A_0 = H_0$$

Careful design of theoretical clean angular observables

Egede, Hurth, Matias, Ramon, Reece, arXiv:0807.2589, arXiv:1005.0571

- Dependence of soft form factors, ξ_{\perp} and ξ_{\parallel} , to be minimized !
form factors should cancel out exactly at LO, best for all s
- unknown Λ/m_b power corrections

$$A_{\perp,\parallel,0} = A_{\perp,\parallel,0}^0 (1 + c_{\perp,\parallel,0}) \text{ vary } c_i \text{ in a range of } \pm 10\% \text{ and also of } \pm 5\%$$




The experimental errors assuming SUSY scenario (b) with large-gluino mass and positive mass insertion, is compared to the theoretical errors assuming the SM.

Careful design of theoretical clean angular observables

Egede, Hurth, Matias, Ramon, Reece, arXiv:0807.2589, arXiv:1005.0571

- Dependence of soft form factors, ξ_{\perp} and ξ_{\parallel} , to be minimized !
form factors should cancel out exactly at LO, best for all s
- unknown Λ/m_b power corrections

$$A_{\perp,\parallel,0} = A_{\perp,\parallel,0}^0 (1 + c_{\perp,\parallel,0}) \text{ vary } c_i \text{ in a range of } \pm 10\% \text{ and also of } \pm 5\%$$



Optimised basis of clean (formfactor-independent)

observables: P_i Descotes-Genon, Hurth, Matias, Virto arXiv:1303.5794

Definition of P'_5

$$n_{\parallel} = \begin{pmatrix} A_{\parallel}^L \\ A_{\parallel}^{R*} \end{pmatrix}, \quad n_{\perp} = \begin{pmatrix} A_{\perp}^L \\ -A_{\perp}^{R*} \end{pmatrix}, \quad n_0 = \begin{pmatrix} A_0^L \\ A_0^{R*} \end{pmatrix}$$

$$P_1 = \frac{|n_{\perp}|^2 - |n_{\parallel}|^2}{|n_{\perp}|^2 + |n_{\parallel}|^2} = \frac{J_3}{2J_{2s}},$$

$$P_2 = \frac{\text{Re}(n_{\perp}^{\dagger} n_{\parallel})}{|n_{\parallel}|^2 + |n_{\perp}|^2} = \beta_{\ell} \frac{J_{6s}}{8J_{2s}},$$

$$P_3 = \frac{\text{Im}(n_{\perp}^{\dagger} n_{\parallel})}{|n_{\parallel}|^2 + |n_{\perp}|^2} = -\frac{J_9}{4J_{2s}},$$

$$P_4 = \frac{\text{Re}(n_0^{\dagger} n_{\parallel})}{\sqrt{|n_{\parallel}|^2 |n_0|^2}} = \frac{\sqrt{2}J_4}{\sqrt{-J_{2c}(2J_{2s} - J_3)}},$$

$$P_5 = \frac{\text{Re}(n_0^{\dagger} n_{\perp})}{\sqrt{|n_{\perp}|^2 |n_0|^2}} = \frac{\beta_{\ell} J_5}{\sqrt{-2J_{2c}(2J_{2s} + J_3)}},$$

$$P_6 = \frac{\text{Im}(n_0^{\dagger} n_{\parallel})}{\sqrt{|n_{\parallel}|^2 |n_0|^2}} = -\frac{\beta_{\ell} J_7}{\sqrt{-2J_{2c}(2J_{2s} - J_3)}},$$

Redefinition:



$$P'_4 \equiv P_4 \sqrt{1 - P_1} = \frac{J_4}{\sqrt{-J_{2c}J_{2s}}}$$

$$P'_5 \equiv P_5 \sqrt{1 + P_1} = \frac{J_5}{2\sqrt{-J_{2c}J_{2s}}}$$

$$P'_6 \equiv P_6 \sqrt{1 - P_1} = -\frac{J_7}{2\sqrt{-J_{2c}J_{2s}}}$$

Measurements of forward-backward asymmetry in $B \rightarrow K^* \mu^+ \mu^-$

$$A_{FB}(s = m_{\mu^+ \mu^-}^2) = \frac{N_F - N_B}{N_F + N_B}$$







Excellent agreement with SM at current level of precision.



However:

Many more angular observables in $B \rightarrow K^* \mu \mu$ to be measured, more sensitive to NP than AFB. New flavour structures needed !

LHCb arXiv:1304.6325



Good agreement with SM in P'_4 , P'_6 and P'_8 ,
 but a 4.0σ deviation in the third bin in P'_5



LHCb Anomaly

a statistical fluctuation, an underestimation of Λ/m_b corrections or new physics in C_9 ?

$$C_7 \quad (B \rightarrow X_s \gamma) \quad C_{10} \quad (B \rightarrow \mu^+ \mu^-)$$

- **Power corrections:** No strict theory: $A'_i = A_i(1 + C_i)$, $|C_i| 10\%$

3% on the observable level: 4.0σ

More realistic: 10% on the observable level: 3.6σ

Dimensional estimate, some soft arguments

Assume 30% : 2.2σ

Descotes, Matias, Virto arXiv:1307.5683

- **Validity of QCDf and of perturbative description of charm loops:** $[1\text{GeV}^2, 6\text{GeV}^2]$,
but local bin is $q^2 \in [4.3, 8.63]\text{GeV}^2$

- **Issue of charm loops** Khodjamirian et al. arXiv:1006.4945

Only soft gluon (but no spectator) contributions included yet

- **Analysis of factorizable power corrections**

Descotes, Hofer, Matias, Virto, arXiv:1407.8526

$$A_2(q^2) = \frac{m_B}{m_B - m_{K^*}} [\xi_{\perp}(q^2) - \xi_{\parallel}(q^2)] + \Delta A_2^{\alpha_s}(q^2) + \Delta A_2^{\Lambda}(q^2),$$

$$A_0(q^2) = \frac{E}{m_{K^*}} \xi_{\parallel}(q^2) + \Delta A_0^{\alpha_s}(q^2) + \Delta A_0^{\Lambda}(q^2), \quad \xi_{\parallel}^{(2)}(q^2) \equiv \frac{m_{K^*}}{E} A_0(q^2)$$

Only central value of power corrections fixed

Present error (without knowing the correlations)
in the LCSR calculation of formfactors too large

Nonfactorizable power corrections still open

- **Analysis of factorizable power corrections**

Descotes, Hofer, Matias, Virto, arXiv:1407.8526

$$A_2(q^2) = \frac{m_B}{m_B - m_{K^*}} [\xi_{\perp}(q^2) - \xi_{\parallel}(q^2)] + \Delta A_2^{\alpha_s}(q^2) + \Delta A_2^{\Lambda}(q^2),$$

$$A_0(q^2) = \frac{E}{m_{K^*}} \xi_{\parallel}(q^2) + \Delta A_0^{\alpha_s}(q^2) + \Delta A_0^{\Lambda}(q^2), \quad \xi_{\parallel}^{(2)}(q^2) \equiv \frac{m_{K^*}}{E} A_0(q^2)$$

Only central value of power corrections fixed

Present error (without knowing the correlations)
in the LCSR calculation of formfactors too large

Nonfactorizable power corrections still open

- **Suggestions beyond guessing numbers**

Direct calculation of QCD formfactors including correlations

see Altmannshofer et al., arXiv:0811.1214

Methods used in an analysis of $B \rightarrow K \ell \ell$

Kjodjamirian, Mannel, Wang, arXiv:1211.0234

Kjodjamirian et al., arXiv:1006.4945

- **Analysis of factorizable power corrections**

Descotes, Hofer, Matias, Virto, arXiv:1407.8526

$$A_2(q^2) = \frac{m_B}{m_B - m_{K^*}} [\xi_{\perp}(q^2) - \xi_{\parallel}(q^2)] + \Delta A_2^{\alpha_s}(q^2) + \Delta A_2^{\Lambda}(q^2),$$

$$A_0(q^2) = \frac{E}{m_{K^*}} \xi_{\parallel}(q^2) + \Delta A_0^{\alpha_s}(q^2) + \Delta A_0^{\Lambda}(q^2), \quad \xi_{\parallel}^{(2)}(q^2) \equiv \frac{m_{K^*}}{E} A_0(q^2)$$

Only central value of power corrections fixed

Present error (without knowing the correlations)




in the LCSR calculation of formfactors too large

Nonfactorizable power corrections still open

- **Use data to estimate power corrections**




see i.e. Straub, Altmannshofer arXiv:1411.3161

- If new physics (negative C_9 and less significant nontrivial C'_9) then it is compatible with the hypothesis of Minimal Flavour Violation





MFV predictions for P_2 , P_4 , and P_5



- **Global fit to the NP contributions δC_i in the MFV (Update)**
effective theory





● Crosscheck with Inclusive mode (Update)



Exclusive observables
($B \rightarrow K \mu^+ \mu^-$)



Exclusive observables
($B \rightarrow K^* \mu^+ \mu^-$)





Inclusive observables

● **Future opportunities:** (Update)

LHCb upgrade: $5fb^{-1}$ to $50fb^{-1}$

Super-B Factory Belle-II: $50ab^{-1}$



- **New physics explanations**

- "The usual suspects, such as the MSSM, warped extra dimension scenarios, or models with partial compositeness, cannot accommodate the observed deviations"

Gauld, Goertz, Haisch arXiv:1308.1959;1310.1082

Altmannshofer, Straub arXiv:1308.1501

Coefficient	1σ	2σ	3σ
C_7^{NP}	$[-0.05, -0.01]$	$[-0.06, 0.01]$	$[-0.08, 0.03]$
C_9^{NP}	$[-1.6, -0.9]$	$[-1.8, -0.6]$	$[-2.1, -0.2]$
C_{10}^{NP}	$[-0.4, 1.0]$	$[-1.2, 2.0]$	$[-2.0, 3.0]$
$C_{7'}^{\text{NP}}$	$[-0.04, 0.02]$	$[-0.09, 0.06]$	$[-0.14, 0.10]$
$C_{9'}^{\text{NP}}$	$[-0.2, 0.8]$	$[-0.8, 1.4]$	$[-1.2, 1.8]$
$C_{10'}^{\text{NP}}$	$[-0.4, 0.4]$	$[-1.0, 0.8]$	$[-1.4, 1.2]$

Model-independent analysis Descotes,Matias,Virto arXiv:1307.5683

- 1σ solutions: Z' -models (331-models...): only change C_9

Descotes,Matias,Virto arXiv:1307.5683

Altmannshofer, Straub arXiv:1308.1501

Gauld, Goertz, Haisch arXiv:1308.1959;1310.1082

Buras,De Fazio,Girrbach arXiv:1311.6729




Altmannshofer,Gori,Pospelov,Yavin arXiv:1403.1269

- SUSY are compatible with the anomaly at the 2σ level




Overall fit at the 1σ level

Mahmoudi, Neshatpour, Virto arXiv:1401.2145

CMSSM



pMSSM



Let us wait for the new LHCb analysis based on the $3fb^{-1}$ data set !

Signs for lepton non-universality ?

$$R_K \equiv \frac{\mathcal{B}(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\mathcal{B}(B^+ \rightarrow K^+ e^+ e^-)} = 0.745_{-0.074}^{+0.090} (\text{stat}) \pm 0.036 (\text{syst}) \quad \text{LHCb; arXiv:1406.6482}$$


2.6 σ deviation from SM

Hiller, Schmaltz; Ghosh et al.; Biswas et al.;
Straub et al.; Hurth et al.; Glashow et al.
Crivellin et al.


Global fits to the $b \rightarrow sll$ data

Hurth, Mahmoudi, Neshatpour, arXiv:1410.4545





Fit results for two operators



Tensors, scalars difficult, sign for tension in C_9^{μ}



Fit results for four operators $\{C_9, C'_9, C_{10}, C'_{10}\}$ Larger new physics contributions are allowed within 1σ



$\{C_9, C'_9\}$

$\{C_9, C_{10}\}$

$\{C_9, C'_9, C_{10}, C'_{10}\}$

Best fit point: χ^2 : 52 (52)





52 (52)

51 (50)

Adding $C_{10}^{(\prime)}$ or primed operators does not improve the fit !

$$\{C_9^\mu, C_9^{\prime\mu}, C_9^e, C_9^{\prime e}\}$$

Larger new physics contributions are allowed within 1σ



$$\{C_9, C_9'\}$$

$$\{C_9^\mu, C_9^{\prime\mu}, C_9^e, C_9^{\prime e}\}$$

Best fit point: χ^2 : 52 (52)





42 (50)

Fit improved by 2.6σ

Assuming that this specific four-operator scenario is correct, the one with the two-operators is ruled out by 2.6σ

$\{C_9^\mu, C_9^e, C_{10}^\mu, C_{10}^e\}$

Larger new physics contributions are allowed within 1σ



χ^2 : $\{C_9, C_{10}\}$
52 (52)

$\{C_9^\mu, C_9^e\}$.
44 (52)

$\{C_9^\mu, C_9^e, C_{10}^\mu, C_{10}^e\}$
43 (50)

Again χ^2 favours the non-universal extension against the universal one

Correlations between $b \rightarrow sll$ and $b \rightarrow s\nu\bar{\nu}$ Buras et al., arXiv:1409.4557

Gauge-invariant effective theory:

$$\begin{aligned} Q_{Hq}^{(1)} &= i(\bar{q}_L \gamma_\mu q_L) H^\dagger D^\mu H, & Q_{ql}^{(1)} &= (\bar{q}_L \gamma_\mu q_L) (\bar{l}_L \gamma^\mu l_L), \\ Q_{Hq}^{(3)} &= i(\bar{q}_L \gamma_\mu \tau^a q_L) H^\dagger D^\mu \tau_a H, & Q_{ql}^{(3)} &= (\bar{q}_L \gamma_\mu \tau^a q_L) (\bar{l}_L \gamma^\mu \tau_a l_L), \\ Q_{Hd} &= i(\bar{d}_R \gamma_\mu d_R) H^\dagger D^\mu H, & Q_{dl} &= (\bar{d}_R \gamma_\mu d_R) (\bar{l}_L \gamma^\mu l_L), \\ Q_{de} &= (\bar{d}_R \gamma_\mu d_R) (\bar{e}_R \gamma^\mu e_R), & Q_{qe} &= (\bar{q}_L \gamma_\mu q_L) (\bar{e}_R \gamma^\mu e_R). \end{aligned}$$

Correlations between $b \rightarrow sll$ and $b \rightarrow s\nu\bar{\nu}$ Buras et al., arXiv:1409.4557

Gauge-invariant effective theory:

$$\begin{aligned}
 Q_{Hq}^{(1)} &= i(\bar{q}_L \gamma_\mu q_L) H^\dagger D^\mu H, & Q_{ql}^{(1)} &= (\bar{q}_L \gamma_\mu q_L) (\bar{l}_L \gamma^\mu l_L), \\
 Q_{Hq}^{(3)} &= i(\bar{q}_L \gamma_\mu \tau^a q_L) H^\dagger D^\mu \tau_a H, & Q_{ql}^{(3)} &= (\bar{q}_L \gamma_\mu \tau^a q_L) (\bar{l}_L \gamma^\mu \tau_a l_L), \\
 Q_{Hd} &= i(\bar{d}_R \gamma_\mu d_R) H^\dagger D^\mu H, & Q_{dl} &= (\bar{d}_R \gamma_\mu d_R) (\bar{l}_L \gamma^\mu l_L), \\
 Q_{de} &= (\bar{d}_R \gamma_\mu d_R) (\bar{e}_R \gamma^\mu e_R), & Q_{qe} &= (\bar{q}_L \gamma_\mu q_L) (\bar{e}_R \gamma^\mu e_R).
 \end{aligned}$$

Match on standard electroweak effective theory:

$$\begin{aligned}
 \mathcal{O}_9^{(l)} &= (\bar{s} \gamma_\mu P_{L(R)} b) (\bar{l} \gamma^\mu l), & \mathcal{O}_{10}^{(l)} &= (\bar{s} \gamma_\mu P_{L(R)} b) (\bar{l} \gamma^\mu \gamma_5 l). \\
 \mathcal{O}_L &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_L b) (\bar{\nu} \gamma^\mu (1 - \gamma_5) \nu), & \mathcal{O}_R &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_R b) (\bar{\nu} \gamma^\mu (1 - \gamma_5) \nu).
 \end{aligned}$$

Correlations between $b \rightarrow sll$ and $b \rightarrow s\nu\bar{\nu}$ Buras et al., arXiv:1409.4557

Gauge-invariant effective theory:


$$\begin{aligned}
 Q_{Hq}^{(1)} &= i(\bar{q}_L \gamma_\mu q_L) H^\dagger D^\mu H, & Q_{ql}^{(1)} &= (\bar{q}_L \gamma_\mu q_L)(\bar{l}_L \gamma^\mu l_L), \\
 Q_{Hq}^{(3)} &= i(\bar{q}_L \gamma_\mu \tau^a q_L) H^\dagger D^\mu \tau_a H, & Q_{ql}^{(3)} &= (\bar{q}_L \gamma_\mu \tau^a q_L)(\bar{l}_L \gamma^\mu \tau_a l_L), \\
 Q_{Hd} &= i(\bar{d}_R \gamma_\mu d_R) H^\dagger D^\mu H, & Q_{dl} &= (\bar{d}_R \gamma_\mu d_R)(\bar{l}_L \gamma^\mu l_L), \\
 Q_{de} &= (\bar{d}_R \gamma_\mu d_R)(\bar{e}_R \gamma^\mu e_R), & Q_{qe} &= (\bar{q}_L \gamma_\mu q_L)(\bar{e}_R \gamma^\mu e_R).
 \end{aligned}$$

Match on standard electroweak effective theory:

$$\begin{aligned}
 \mathcal{O}_9^{(l)} &= (\bar{s} \gamma_\mu P_{L(R)} b)(\bar{l} \gamma^\mu l), & \mathcal{O}_{10}^{(l)} &= (\bar{s} \gamma_\mu P_{L(R)} b)(\bar{l} \gamma^\mu \gamma_5 l). \\
 \mathcal{O}_L &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_L b)(\bar{\nu} \gamma^\mu (1 - \gamma_5) \nu), & \mathcal{O}_R &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_R b)(\bar{\nu} \gamma^\mu (1 - \gamma_5) \nu).
 \end{aligned}$$


$$\begin{aligned}
 C_L &= C_L^{\text{SM}} + \tilde{c}_{ql}^{(1)} - \tilde{c}_{ql}^{(3)} + \tilde{c}_Z, & C_R &= \tilde{c}_{dl} + \tilde{c}'_Z & \tilde{c}_Z &= \frac{1}{2}(\tilde{c}_{Hq}^{(1)} + \tilde{c}_{Hq}^{(3)}) \\
 C_9 &= C_9^{\text{SM}} + \tilde{c}_{qe} + \tilde{c}_{ql}^{(1)} + \tilde{c}_{ql}^{(3)} - \zeta \tilde{c}_Z, & C'_9 &= \tilde{c}_{de} + \tilde{c}_{dl} - \zeta \tilde{c}'_Z & \tilde{c}'_Z &= \frac{1}{2}\tilde{c}_{Hd} \\
 C_{10} &= C_{10}^{\text{SM}} + \tilde{c}_{qe} - \tilde{c}_{ql}^{(1)} - \tilde{c}_{ql}^{(3)} + \tilde{c}_Z, & C'_{10} &= \tilde{c}_{de} - \tilde{c}_{dl} + \tilde{c}'_Z & \zeta &= 1 - 4s_w^2 \approx 0.08
 \end{aligned}$$

No model-independent correlations via gauge-invariant operators



Separate variation:
 up to 30% deviation
 from SM possible

Impact of the Wilson coefficients of the SM-EFT on $B \rightarrow K^{(*)} \nu \bar{\nu}$ vs. $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, assuming MFV and LFU, varied within their 2σ ranges allowed by the global fit to $b \rightarrow s \mu \mu$ data. Blue: \tilde{c}_Z , green: $\tilde{c}_{ql}^{(1)}$, red: $\tilde{c}_{ql}^{(3)}$. The dashed green line shows the case $\tilde{c}_{ql}^{(1)} = -\tilde{c}_{ql}^{(3)}$,



Separate variation:
up to 30% deviation
from SM possible

Impact of the Wilson coefficients of the SM-EFT on $B \rightarrow K^{(*)} \nu \bar{\nu}$ vs. $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, assuming MFV and LFU, varied within their 2σ ranges allowed by the global fit to $b \rightarrow s \mu \mu$ data. Blue: \tilde{c}_Z , green: $\tilde{c}_{ql}^{(1)}$, red: $\tilde{c}_{ql}^{(3)}$. The dashed green line shows the case $\tilde{c}_{ql}^{(1)} = -\tilde{c}_{ql}^{(3)}$,

Correlations exist if additional assumptions made:

i.e. NP only through flavour changing Z couplings: $\tilde{c}_Z, \tilde{c}_Z^J$

- **Extra Slides**

Recent theory effort to eliminate perturbative uncertainties of 7%

NLO QCD corrections

Buchalla, Buras 1999, Misiak, Urban 1999

→ NNLO QCD corrections

Hermann, Misiak, Steinhauser arXiv:1311.1347

Leading- m_t NLO electroweak corrections

Buchalla, Buras 1998

→ NLO electroweak corrections

Bobeth, Gorbahn, Stamou arXiv:1311.1348

Experiment versus Theory

$$\overline{\mathcal{B}}_{s\mu}^{\text{exp}} = (2.9 \pm 0.7) \times 10^{-9}$$

$$\overline{\mathcal{B}}_{s\mu}^{\text{th}} = (3.65 \pm 0.23) \times 10^{-9}$$


$$\overline{\mathcal{B}}_{d\mu}^{\text{exp}} = (3.6_{-1.4}^{+1.6}) \times 10^{-10}$$

$$\overline{\mathcal{B}}_{d\mu}^{\text{th}} = (1.06 \pm 0.09) \times 10^{-10}$$

Error budget:


	f_{B_s}	CKM	τ_H^s	M_t	α_s	other param.	non-param.	Σ
\bar{B}_{sl}	4.0%	4.3%	1.3%	1.6%	0.1%	< 0.1%	1.5%	6.4%


Scale dependence:




Implications of the latest measurements of $B_s \rightarrow \mu\mu$

$$A_{SM} \sim m_\mu/m_b \Leftrightarrow A_{H^0, A^0} \sim \tan^3 \beta$$





Black line corresponds to direct search: ATLAS with 20.3 fb^{-1}



Black line corresponds to direct search: ATLAS with 20.3 fb^{-1}

Forthcoming theory analysis including all three independent angular observables ($z = \cos\theta$)

Huber, Hurth, Lunghi

- Dependence on Wilson coefficients

$$H_T(q^2) \propto 2s(1-s)^2 \left[\left| C_9 + \frac{2}{s} C_7 \right|^2 + |C_{10}|^2 \right]$$

$$H_A(q^2) \propto -4s(1-s)^2 \operatorname{Re} \left[C_{10} \left(C_9 + \frac{2}{s} C_7 \right) \right]$$

H_T suppressed in low- q^2 window

$$H_L(q^2) \propto (1-s)^2 \left[|C_9 + 2 C_7|^2 + |C_{10}|^2 \right]$$

- Devide low- q^2 bin in two bins (zero of H_A in low- q^2)
- Most important input parameters

$$m_b^{\text{IS}} = (4.691 \pm 0.037) \text{GeV}, \quad \bar{m}_c(\bar{m}_c) = (1.275 \pm 0.025) \text{GeV}$$
$$|V_{ts}^* V_{tb} / V_{cb}|^2 = 0.9621 \pm 0.0027, \quad BR_{b \rightarrow c e \nu}^{\text{exp.}} = (10.51 \pm 0.13) \%$$

- Perturbative expansion (NNLO QCD + NLO QED) α_s $\kappa = \alpha_{\text{em}} / \alpha_s$

$$A = \kappa \left[A_{LO} + \alpha_s A_{NLO} + \alpha_s^2 A_{NNLO} + \mathcal{O}(\alpha_s^3) \right]$$
$$+ \kappa^2 \left[A_{LO}^{\text{em}} + \alpha_s A_{NLO}^{\text{em}} + \alpha_s^2 A_{NNLO}^{\text{em}} + \mathcal{O}(\alpha_s^3) \right] + \mathcal{O}(\kappa^3)$$

Forthcoming theory analysis including all three independent angular observables ($z = \cos\theta$)

Huber, Hurth, Lunghi

- Collinear Photons give rise to log-enhanced QED corrections $\alpha_{\text{em}} \log(m_b^2/m_\ell^2)$
- Higher powers of z in double differential decay width
 - Definition of H_i ? Sensitivity for QED observables ?
- Size of logs depend on experimental set-up

$$q^2 = (p_{e^+} + p_{e^-})^2 \quad \text{vs.} \quad q^2 = (p_{e^+} + p_{e^-} + p_{\gamma, \text{coll}})^2$$

- We assume no photons are included in the definition of q^2 (di-muon channel at Babar/Belle, di-electron at Belle)
- Babar's di-electron channel: Photons that are emitted in a cone of 35 mrad angular opening are included in q^2

Monte Carlo techniques needed to estimate this effect

$$\frac{[\mathcal{B}_{ee}^{\text{low}}]_{q=p_{e^+}+p_{e^-}+p_{\gamma, \text{coll}}}}{[\mathcal{B}_{ee}^{\text{low}}]_{q=p_{e^+}+p_{e^-}}} - 1 = 1.65\%$$



$$\frac{[\mathcal{B}_{ee}^{\text{high}}]_{q=p_{e^+}+p_{e^-}+p_{\gamma, \text{coll}}}}{[\mathcal{B}_{ee}^{\text{high}}]_{q=p_{e^+}+p_{e^-}}} - 1 = 6.8\%$$

Forthcoming theory analysis including all three independent angular observables ($z = \cos\theta$)



Huber, Hurth, Lunghi

Monte Carlo analysis

(event generator EVTGEN, hadronization JETSET, EM radiation PHOTOS)



Validation of MC analysis (without Photos)



Results

Low- q^2 ($1\text{GeV}^2 < q^2 < 6\text{GeV}^2$)

$$BR(B \rightarrow X_s ee) = (1.67 \pm 0.10) 10^{-6} \text{ (preliminary)}$$

$$BR(B \rightarrow X_s \mu\mu) = (1.62 \pm 0.09) 10^{-6} \text{ (preliminary)}$$

Babar: $BR(B \rightarrow X_s ll) =$

$$= (1.60 (+0.41 - 0.39)_{stat} (+0.17 - 0.13)_{syst} (\pm 0.18)_{mod}) 10^{-6}$$

good agreement with SM

Forthcoming theory analysis including all three independent angular observables ($z = \cos\theta$)

Huber, Hurth, Lunghi

Results

High- q^2 , Theory: $q^2 > 14.4 \text{ GeV}^2$, Babar: $q^2 > 14.2 \text{ GeV}^2$

$$BR(B \rightarrow X_s ee) = (0.220 \pm 0.070) 10^{-6} \text{ (preliminary)}$$

$$BR(B \rightarrow X_s \mu\mu) = (0.253 \pm 0.070) 10^{-6} \text{ (preliminary)}$$

Babar: $BR(B \rightarrow X_s ll) =$

$$(0.57 (+0.16 - 0.15)_{stat} (+0.03 - 0.02)_{syst}) 10^{-6}$$

2σ higher than SM

Comparison with $B \rightarrow K^* ll$ data >>>

Further refinement

- Normalization to semileptonic $B \rightarrow X_u \ell \nu$ decay rate **with the same cut** reduces the impact of $1/m_b$ corrections in the high- q^2 region significantly.

Ligeti, Tackmann arXiv:0707.1694

Theory prediction for ratio (preliminary)

$$R(s_0)_{ee} = (2.25 \pm 0.31) 10^{-3}$$

$$R(s_0)_{\mu\mu} = (2.62 \pm 0.30) 10^{-3}$$

- Additional $O(5\%)$ uncertainty due to nonlocal power corrections $O(\alpha_s \Lambda/m_b)$

Forthcoming theory analysis including all three independent angular observables ($z = \cos\theta$)

Huber,Hurth,Lunghi

Further results in units of 10^{-6} (preliminary)

$$H_L[1, 3.5]_{ee} = 0.64 \pm 0.03$$

$$H_L[1, 3.5]_{\mu\mu} = 0.68 \pm 0.04$$

$$H_L[3.5, 6]_{ee} = 0.50 \pm 0.03$$

$$H_L[3.5, 6]_{\mu\mu} = 0.53 \pm 0.03$$

$$H_L[1, 6]_{ee} = 1.13 \pm 0.06$$

$$H_L[1, 6]_{\mu\mu} = 1.21 \pm 0.07$$

$$H_T[1, 3.5]_{ee} = 0.29 \pm 0.02$$

$$H_T[1, 3.5]_{\mu\mu} = 0.21 \pm 0.01$$

$$H_T[3.5, 6]_{ee} = 0.24 \pm 0.02$$

$$H_T[3.5, 6]_{\mu\mu} = 0.19 \pm 0.02$$

$$H_T[1, 6]_{ee} = 0.53 \pm 0.04$$

$$H_T[1, 6]_{\mu\mu} = 0.40 \pm 0.03$$

$$H_A[1, 3.5]_{ee} = -0.103 \pm 0.005$$

$$H_A[1, 3.5]_{\mu\mu} = -0.110 \pm 0.005$$

$$H_A[3.5, 6]_{ee} = +0.073 \pm 0.012$$

$$H_A[3.5, 6]_{\mu\mu} = +0.067 \pm 0.012$$

$$H_A[1, 6]_{ee} = -0.029 \pm 0.016$$

$$H_A[1, 6]_{\mu\mu} = -0.042 \pm 0.016$$

Total error $\mathcal{O}(5 - 8\%)$. Still dominated by scale uncertainty.

