


27th Rencontres de Blois

Château Royal de Blois — June 2nd, 2015

Leptonic flavour models

Christoph Luhn



Outline

- ▶ motivation
 - triplication of chiral families
 - observation of mixing
 - simple (approximate) patterns
- ▶ non-Abelian discrete family symmetries
 - finite groups
 - direct implementation
 - spontaneous breaking of family symmetry (flavon alignment)
- ▶ accommodating large θ_{13}
 - new family symmetries with more structure
 - perturbation of simple mixing patterns
 - testable correlations among mixing parameters

Fermion mixing

- ▶ mismatch of flavour (weak) and mass eigenstates

$$\Psi_{\text{flavour}} = V^\dagger \Psi_{\text{mass}}$$

- ▶ quark sector: V_L^u and V_L^d

$$U_{\text{CKM}} = V_L^u V_L^{d\dagger} \approx \begin{pmatrix} 1 & \lambda & \lambda^3 \\ \lambda & 1 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix} \quad \lambda \sim 0.22$$

- ▶ lepton sector: V_L^e and V_L^ν

$$U_{\text{PMNS}} = V_L^e V_L^{\nu\dagger} \approx \begin{pmatrix} 0.82 & 0.55 & 0.15 \\ 0.37 & 0.57 & 0.70 \\ 0.39 & 0.59 & 0.68 \end{pmatrix}$$

www.nu-fit.org (2014)

neutrino mixing pattern suggestive of some underlying principle

Simple mixing patterns – tri-bimaximal



Harrison



Perkins




Scott

$$U_{\text{PMNS}} \approx U_{\text{TB}} \equiv \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix}$$

$$\theta_{12} \approx 35.3^\circ \quad \theta_{23} = 45^\circ \quad \theta_{13} = 0^\circ$$

Simple mixing patterns – golden ratio



$$\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618$$

$$\tan \theta_{12} = \frac{1}{\varphi}$$


$$U_{\text{PMNS}} \approx U_{\text{GR}} \equiv \begin{pmatrix} \cos \theta_{12} & \sin \theta_{12} & 0 \\ -\frac{\sin \theta_{12}}{\sqrt{2}} & \frac{\cos \theta_{12}}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{\sin \theta_{12}}{\sqrt{2}} & \frac{\cos \theta_{12}}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}$$

$$\theta_{12} \approx 31.7^\circ \quad \theta_{23} = 45^\circ \quad \theta_{13} = 0^\circ$$

Family symmetries (aka horizontal symmetries)

Candidate symmetry groups G

- **non-Abelian** to unify families (G should have triplet representations)
- **discrete** to facilitate obtaining simple mixing patterns



Essentials of finite group theory – S_4 example

e.g. Ramond, Group theory: a physicist's survey (2010)


- finite number of group elements
- construct all elements from a small number of generators
- presentation of S_4 : generators S, T, U which satisfy
$$S^2 = T^3 = U^2 = (ST)^3 = (SU)^2 = (TU)^2 = (STU)^4 = 1$$
- matrix representations for $S, T, U \rightarrow$ irreps $\mathbf{1} \quad \mathbf{1}' \quad \mathbf{2} \quad \mathbf{3} \quad \mathbf{3}'$
- in physics we are mainly interested in multiplication rules
- Kronecker products e.g. $\mathbf{3} \otimes \mathbf{3} = \mathbf{1} + \mathbf{2} + \mathbf{3} + \mathbf{3}'$
- Clebsch-Gordan coefficients e.g. $\mathbf{3} \otimes \mathbf{3} \rightarrow \mathbf{1}$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} \rightarrow \alpha_1\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_2 \quad \text{basis dependent !!}$$

Two model building strategies


direct models

(mixing from residual symmetries)



indirect models

(mixing from flavon alignments Φ)



King, Luhn (JHEP 10, 2009)

Building a direct model with tri-bimaximal mixing

- choose family symmetry group – S_4
- identify VEV configurations for family symmetry breaking fields ϕ

$$S\langle\phi^\nu\rangle = U\langle\phi^\nu\rangle = \langle\phi^\nu\rangle \quad T\langle\phi^\ell\rangle = \langle\phi^\ell\rangle \quad \text{flavon VEVs}$$

S_4	S	U	T	$\langle\phi^\nu\rangle$	$\langle\phi^\ell\rangle$
$\mathbf{1}, \mathbf{1}'$	1	± 1	1	$\mathbf{1}$	$\mathbf{1}, \mathbf{1}'$
$\mathbf{2}$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}$	$\mathbf{2} \propto \begin{pmatrix} 1 \\ 1 \end{pmatrix}$	–
$\mathbf{3}, \mathbf{3}'$	$\frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$	$\mp \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}$	$\mathbf{3}' \propto \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$	$\mathbf{3}, \mathbf{3}' \propto \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

- control coupling of flavons to fermions by extra Z_N or $U(1)$ symmetry

$$\frac{\phi^\nu}{\Lambda^2} LH_u LH_u \quad \rightarrow \quad M_\nu = S^T M_\nu S = U^T M_\nu U \quad \rightarrow \quad \text{tri-bimaximal } M_\nu$$


$$\frac{\phi^\ell}{\Lambda} L \ell^c H_d \quad \rightarrow \quad [M_\ell M_\ell^\dagger] = T^T [M_\ell M_\ell^\dagger] T^* \quad \rightarrow \quad \text{diagonal } [M_\ell M_\ell^\dagger]$$

Accommodating $\theta_{13} \sim 9^\circ$

King, Luhn (Rept. Prog. Phys. 76, 2013)

King et al. (New J. Phys. 16, 2014)

Direct models after Daya Bay and RENO



mixing patterns:

	θ_{13}	θ_{23}	θ_{12}
TB	0°	45°	35.3°
BM	0°	45°	45°
GR	0°	45°	31.7°
BT	12.2°	36.2°	36.2°
TM	$\neq 0^\circ$	$\neq 45^\circ$	35.3°

TB = tri-bimaximal
 BM = bimaximal
 GR = golden ratio
 BT = bi-trimaximal
 TM = trimaximal

New family symmetries

- scans of “small” finite groups [Holthausen, Lim, Lindner \(Phys. Lett. B721, 2013\)](#)

$$\Delta(6 \cdot 10^2) \quad \Delta(6 \cdot 16^2) \quad (Z_{18} \times Z_6) \rtimes S_3 \subset \Delta(6 \cdot 18^2)$$

- mixing patterns derived from $\Delta(6n^2)$ [King, Neder \(Phys. Lett. B726, 2013\)](#)

$$U_{\Delta(6n^2)} = \begin{pmatrix} \sqrt{\frac{2}{3}} \cos \vartheta & \sqrt{\frac{1}{3}} & \sqrt{\frac{2}{3}} \sin \vartheta \\ -\sqrt{\frac{2}{3}} \sin(\frac{\pi}{6} + \vartheta) & \sqrt{\frac{1}{3}} & \sqrt{\frac{2}{3}} \cos(\frac{\pi}{6} + \vartheta) \\ \sqrt{\frac{2}{3}} \sin(\frac{\pi}{6} - \vartheta) & -\sqrt{\frac{1}{3}} & \sqrt{\frac{2}{3}} \cos(\frac{\pi}{6} - \vartheta) \end{pmatrix} \quad \text{with} \quad \vartheta = \pi \frac{p}{n}$$

- all possible mixing patterns from finite groups [Fonseca, Grimus \(JHEP 09, 2014\)](#)
 - 17 sporadic cases, but all incompatible with observed mixing
 - one series of mixing patterns related to $\Delta(6n^2)$

limited possibilities for lepton mixing from residual symmetries alone

Perturbation I – solar mixing sum rule

- T symmetry of charged lepton sector “slightly” broken (e.g. GUTs)
- $U_{\text{PMNS}} = V_{\ell_L} V_{\nu_L}^\dagger$ and $V_{\nu_L}^\dagger = U_{\text{TB}}$

$$U_{\text{PMNS}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & \hat{s}_{23} \\ 0 & -\hat{s}_{23}^* & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & \hat{s}_{13} \\ 0 & 1 & 0 \\ -\hat{s}_{13}^* & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & \hat{s}_{12} & 0 \\ -\hat{s}_{12}^* & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$s_{12} e^{i\delta_{12}} \approx \frac{1}{\sqrt{3}} \left(e^{i\delta_{12}^\nu} - \theta_{12}^\ell e^{i\delta_{12}^\ell} + \theta_{13}^\ell e^{i(\delta_{13}^\ell - \delta_{23}^\nu)} \right)$$

$$s_{23} e^{i\delta_{23}} \approx \frac{1}{\sqrt{2}} \left(e^{i\delta_{23}^\nu} - \theta_{23}^\ell e^{i\delta_{23}^\ell} \right)$$

$$s_{13} e^{i\delta_{13}} \approx \frac{1}{\sqrt{2}} \left(-\theta_{12}^\ell e^{i(\delta_{12}^\ell + \delta_{23}^\nu)} - \theta_{13}^\ell e^{i\delta_{13}^\ell} \right)$$

$$c_{ij} = \cos \theta_{ij}$$


$$\hat{s}_{ij} = \sin \theta_{ij} e^{-i\delta_{ij}}$$

- $\theta_{12}^\ell \sim \theta_C \sim 0.22 \rightarrow \theta_{13} \sim 9^\circ$

- first order relation $\theta_{12} \approx 35.3^\circ + \theta_{13} \cos \delta$

Testing the solar sum rule

- JUNO will measure θ_{12} with high precision
- wide-band superbeam (LBNO/LBNE) could access Dirac phase δ
- expected sensitivity for ruling out solar sum rule



Ballett et al.
(JHEP 12, 2014)

Perturbation II – atmospheric mixing sum rule


- U symmetry of neutrino sector “slightly” broken $\rightarrow U_{\text{PMNS}}^{13} \neq 0$
- conserve one Z_2 symmetry, either S or SU

	<u>trimaximal 1 (TM₁)</u>	<u>trimaximal 2 (TM₂)</u>
unbroken Z_2	$SU = -\frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & -1 \\ 2 & -1 & 2 \end{pmatrix}$	$S = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$
PMNS mixing	$\frac{1}{\sqrt{6}} \begin{pmatrix} 2 & \cdot & \cdot \\ -1 & \cdot & \cdot \\ -1 & \cdot & \cdot \end{pmatrix}$	$\frac{1}{\sqrt{3}} \begin{pmatrix} \cdot & 1 & \cdot \\ \cdot & 1 & \cdot \\ \cdot & 1 & \cdot \end{pmatrix}$
solar angle	$\theta_{12} \approx 34.2^\circ$	$\theta_{12} \approx 35.8^\circ$
first order relation	$\theta_{23} \approx 45^\circ + \sqrt{2} \theta_{13} \cos \delta$	$\theta_{23} \approx 45^\circ - \frac{1}{\sqrt{2}} \theta_{13} \cos \delta$


Testing the atmospheric sum rule

- low energy neutrino factory could measure θ_{23} and δ to high precision
- expected sensitivity for ruling out atmospheric sum rule

trimaximal 1



trimaximal 2



Ballett et al.

(Phys. Rev. D89, 2014)

Summary

- ▶ non-Abelian (discrete) symmetries
 - unify three families of chiral fermions
 - still attractive despite $\theta_{13} \sim 9^\circ$
- ▶ direct models with realistic θ_{13}
 - large family symmetries [e.g. $\Delta(600)$]
 - small family symmetries [e.g. S_4] plus perturbations
 - testable **mixing sum rules**
involving CP phase δ
- ▶ gain predictivity by imposing CP symmetry → talk by F. Feruglio
- ▶ crucial to measure mixing parameters (incl. δ) to a high precision
→ talk by M. Zito

Indirect models

Building an indirect model with tri-bimaximal mixing

- family symmetry $G \subset SU(3)$
- diagonal charged leptons
- type I seesaw with 2 or 3 ν_a^c in singlet representation of G
- diagonal right-handed neutrino mass matrix (e.g. due to Z_2 symmetry)


$$\mathcal{L}_\nu \sim \sum_a \frac{\phi_a^\nu}{\Lambda} L H_u \nu_a^c + M_a \nu_a^c \nu_a^c$$

- $\phi_a^\nu \sim \bar{\mathbf{3}}$ and $L \sim \mathbf{3}$ of G
- G or $SU(3)$ invariant $\rightarrow \phi_{a1}^\nu L_1 + \phi_{a2}^\nu L_2 + \phi_{a3}^\nu L_3 = \phi_a^{\nu T} L$
- integrate out ν_a^c (seesaw formula)

$$\mathcal{L}_\nu \sim L^T \sum_a \left(\frac{\langle \phi_a^\nu \rangle}{\Lambda} \cdot \frac{1}{M_a} \cdot \frac{\langle \phi_a^\nu \rangle^T}{\Lambda} \right) L H_u H_u$$

- tri-bimaximal if $\left[\langle \phi_1^\nu \rangle \propto \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix} \right] \quad \langle \phi_2^\nu \rangle \propto \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \langle \phi_3^\nu \rangle \propto \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$

Indirect models after Daya Bay and RENO



flavon alignments:

	$\langle \Phi_2 \rangle$	$\langle \Phi_3 \rangle$
CSD	$\frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$	$\frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$
CSD2	$\frac{1}{\sqrt{5}} \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$	$\frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$
PCSD	$\frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$	$\frac{1}{\sqrt{2}} \begin{pmatrix} \epsilon \\ 1 \\ -1 \end{pmatrix}$

other alignments possible as well

e.g. CSD3, CSD4,

King (JHEP 07, 2013 &
Phys. Lett. B724, 2013)

Variations of constrained sequential dominance (CSD)

$$m_\nu = m_2^0 \Phi_2 \Phi_2^T + m_3^0 \Phi_3 \Phi_3^T$$

$$m_2^0 \ll m_3^0$$

- CSD King (JHEP 08, 2005)

tri-bimaximal

$$\frac{m_2^0}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} + \frac{m_3^0}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix} \implies$$

$$\begin{array}{ll} \theta_{13} = 0 & m_1^\nu = 0 \\ \theta_{23} = 45^\circ & m_2^\nu = m_2^0 \\ \theta_{12} = 35.3^\circ & m_3^\nu = m_3^0 \end{array}$$

- CSD2 Antusch et al. (Nucl. Phys. B856, 2012)

trimaximal 1 (TM₁)

$$\frac{m_2^0}{5} \begin{pmatrix} 1 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{m_3^0}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix}$$

$$\begin{array}{ll} \theta_{13} \approx \frac{\sqrt{2}}{3} \frac{m_2^\nu}{m_3^\nu} & m_1^\nu = 0 \\ \theta_{23} \approx 45^\circ + \sqrt{2} \theta_{13} \cos \delta & m_2^\nu \approx \frac{3}{5} m_2^0 \\ \theta_{12} \approx 35.3^\circ & m_3^\nu \approx m_3^0 \end{array}$$

- P(artially)CSD King (Phys. Lett. B675, 2009)

tri-bimaximal-reactor

$$\frac{m_2^0}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} + \frac{m_3^0}{2} \begin{pmatrix} \epsilon^2 & \epsilon & -\epsilon \\ \epsilon & 1 & -1 \\ -\epsilon & -1 & 1 \end{pmatrix} \implies$$

$$\begin{array}{ll} \theta_{13} \approx \frac{\epsilon}{\sqrt{2}} & m_1^\nu = 0 \\ \theta_{23} \approx 45^\circ & m_2^\nu \approx m_2^0 \\ \theta_{12} \approx 35.3^\circ & m_3^\nu \approx m_3^0 \end{array}$$

Flavon alignment

Aligning triplet flavons in $\Delta(27)$, $Z_7 \rtimes Z_3$, A_4

$$V(\phi) = -m^2 \sum_i \phi_i^\dagger \phi_i + \lambda \left(\sum_i \phi_i^\dagger \phi_i \right)^2 + \Delta V$$

central terms in ΔV

$$(i) \quad \kappa \sum_i \phi_i^\dagger \phi_i \phi_i^\dagger \phi_i \quad \kappa > 0 \rightarrow \langle \phi \rangle \propto \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

$$\kappa < 0 \rightarrow \langle \phi \rangle \propto \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

$$(ii) \quad \tilde{\kappa} \sum_{i,j} (\phi_i^\dagger \tilde{\phi}_i)(\tilde{\phi}_j^\dagger \phi_j) \quad \tilde{\kappa} > 0 \rightarrow \text{orthogonality condition } \langle \phi \rangle \perp \langle \tilde{\phi} \rangle$$

$$\text{e.g. } \langle \tilde{\phi} \rangle \propto \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$$

$$\dots \langle \tilde{\phi} \rangle \propto \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}$$

Flavon alignment in supersymmetry

- SUSY unbroken at scale of family symmetry breaking
- introduce so-called driving fields X which couple to flavons
- flavon superpotential W^{flavon} linear in X due to $U(1)_R$ symmetry
- F -terms of driving fields need to vanish

$$F_{X_i}^* = -\frac{\partial W^{\text{flavon}}}{\partial X_i} = 0$$

- two examples in S_4

$$W^{\text{flavon}} \sim X_1 \phi_{\mathbf{2}} \phi_{\mathbf{2}} = X_1 (\phi_{\mathbf{2},1} \phi_{\mathbf{2},2} + \phi_{\mathbf{2},2} \phi_{\mathbf{2},1}) = 2X_1 \phi_{\mathbf{2},1} \phi_{\mathbf{2},2}$$

$$\longrightarrow \langle \phi_{\mathbf{2}} \rangle \propto \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{or} \quad \langle \phi_{\mathbf{2}} \rangle \propto \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$W^{\text{flavon}} = g_0 X_{\mathbf{3}} \phi_{\mathbf{3}'} \phi_{\mathbf{2}} + X_{\mathbf{3}'} (g_1 \phi_{\mathbf{3}'} \phi_{\mathbf{3}'} + g_2 \phi_{\mathbf{3}'} \phi_{\mathbf{2}} + g_3 \phi_{\mathbf{3}'} \phi_{\mathbf{1}})$$

$$\longrightarrow \langle \phi_{\mathbf{3}'} \rangle = \varphi_{\mathbf{3}'} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \langle \phi_{\mathbf{2}} \rangle = \varphi_{\mathbf{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \varphi_{\mathbf{2}} = -\frac{g_3}{2g_2} \varphi_{\mathbf{1}}$$

- flavon alignments independent of g_i

An S_4 benchmark model

King, Luhn (JHEP 09, 2011)

Direct model of leptons based on S_4

- discrete family symmetry with 24 elements
- irreducible S_4 representations: $\mathbf{1}$ $\mathbf{1}'$ $\mathbf{2}$ $\mathbf{3}$ $\mathbf{3}'$
- diagonal charged leptons (enforced by T symmetry)
- in neutrino sector ($L \sim N^c \sim \mathbf{3}$ $H_u \sim \mathbf{1}$)



$$W_\nu \sim y_D L N^c H_u + (y_{\mathbf{3}'} \phi_{\mathbf{3}'} + y_{\mathbf{2}} \phi_{\mathbf{2}} + y_{\mathbf{1}} \phi_{\mathbf{1}}) N^c N^c + \frac{y_{\mathbf{1}'}}{M} \tilde{\phi}_{\mathbf{1}'} \phi_{\mathbf{2}} N^c N^c$$

S_4 irrep	S	U	VEV alignment
$\mathbf{3}'$	$\frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$	$\langle \phi_{\mathbf{3}'} \rangle \propto \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$
$\mathbf{2}$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	$\langle \phi_{\mathbf{2}} \rangle \propto \begin{pmatrix} 1 \\ 1 \end{pmatrix}$
$\mathbf{1}$	1	1	$\langle \phi_{\mathbf{1}} \rangle \propto 1$
$\mathbf{1}'$	1	-1	$\langle \tilde{\phi}_{\mathbf{1}'} \rangle \propto 1$

U broken & S conserved \longrightarrow TM_2 mixing

Flavon alignment

$$\langle \phi_{\mathbf{3}'} \rangle = v_{\mathbf{3}'} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \langle \phi_{\mathbf{2}} \rangle = v_{\mathbf{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \langle \phi_{\mathbf{1}} \rangle = v_{\mathbf{1}} \quad \langle \tilde{\phi}_{\mathbf{1}'} \rangle = \tilde{v}_{\mathbf{1}'}$$

- SUSY unbroken at scale of family symmetry breaking
- F -terms of driving fields $\phi_{\mathbf{r}}^0$ need to vanish

$$\begin{aligned} W_{\nu}^{\text{flavon}} &= \phi_{\mathbf{3}'}^0 (g_1 \phi_{\mathbf{3}'} \phi_{\mathbf{3}'} + g_2 \phi_{\mathbf{3}'} \phi_{\mathbf{2}} + g_3 \phi_{\mathbf{3}'} \phi_{\mathbf{1}}) \\ &\quad + \phi_{\mathbf{3}}^0 (g_4 \phi_{\mathbf{3}'} \phi_{\mathbf{2}}) \\ &\quad + \phi_{\mathbf{1}}^0 (g_5 \phi_{\mathbf{3}'} \phi_{\mathbf{3}'} + g_6 \phi_{\mathbf{2}} \phi_{\mathbf{2}} + g_7 \phi_{\mathbf{1}} \phi_{\mathbf{1}}) \\ &\quad + \tilde{\phi}_{\mathbf{1}}^0 (g_8 \tilde{\phi}_{\mathbf{1}'} \tilde{\phi}_{\mathbf{1}'} + M^2) \end{aligned}$$

- previously assumed **flavon alignments independent of g_i** with

$$v_{\mathbf{2}} = -\frac{g_3}{2g_2} v_{\mathbf{1}} \quad v_{\mathbf{3}'}^2 = -\frac{1}{3g_5} \left(g_7 + \frac{g_3^2 g_6}{2g_2^2} \right) v_{\mathbf{1}}^2 \quad \tilde{v}_{\mathbf{1}'}^2 = -\frac{1}{g_8} M^2$$

Imposing CP symmetry (straightforward in S_4)

$$M_R = y_{\mathbf{3}'} v_{\mathbf{3}'} \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + y_{\mathbf{2}} v_{\mathbf{2}} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + y_{\mathbf{1}} v_{\mathbf{1}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \\ + \frac{y_{\mathbf{1}'}}{M} \tilde{v}_{\mathbf{1}'} v_{\mathbf{2}} \begin{pmatrix} 0 & 1 & -1 \\ 1 & -1 & 0 \\ -1 & 0 & 1 \end{pmatrix}$$

$$v_{\mathbf{2}} = -\frac{g_3}{2g_2} v_{\mathbf{1}} \quad v_{\mathbf{3}'}^2 = -\frac{1}{3g_5} \left(g_7 + \frac{g_3^2 g_6}{2g_2^2} \right) v_{\mathbf{1}}^2 \quad \tilde{v}_{\mathbf{1}'}^2 = -\frac{1}{g_8} M^2$$

- CP symmetry \rightarrow couplings y_i and g_i real
- phases of $v_{\mathbf{1}}$, $v_{\mathbf{2}}$, $v_{\mathbf{3}'}$ identical up to π or $\pm\pi/2$
- absorb phase of $v_{\mathbf{1}}$ into redefinition of N^c

	$v_{\mathbf{1}}$	$v_{\mathbf{2}}$	$v_{\mathbf{3}'}$	$\tilde{v}_{\mathbf{1}'}$
(A)	real	real	real	real
(B)	real	real	real	imaginary
(C)	real	real	imaginary	real
(D)	real	real	imaginary	imaginary

Predictions with CP symmetry

► seesaw mechanism

$$M_\nu = m_D M_R^{-1} m_D^T \quad \text{with} \quad m_D \propto \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\longrightarrow \boxed{U_{\text{PMNS}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} U_R} \quad \text{with} \quad U_R^T M_R U_R = M_R^{\text{diag}}$$

► resulting PMNS parameters

	θ_{13}	θ_{23}	θ_{12}	δ	(α_1, α_2)
(A)	free	$45^\circ \mp \frac{1}{\sqrt{2}}\theta_{13}$	35.3°	0 or π	0 or π
(B)	free	45°	35.3°	$\pm \frac{\pi}{2}$	0 or π
(C)	unphysical: two degenerate neutrino masses				
(D)	unphysical: $\theta_{13} = 35.3^\circ$				

\implies **five low-energy predictions** with imposed CP symmetry
(up to a finite choice)