

Usefulness of EFT for Boost Higgs Production

Ian Lewis
SLAC National Accelerator Laboratory


Dawson, IL, Zeng PRD90 (2014) 093007; PRD91 (2015) 074012

August 6, 2015
DPF 2015
Ann Arbor, MI

Discovered a Higgs Boson-Remarkably Standard Model Like

ATLAS Preliminary

$m_H = 125.36 \text{ GeV}$



$\sqrt{s} = 7 \text{ TeV} \int L dt = 4.5\text{-}4.7 \text{ fb}^{-1}$

$\sqrt{s} = 8 \text{ TeV} \int L dt = 20.3 \text{ fb}^{-1}$

released 12.01.2015

Signal strength (μ)

EFT for Boosted Higgs

$19.7 \text{ fb}^{-1} (8 \text{ TeV}) + 5.1 \text{ fb}^{-1} (7 \text{ TeV})$

CMS

$m_H = 125 \text{ GeV}$

$p_{SM} = 0.96$

Combined
 $\mu = 1.00 \pm 0.14$

$H \rightarrow \gamma\gamma$ tagged
 $\mu = 1.12 \pm 0.24$

$H \rightarrow ZZ$ tagged
 $\mu = 1.00 \pm 0.29$

$H \rightarrow WW$ tagged
 $\mu = 0.83 \pm 0.21$

$H \rightarrow \tau\tau$ tagged
 $\mu = 0.91 \pm 0.28$

$H \rightarrow bb$ tagged
 $\mu = 0.84 \pm 0.44$

0 0.5 1 1.5 2

Best fit σ/σ_{SM}

Ian Lewis (SLAC)


August 6, 2015

2 / 25


Future Measurements

ATLAS Simulation Preliminary


$\sqrt{s} = 14 \text{ TeV}$: $\int L dt = 300 \text{ fb}^{-1}$; $\int L dt = 3000 \text{ fb}^{-1}$



CMS Projection




CMS Projection



- Thoroughly testing mechanism of electroweak symmetry breaking.

Single Higgs Production

gg → h rate within δ of SM prediction



- Assume new physics heavy and parameterize as effective coupling:

$$\mathcal{L} = -\kappa_t \left(\frac{m_t}{v} \right) \bar{t} t h + \kappa_g \left(\frac{\alpha_s}{12\pi v} \right) G^{A,\mu\nu} G_{\mu\nu}^A h$$

- Degeneracy in Higgs production rate.
- Need to break degeneracy: tth or exploit different energy scalings between the SM and new physics contributions to the loop.


Higgs Plus Jet



;

- High energy jet $p_T \gtrsim m_{NP}$ can resolve physics inside of loop.
- Much work trying to study this possibility. Try to resolve if new physics is from shift in Yukawa coupling or new particles in loop. [Grojean et al JHEP \(2014\) 1405:022](#); [Azatov, Paul JHEP \(2014\) 1401:014](#); [Buschmann et al PRD90 \(2014\) 013010](#); [Buschmann et all JHEP 1502 \(2015\) 038](#); [Schlaffer et al EPJ C74 2014 3120](#); [Banfi, Martin, Sanz JHEP \(2014\) 1408:053](#)
- Bounds on new heavy particles quite high, try to simplify life and use EFT.


Higgs Plus Jet EFT



- EFT and SM contributions scale different with energy.
- Simple to keep single Higgs rate SM-like: $\kappa_t + \kappa_g = 1$.
- Allows more model independent approach: don't have to assume what new colored physics is giving new operator.

$$\mathcal{L} = -\kappa_t \left(\frac{m_t}{v} \right) \bar{t} t h + \kappa_g \left(\frac{\alpha_s}{12\pi v} \right) G^{A,\mu\nu} G_{\mu\nu}^A h$$

Higgs Plus Jet EFT




- Impose $\kappa_t + \kappa_g = 1$.
- Can see some deviation in tail.
- Direction of deviation determines direction that relevant couplings change relative to the Standard Model.

Schlaffer, Spannowsky, Takeuchi, Weiler, Wymant, Eur.Phys.J C74

(2014) 10, 3120

Effectiveness of EFT



- Previous results calculated using a Dimension-5 operator.

$$h G_{\mu\nu}^A G^{A,\mu\nu}$$

- Coefficient calculated by taking exact loop and expanding in $1/m^2$ and keeping LO in the expansion.
- How well does this EFT work.
 - Can calculate higher order, dimension-7, operators using next order in expansion.
 - Give an overview of the operator basis.
 - First present results in the SM, then exotic new physics.

Dimension 7 Operators

- Have Lagrangian:

$$\mathcal{L} = \mathcal{L}_{SM} + (\kappa_t - 1) \left(-\frac{m_t}{v} \right) \bar{t} t h + \mathcal{L}_5 + \mathcal{L}_7$$

- Dimension 5 operator:

$$\mathcal{L}_5 \equiv \hat{C}_1 O_1 = \hat{C}_1 G_{\mu\nu}^A G^{\mu\nu,A} h$$

- Four independent Dimension-7 operators relevant for Higgs production:

$$\mathcal{L}_7 = \sum_{i=2,3,4,5} \hat{C}_i O_i$$

$$O_2 = D_\sigma G_{\mu\nu}^A D^\sigma G^{\mu\nu,A} h \qquad \qquad O_3 = f_{ABC} G_v^{A,\mu} G_\sigma^{B,v} G_\mu^{C,\sigma} h$$

$$O_4 = g_s^2 h \sum_{i,j=1}^{n_{lf}} \overline{\Psi}_i \gamma_\mu T^A \Psi_i \overline{\Psi}_j \gamma^\mu T^A \Psi_j \qquad \qquad O_5 = g_s h \sum_{i,j=1}^{n_{lf}} G_{\mu\nu}^A D^\mu \overline{\Psi}_i \gamma^\nu T^A \Psi_i$$

Dimension 7 Operators

- Equations of motion can relate fermion operators to operators involving gluons only:

$$O_4 = g_s^2 h \sum_{i,j=1}^{n_{lf}} \bar{\Psi}_i \gamma_\mu T^A \Psi_i \bar{\Psi}_j \gamma^\mu T^A \Psi_j \rightarrow D^\sigma G_{\sigma\nu}^A D_\rho G^{A,\rho,\nu} h$$

$$O_5 = g_s h \sum_{i,j=1}^{n_{lf}} G_{\mu\nu}^A D^\mu \bar{\Psi}_i \gamma^\nu T^A \Psi_i \rightarrow G_{\sigma\nu}^A D^\nu D^\rho G_\rho^{A,\sigma} h$$

- Another useful operator is

$$O_6 = -D^\rho D_\rho (G_{\mu\nu}^A G^{\mu\nu,A}) h = m_h^2 O_1$$

- Using Jacobi identities:


$$O_6 = m_h^2 O_1 = -2O_2 + 4g_s O_3 + 4O_5$$

- Finally, choose basis $O_6 = m_h^2 O_1, O_3, O_4, O_5$, and the effective Lagrangian is

$$\mathcal{L}_{eff} = C_1 O_1 + (C_3 O_3 + C_4 O_4 + C_5 O_5)$$


- Since O_4 involves 4 light fermions, contributes to Higgs+jet starting at NLO.

LO Relative Contributions




- Operators matched onto SM.
- O_5 only contributes to $qg \rightarrow qh$: $O_5 = g_s h \sum_{i,j=1}^{n_{lf}} G_{\mu\nu}^A D^\mu \bar{\Psi}_i \gamma^5 T^A \Psi_i$
- O_3 only contributes to $gg \rightarrow gh$: $O_3 = f_{ABC} G_V^{A,\mu} G_\sigma^{B,\nu} G_\mu^{C,\sigma} h$
- qg production important in tail, O_5 more important to total distribution.

LO Relative Contributions



- Operators matched onto SM with $m_t \rightarrow \infty$
- O_3 makes little difference.
- O_5 important in tail of distribution, helps convergence.

Relative Contributions




- Operators matched onto SM.
- Higher order terms help convergence for $p_T \gtrsim 150 \text{ GeV}$.

Higher order corrections


- Higher order corrections have been calculated with dimension-5 operator.
 - Used for SM Higgs production at NNLO [Harlander, Kilgore, PRL88 \(2002\) 201801; Ravindran, Smith, van Neerven, NPB665 \(2003\) 325; Anastasiou, Melnikov NPB646 \(2002\) 220](#)
 - Used for SM Higgs p_T distribution [Anastasiou, Melnikov, Petriello NPB724 \(2005\) 197; Catani, Grazzini PRL98 \(2007\) 222002; Ravindran, Smith, van Neerven, NPB634 \(2002\) 247](#)
 - SM Higgs production to N^3LO [Anastasiou, Duhr, Dulat, Herzog, Mistlberger PRL114 \(2015\) 212001](#)
 - Higgs+jet at NNLO [Boughezal, Caola, Melnikov, Petriello, Schulze, arXiv:1504.07922](#)
- New operators could make a difference to in QCD corrections.
 - Different operators have different structures.
 - Could effect K-factors.
 - Most important in Higgs production with jets.
- Will present the numerical results for NLO Higgs+jet production with Wilson Coefficient matched onto SM.
- Can find equations for virtual corrections with arbitrary coefficients in [Dawson, IL, Zeng, PRD90 \(2014\), 034016](#)

NLO by operator




- O_1 is still dominant contribution.
- O_5 more important in the tail.
- O_3 smallest contribution.

NLO by channel





- Operators matched onto SM.
- O_5 now makes subdominant contribution to gg initial state.
- O_3 now makes subdominant contribution to qg initial state.

K-factor by Operator




- $O_5 = g_s h \sum_{i,j=1}^{n_{lf}} G_{\mu\nu}^A D^\mu \bar{\Psi}_i \gamma^\nu T^A \Psi_i$
- $O_3 = f_{ABC} G_\nu^{A,\mu} G_\sigma^{B,\nu} G_\mu^{C,\sigma} h$
- Flat K-factor for O_5 and O_1 .
- Increasing K-factor for O_3 .

K-factor by Channel



- New operators effect K-factor for $p_T \gtrsim 150 \text{ GeV}$.
- K-factor for qg initial state separates from other intial states for $p_T \gtrsim 150 \text{ GeV}$.

New Physics



- Add new physics, including higher dimensional operators.
- Test conclusions from SM in more exotic scenarios.
- Keep SM to exactly, perform comparison at LO in QCD.
- Consider two models:
 - Singlet top partner:

$$\begin{aligned} \mathcal{L} = & - \left\{ \cos^2 \theta_L \frac{m_t}{v} \bar{t}_L t_R h + \sin^2 \theta_L \frac{M_T}{v} \bar{T}_L T_R h \right. \\ & \left. + \frac{M_T}{2v} \sin(2\theta_L) \bar{t}_L T_R h + \frac{m_t}{2v} \sin(2\theta_L) \bar{T}_L t_R H + H.c. \right\} \end{aligned}$$

- Colored scalar:

$$V = V_{SM}(H) + m_i^2 \phi_i^\dagger \phi_i + \frac{C_h}{v} \phi_i^\dagger \phi_i H^\dagger H - \lambda_4 (\phi_i^\dagger \phi_t)^2$$



New Physics

- Besides Standard Model, can check validity of EFT in New physics.
 - Matched singlet top partner and colored scalar models onto Dim-7 EFT.

	Dirac Fermion	$SU(3)$ Triplet Scalar	$SU(3)$ Octet Scalar
$C_1(\Lambda)$	$\frac{\alpha_s \kappa_F}{12\pi v} \left[1 + \frac{7m_h^2}{120m_F^2} \right]$	$-\frac{\alpha_s}{96\pi M_S^2} C_h \left[1 + \frac{2m_h^2}{15M_S^2} \right]$	$-\frac{\alpha_s}{16\pi M_S^2} C_h \left[1 + \frac{2m_h^2}{15M_S^2} \right]$
$C_3(\Lambda)$	$-\frac{g_s \alpha_s \kappa_F}{360\pi v m_F^2}$	$-\frac{g_s \alpha_s}{1440 M_S^4} C_h$	$-\frac{g_s \alpha_s}{240 M_S^4} C_h$
$C_5(\Lambda)$	$\frac{11\kappa_F \alpha_s}{360\pi v m_F^2}$	$-\frac{\alpha_s}{360\pi M_S^4} C_h$	$-\frac{\alpha_s}{60\pi M_S^4} C_h$


- $\kappa_F = \sin^2 \theta_L$, θ_L is left-handed mixing angle between top and top partner.
- $C_h = 3m_Z$ is the scalar-scalar-Higgs boson triple coupling.

Triplet Scalar and Top Partner Results



- EFT error is relative deviation from exact LO QCD calculation.
- Dimension 5 works well for $p_T \lesssim$ new physics scale..
- Results with a few percent of exact result.
- Top quark contribution included exactly.



Triplet Scalar By Channel



- gg channel
 - Dimension 7 operators make mild difference.
 - Help convergence for $p_T \lesssim M_S$.

- qg channel
 - Dimension 7 operators, help convergence.

Distributions





- New physics makes little contribution to the differential or inclusive distribution.
- Inclusive distribution defined as:

$$\sigma(p_T > p_{T\text{cut}}) = \int_{p_{T\text{cut}}}^{\infty} dp_T \frac{d\sigma}{dp_T}$$

- How big of a new physics effect is needed to make appreciable difference?

Scaling Operators





- Red: SM
- Blue: Scaling of top yukawa, O_1
- Violet: Scaling of O_3
- $O_5 = g_s h \sum_{i,j=1}^{n_f} G_{\mu\nu}^A D^\mu \bar{\Psi}_i \gamma^5 T^A \Psi_i$, $O_3 = f_{ABC} G_v^{A,\mu} G_\sigma^{B,\nu} G_\mu^{C,\sigma} h$
- $O_1 = G_{\mu\nu}^A G^{A,\mu\nu} h$
- Kept single Higgs rate SM like.
- Scale operators by factor of 25 w.r.t. 500 GeV scalar triplet.

Conclusions

- Investigated how higher order operators effect Higgs+jet production.
- Standard Model:
 - Dimension-7 operators important and help convergence for $p_T \gtrsim 150$ GeV.
 - Main effect came from O_5 which effects qg initial state.
 - Reproduces exact LO p_T distribution within 10% for $p_T \lesssim 300$ GeV
 - At NLO, dimension-7 operators effect K-factor by a few to 10 percent for $p_T \gtrsim 150$ GeV.
 - Mainly because of K-factor for qg channel.
- New physics:
 - Investigated singlet top partner and color triplet scalar extensions of SM, specifically how important dimension-7 operators are to the rate.
 - Kept exact LO result from SM top quark.
 - Dimension-7 operators made little difference to p_T distribution for p_T below heavy particle's mass.
 - To make considerable contribution to p_T distributions need very large rescaling of dimension-7 operators.

EXTRA SLIDES

Higgs Plus Jet with Top Partner



Banfi, Martin, Sanz. JHEP 1408 (2014) 053

- Standard Model: Blue
- For $p_T < m_T$, see overall deviation insensitive to absolute scale of new physics.
- For $p_T \gtrsim m_T$, deviation changes.
- With current top partner limits $\sim 700 - 800 \text{ GeV}$, suggests using an EFT.