



Contribution ID: 324

Type: not specified

Fermi/LAT observations of Dwarf Galaxies highly constrain a Dark Matter Interpretation of Excess Positrons seen in AMS-02, HEAT, and PAMELA

Tuesday, 4 August 2015 16:30 (18 minutes)

It is shown that a Weakly Interacting Massive dark matter Particle (WIMP) interpretation for the positron excess observed in a variety of experiments, HEAT, PAMELA, and AMS-02, is highly constrained by the Fermi/LAT observations of dwarf galaxies. In particular, this paper has focused on the annihilation channels that best fit the current AMS-02 data (Boudaud et al., 2014). The Fermi satellite has surveyed the γ -ray sky, and its observations of dwarf satellites are used to place strong bounds on the annihilation of WIMPs into a variety of channels. For the single channel case, we find that dark matter annihilation into $\{b\bar{b}, e^+e^-, \mu^+\mu^-, \tau^+\tau^-, 4e, \text{ or } 4\tau\}$ is ruled out as an explanation of the AMS positron excess (here b quarks are a proxy for all quarks, gauge and Higgs bosons). In addition, we find that the Fermi/LAT 2σ upper limits, assuming the best-fit AMS-02 branching ratios, exclude multichannel combinations into $b\bar{b}$ and leptons. The tension between the results might relax if the branching ratios are allowed to deviate from their best-fit values, though a substantial change would be required. Of all the channels we considered, the only viable channel that survives the Fermi/LAT constraint and produces a good fit to the AMS-02 data is annihilation (via a mediator) to 4μ , or mainly to 4μ in the case of multichannel combinations.

Oral or Poster Presentation

Oral

Primary author: LOPEZ, Alejandro

Co-authors: SAVAGE, Christopher (NORDITA); ADAMS, Doug (Stockholm University); SPOLYAR, Doug (Stockholm University)

Presenter: LOPEZ, Alejandro

Session Classification: AstroParticle, Cosmology, Dark Matter Searches, and CMB

Track Classification: Astroparticles/Cosmic Rays