



Contribution ID: 358

Type: **not specified**

## Anatomy of Coannihilation with a Scalar Top Partner

*Friday 7 August 2015 16:20 (20 minutes)*

We present a simplified dark matter model where a Majorana fermion  $\chi$  coannihilates with a colored scalar top partner  $\tilde{t}$ . We explore the cosmological history, with particular emphasis on the most relevant low-energy parameters: the mass splitting between the dark matter and the coannihilator, and the Yukawa coupling that connects these fields to the Standard Model top quarks. We also allow a free quartic coupling between a pair of Higgs bosons and  $\tilde{t}$  pairs. We pay special attention to the case where the values take on those expected where  $\tilde{t}$  corresponds to the superpartner of the right-handed top, and  $\chi$  is a bino. Direct detection, indirect detection, and colliders are complementary probes of this simple model. We will also discuss implications for stop coannihilations within the MSSM.

### Oral or Poster Presentation

Oral

**Author:** Prof. PIERCE, Aaron (University of Michigan)**Co-authors:** Prof. IBARRA, Alejandro (Munich); Dr SHAH, Nausheen (University of Michigan); Dr VOGL, Stefan (Stockholm)**Presenter:** Prof. PIERCE, Aaron (University of Michigan)**Session Classification:** BSM Physics**Track Classification:** BSM Theory