



Contribution ID: 138

Type: parallel talk

Muon $(g - 2)_\mu$, Neutralino Dark Matter and the LHC

Monday 4 May 2015 16:45 (15 minutes)

We study the muon $(g - 2)_\mu$ anomaly in light of neutralino dark matter and the LHC. We scan the MSSM parameters relevant to $(g - 2)_\mu$ and focus on three different cases with different neutralino compositions. For $\tan \beta = 10$, we find that the 2σ range of $(g - 2)_\mu$ requires the smuon ($\tilde{\mu}_1$) to be lighter than ~ 500 GeV. The neutralinos, $\tilde{\chi}_1^0, \tilde{\chi}_2^0$, have to be lighter than ~ 300 GeV and 900 GeV respectively. In contrast, for $\tan \beta = 50$, the corresponding upper limits are ~ 1 TeV for $\tilde{\mu}_1$ and ~ 650 GeV for $\tilde{\chi}_1^0$ and 1 TeV for $\tilde{\chi}_2^0$. We thereby study the prospects of searching these light smuons and neutralinos at the LHC, in conjunction with constraints coming from indirect detection experiments of dark matter.

Authors: Dr AJAIB, M. Adeel (Ursinus Coll.); Mr GHOSH, Tathagata (Texas A&M University)

Co-authors: Dr DUTTA, Bhaskar (Texas A&M University); Dr GOGOLADZE, Ilia (University of Delaware); Dr SHAFI, Qaisar (University of Delaware)

Presenter: Mr GHOSH, Tathagata (Texas A&M University)

Session Classification: Dark Matter II