

Baryonic Dark Matter

Michael Duerr

Max-Planck-Institut für Kernphysik, Heidelberg



MAX-PLANCK-GESELLSCHAFT

Based on:

MD, P. Fileviez Pérez, M. B. Wise, **PRL 110**, 231801 (2013)

MD, P. Fileviez Pérez, **PLB 732** (2014) 101

MD, P. Fileviez Pérez, **PRD 91**, 095001 (2015)



MAX-PLANCK-INSTITUT
FÜR KERNPHYSIK

PHENO 2015

University of Pittsburgh

5 May 2015

The Great Desert




Figure: Wikipedia

Low scale
Electroweak scale
($\Lambda_{\text{EW}} \sim 10^2 \text{ GeV}$)



Figure: Wikipedia

$$\frac{c_6}{\Lambda_B^2} QQQL$$

S. Weinberg, PRL 43 (1979) 1566

High scale
e.g. GUT scale
($\Lambda_{\text{GUT}} \sim 10^{15} \text{ GeV}$)

Energy

The Great Desert

fermions		bosons	
generation	particle	generation	particle
1st	u, c, t, b quarks; e, muon, tau leptons	1st	W, Z bosons
2nd	d, s, b quarks; muon neutrino	2nd	Higgs boson
3rd	tau neutrino		

GALILEAN BOSONS

Figure: Wikipedia

Low scale
 Electroweak scale
 $(\Lambda_{EW} \sim 10^2 \text{ GeV})$



Figure: Wikipedia

$$\frac{C_6}{\Lambda_B^2} Q\bar{Q}L\bar{L}$$

S. Weinberg, PRL 43 (1979) 1566

High scale
 e.g. GUT scale
 $(\Lambda_{GUT} \sim 10^{15} \text{ GeV})$

Energy

The Great Desert

up	u	down	d	charm	s	top	t	bottom	b	gluon	g	Higgs boson	H
CHARGED LEPTONS													
electron	e	muon	μ	tau	τ	Z boson	Z	NEUTRAL BOSONS					
electron neutrino	ν_e	muon neutrino	ν_μ	tau neutrino	ν_τ	W boson	W						
neutrino	ν	neutrino	$\bar{\nu}$	neutrino	$\bar{\nu}$	neutrino	$\bar{\nu}$	neutrino	$\bar{\nu}$	neutrino	$\bar{\nu}$	neutrino	$\bar{\nu}$

Figure: Wikipedia

Low scale
 Electroweak scale
 $(\Lambda_{EW} \sim 10^2 \text{ GeV})$



Figure: Wikipedia

$$\frac{C_5}{\Lambda_L} LLHH$$

$$\frac{C_6}{\Lambda_B^2} QQQL$$

S. Weinberg, PRL 43 (1979) 1566

High scale
 e.g. GUT scale
 $(\Lambda_{GUT} \sim 10^{15} \text{ GeV})$

Energy

The Great Desert

CHARGED		NEUTRAL		CHARGED		NEUTRAL		CHARGED		NEUTRAL	
family	gen	family	gen	family	gen	family	gen	family	gen	family	gen
up	u	down	d	charm	c	top	t	gluon	g	Higgs	H
up	u	down	d	charm	c	top	t	gluon	g	Higgs	H
lepton	e	lepton	e	neutrino	ν_e	lepton	μ	neutrino	ν_μ	lepton	τ
lepton	e	lepton	e	neutrino	ν_e	lepton	μ	neutrino	ν_μ	lepton	τ
lepton	ν_e	lepton	ν_e	neutrino	ν_μ	lepton	ν_μ	neutrino	ν_τ	lepton	ν_τ
lepton	ν_e	lepton	ν_e	neutrino	ν_μ	lepton	ν_μ	neutrino	ν_τ	lepton	ν_τ
GALILEAN BOSONS											
W boson											

Figure: Wikipedia

Low scale
 Electroweak scale
 $(\Lambda_{EW} \sim 10^2 \text{ GeV})$



Figure: Wikipedia

High scale
 e.g. GUT scale
 $(\Lambda_{GUT} \sim 10^{15} \text{ GeV})$

Energy

The Great Desert

Quarks		Leptons		Gauge Bosons	
up	down	electron	neutrino	W boson	Z boson
u	d	e	ν_e		
c	s	μ	ν_μ		
t	b	τ	ν_τ		
g	v				Higgs boson

Figure: Wikipedia

Low scale
 Electroweak scale
 $(\Lambda_{EW} \sim 10^2 \text{ GeV})$

$$\frac{C_5}{\Lambda_L} LLHH$$



Figure: Wikipedia

$$\frac{C_6}{\Lambda_B^2} QQQL$$

S. Weinberg, PRL 43 (1979) 1566

High scale
 e.g. GUT scale
 $(\Lambda_{GUT} \sim 10^{15} \text{ GeV})$

Energy

What I will talk about today

Phenomenological aspects of a
consistent gauge theory for
baryon and lepton numbers
that can be broken at the low scale.

Baryonic and Leptonic Anomalies

- ▶ New gauge group:

$$SU(3) \otimes SU(2) \otimes U(1)_Y \otimes U(1)_B \otimes U(1)_L$$

- ▶ Purely baryonic anomalies:

$$\begin{aligned} \mathcal{A}_1 & (SU(3)^2 \otimes U(1)_B), \quad \mathcal{A}_2 (SU(2)^2 \otimes U(1)_B), \quad \mathcal{A}_3 (U(1)_Y^2 \otimes U(1)_B), \\ \mathcal{A}_4 & (U(1)_Y \otimes U(1)_B^2), \quad \mathcal{A}_5 (U(1)_B), \quad \mathcal{A}_6 (U(1)_B^3). \end{aligned}$$


- ▶ Purely leptonic anomalies:

$$\begin{aligned} \mathcal{A}_7 & (SU(3)^2 \otimes U(1)_L), \quad \mathcal{A}_8 (SU(2)^2 \otimes U(1)_L), \quad \mathcal{A}_9 (U(1)_Y^2 \otimes U(1)_L), \\ \mathcal{A}_{10} & (U(1)_Y \otimes U(1)_L^2), \quad \mathcal{A}_{11} (U(1)_L), \quad \mathcal{A}_{12} (U(1)_L^3). \end{aligned}$$

- ▶ Mixed anomalies:

$$\begin{aligned} \mathcal{A}_{13} & (U(1)_B^2 \otimes U(1)_L), \quad \mathcal{A}_{14} (U(1)_L^2 \otimes U(1)_B), \\ \mathcal{A}_{15} & (U(1)_Y \otimes U(1)_L \otimes U(1)_B). \end{aligned}$$

Field	$SU(3)$	$SU(2)$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
Q_L	3	2	$\frac{1}{6}$	$\frac{1}{3}$	0
u_R	3	1	$\frac{2}{3}$	$\frac{1}{3}$	0
d_R	3	1	$-\frac{1}{3}$	$\frac{1}{3}$	0
l_L	1	2	$-\frac{1}{2}$	0	1
e_R	1	1	0	0	1
H	1	2	$\frac{1}{2}$	0	0



Baryonic and Leptonic Anomalies

- ▶ New gauge group:

$$SU(3) \otimes SU(2) \otimes U(1)_Y \otimes U(1)_B \otimes U(1)_L$$

- ▶ Purely baryonic anomalies:

$$\begin{aligned} \mathcal{A}_1 & (SU(3)^2 \otimes U(1)_B), \quad \mathcal{A}_2 (SU(2)^2 \otimes U(1)_B), \quad \mathcal{A}_3 (U(1)_Y^2 \otimes U(1)_B), \\ \mathcal{A}_4 & (U(1)_Y \otimes U(1)_B^2), \quad \mathcal{A}_5 (U(1)_B), \quad \mathcal{A}_6 (U(1)_B^3). \end{aligned}$$

- ▶ Purely leptonic anomalies:

$$\begin{aligned} \mathcal{A}_7 & (SU(3)^2 \otimes U(1)_L), \quad \mathcal{A}_8 (SU(2)^2 \otimes U(1)_L), \quad \mathcal{A}_9 (U(1)_Y^2 \otimes U(1)_L), \\ \mathcal{A}_{10} & (U(1)_Y \otimes U(1)_L^2), \quad \mathcal{A}_{11} (U(1)_L), \quad \mathcal{A}_{12} (U(1)_L^3). \end{aligned}$$

- ▶ Mixed anomalies:

$$\begin{aligned} \mathcal{A}_{13} & (U(1)_B^2 \otimes U(1)_L), \quad \mathcal{A}_{14} (U(1)_L^2 \otimes U(1)_B), \\ \mathcal{A}_{15} & (U(1)_Y \otimes U(1)_L \otimes U(1)_B). \end{aligned}$$

Field	$SU(3)$	$SU(2)$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
Q_L	3	2	$\frac{1}{6}$	$\frac{1}{3}$	0
u_R	3	1	$\frac{2}{3}$	$\frac{1}{3}$	0
d_R	3	1	$-\frac{1}{3}$	$\frac{1}{3}$	0
l_L	1	2	$-\frac{1}{2}$	0	1
e_R	1	1	0	0	1
H	1	2	$\frac{1}{2}$	0	0

SM + right-handed ν 's

$$\mathcal{A}_2 = -\mathcal{A}_3 = \frac{3}{2},$$

$$\mathcal{A}_8 = -\mathcal{A}_9 = \frac{3}{2}$$

Baryonic and Leptonic Anomalies

- ▶ New gauge group:

$$SU(3) \otimes SU(2) \otimes U(1)_Y \otimes U(1)_B \otimes U(1)_L$$

Some History

- ▶
 - ▶ Early attempts to gauge B and L
 - ▶ A. Pais, PRD **8**, 1844 (1973)
 - ▶ S. Rajpoot, Int. J. Theor. Phys. **27**, 689 (1988)
 - ▶ R. Foot, G. C. Joshi, H. Lew, PRD **40**, 2487 (1989)
 - ▶ C. D. Carone, H. Murayama, PRD **52**, 484 (1995)
 - ▶ H. Georgi, S. L. Glashow, PLB **387**, 341 (1996)
- ▶ First realistic model
 - ▶ P. Fileviez Pérez, M.B. Wise, PRD **82**, 011901 (2010)

- ▶ Mixed anomalies:

$$\begin{aligned} \mathcal{A}_{13} & (U(1)_B^2 \otimes U(1)_L), \quad \mathcal{A}_{14} (U(1)_L^2 \otimes U(1)_B), \\ \mathcal{A}_{15} & (U(1)_Y \otimes U(1)_L \otimes U(1)_B). \end{aligned}$$

Field	$SU(3)$	$SU(2)$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
Q_L	3	2	$\frac{1}{6}$	$\frac{1}{3}$	0
u	3	1	2	1	0
e	0	1	1	1	0

$$\mathcal{A}_8 = -\mathcal{A}_9 = \frac{3}{2}$$

Solution: Vector-Like Lepto-Baryons

Field	$SU(3)$	$SU(2)$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
Ψ_L	1	2	$\pm\frac{1}{2}$	B_1	L_1
Ψ_R	1	2	$\pm\frac{1}{2}$	B_2	L_2
η_R	1	1	± 1	B_1	L_1
η_L	1	1	± 1	B_2	L_2
χ_R	1	1	0	B_1	L_1
χ_L	1	1	0	B_2	L_2

Anomaly cancellation demands: $B_1 - B_2 = L_1 - L_2 = -3$

MD, P. Fileviez Perez, M. B. Wise, arXiv:1304.0576 [hep-ph] (PRL)

Simple Version: Baryon Number Only

- ▶ Gauge group: $G_{\text{SM}} \otimes U(1)_B$
- ▶ Additional fields for an anomaly-free theory:

$$\begin{array}{ll} \Psi_L \sim (\mathbf{1}, \mathbf{2}, -1/2, B_1), & \Psi_R \sim (\mathbf{1}, \mathbf{2}, -1/2, B_2), \\ \eta_R \sim (\mathbf{1}, \mathbf{1}, -1, B_1), & \eta_L \sim (\mathbf{1}, \mathbf{1}, -1, B_2), \\ \chi_R \sim (\mathbf{1}, \mathbf{1}, 0, B_1), & \chi_L \sim (\mathbf{1}, \mathbf{1}, 0, B_2), \end{array}$$

- ▶ New Higgs for spontaneous breaking of B : S_B
- ▶ Condition from anomaly cancellation: $B_1 - B_2 = -3$.

MD, P. Fileviez Pérez, [arXiv:1309.3970 \[hep-ph\]](https://arxiv.org/abs/1309.3970) (PLB)

Spontaneous Symmetry Breaking

- ▶ Relevant interactions of the new fields (for $B_1 \neq -B_2$):

$$\begin{aligned} -\mathcal{L} \supset & h_1 \bar{\Psi}_L H \eta_R + h_2 \bar{\Psi}_L \tilde{H} \chi_R + h_3 \bar{\Psi}_R H \eta_L + h_4 \bar{\Psi}_R \tilde{H} \chi_L \\ & + \lambda_1 \bar{\Psi}_L \Psi_R S_B + \lambda_2 \bar{\eta}_R \eta_L S_B + \lambda_3 \bar{\chi}_R \chi_L S_B + \text{h.c.} \end{aligned}$$

$$S_B \sim (\mathbf{1}, \mathbf{1}, 0, B_1 - B_2)$$

- ▶ $\langle S_B \rangle \neq 0$ generates vector-like masses:

$$-\mathcal{L} \supset M_\Psi \bar{\Psi}_L \Psi_R + M_\eta \bar{\eta}_R \eta_L + M_\chi \bar{\chi}_R \chi_L + \text{h.c.}$$

$$S_B \sim (\mathbf{1}, \mathbf{1}, 0, -3) \rightarrow \Delta B = 3 \Rightarrow \text{no proton decay!}$$

- ▶ Remnant \mathcal{Z}_2 stabilizes lightest new fermion.

Baryonic Dark Matter

- ▶ Dirac DM, SM singlet-like: $\chi = \chi_R + \chi_L$
- ▶ Coupling to the new gauge boson:


$$\mathcal{L} \supset g_B \bar{\chi} \gamma_\mu Z_B^\mu (B_2 P_L + B_1 P_R) \chi$$

Annihilation and direct detection!

- ▶ Model has six free parameters:



$$M_\chi, M_{Z_B}, g_B, B \equiv B_1 + B_2 \text{ and } M_{h_2}, \theta_B$$

⇒ testable by combining LHC, DM direct detection, DM relic density.





MD, P. Fileviez Pérez, arXiv:1409.8165 [hep-ph] (PRD)

Decays of the new Gauge Boson Z_B



MD, P. Fileviez Pérez, [arXiv:1409.8165 \[hep-ph\]](https://arxiv.org/abs/1409.8165) (PRD)


LHC Bounds: $pp \rightarrow Z_B^* \rightarrow \bar{t}t$





$$\sigma(\bar{q}q \rightarrow Z_B^* \rightarrow \bar{t}t)(\hat{s}) = \frac{g_B^4 \sqrt{\hat{s} - 4M_t^2}}{972\pi\sqrt{\hat{s}}} \frac{(2M_t^2 + \hat{s})}{[(\hat{s} - M_{Z_B}^2)^2 + M_{Z_B}^2 \Gamma_{Z_B}^2]}$$

ATLAS-CONF-2013-052


Associated Production: $pp \rightarrow Z_B^* \rightarrow Z_B h_2$



Decays of the new Heavy Higgs h_2





$\bar{t}t$ Plus Missing Energy





$$N(\bar{t}t E_T^{\text{miss}}) = \mathcal{L} \times \sigma(pp \rightarrow Z_B h_2) \times \text{Br}(Z_B \rightarrow \bar{t}t) \times \text{Br}(h_2 \rightarrow \bar{\chi}\chi)$$

Dark Matter Relic Density





► Planck: $\Omega_{\text{DM}} h^2 = 0.1199 \pm 0.0027$

Dark Matter Relic Density



► Planck: $\Omega_{\text{DM}} h^2 = 0.1199 \pm 0.0027$

Dark Matter Direct Detection



► Planck: $\Omega_{\text{DM}} h^2 = 0.1199 \pm 0.0027$

$$M_{Z_B} \in [0.5, 5.0] \text{ TeV}$$

$$g_B \in [0.1, 0.5]$$

Bound on the Symmetry Breaking Scale

- Unsuppressed terms in the annihilation cross section:

$$\sum_q \sigma v(\bar{\chi}\chi \rightarrow Z_B^* \rightarrow \bar{q}q) = \frac{g_B^4 M_\chi^2}{2\pi} \frac{(B_1 + B_2)^2}{[(4M_\chi^2 - M_{Z_B}^2)^2 + M_{Z_B}^2 \Gamma_{Z_B}^2]}$$

- Using $\Omega_{\text{DM}} h^2 \leq 0.12$ and $M_\chi = \lambda_\chi v_B / \sqrt{2}$ and $M_{Z_B} = 3g_B v_B$, one obtains

$$v_B^2 \leq \frac{g_B^4 \lambda_\chi^2 (B_1 + B_2)^2}{\pi} \frac{1.77 \times 10^9 \text{ GeV}^2}{[(2\lambda_\chi^2 - 9g_B^2)^2 + \frac{9}{4\pi^2} g_B^8]} x_f$$

- For $x_f = 20$ and $B_1 + B_2 = 1/2$:

$$M_{Z_B} \leq 35.3 \text{ TeV} \text{ and } M_\chi \leq 17.7 \text{ TeV}$$

MD, P. Fileviez Pérez, arXiv:1409.8165 [hep-ph] (PRD)

Summary

Baryonic Dark Matter:

- ▶ B (and L) are gauge symmetries that can be broken at the low scale
- ▶ no proton decay \Rightarrow no need for a desert
- ▶ Fermionic DM candidate
- ▶ Testable at the LHC

Summary

Baryonic Dark Matter:

- ▶ B (and L) are gauge symmetries that can be broken at the low scale
- ▶ no proton decay \Rightarrow no need for a desert
- ▶ Fermionic DM candidate
- ▶ Testable at the LHC



Thank you!

Backup slides


Other Solution for Anomaly Cancellation

$$\begin{aligned}\Psi_L &\sim (\mathbf{1}, \mathbf{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}), & \Psi_R &\sim (\mathbf{1}, \mathbf{2}, \frac{1}{2}, -\frac{3}{2}, -\frac{3}{2}) \\ \Sigma_L &\sim (\mathbf{1}, \mathbf{3}, 0, -\frac{3}{2}, -\frac{3}{2}), & \chi_L &\sim (\mathbf{1}, \mathbf{1}, 0, -\frac{3}{2}, -\frac{3}{2})\end{aligned}$$

P. Fileviez Pérez, S. Ohmer, H. H. Patel, [arXiv:1403.8029 \[hep-ph\]](https://arxiv.org/abs/1403.8029)


- ▶ Less representations
- ▶ Same degrees of freedom after symmetry breaking
- ▶ Majorana dark matter

Mass of the Heavy Higgs h_2



$$M_{h_2}^2 = M_{h_1}^2 + \frac{2}{3g_B} |\csc 2\theta_B| \nu_0 M_{Z_B} \lambda_{HB}$$

Mixing Angle in the Higgs Sector



Gauging B and L in a SUSY Setup

► BLMSSM:

J. M. Arnold, P. Fileviez Pérez, B. Fornal, S. Spinner, [arXiv:1310.7052 \[hep-ph\]](https://arxiv.org/abs/1310.7052)

- $U(1)_B$ and $U(1)_L$ breaking scale linked to SUSY breaking scale
- R-parity must be spontaneously broken
- Stable DM candidate

Symmetries of the Model

- ▶ Symmetries of the Model:

- ▶ $(B - L)_{\text{SM}}$
- ▶ Accidental η symmetry in the new sector:

$$\Psi_{L,R} \rightarrow e^{i\eta} \Psi_{L,R},$$

$$\eta_{L,R} \rightarrow e^{i\eta} \eta_{L,R},$$

$$\chi_{L,R} \rightarrow e^{i\eta} \chi_{L,R}.$$

⇒ DM stability

- ▶ Total baryon number B_T (above symmetry-breaking scale)

Assume that scale for baryon number breaking is low, close to the EW scale.

Baryon and Dark Matter Asymmetries

- For $\mu \ll T$:

$$\frac{\Delta n}{s} = \frac{n_+ - n_-}{s} = \frac{15g_* \mu}{2\pi^2 g_* \xi T},$$

g : internal degrees of freedom,
 s : entropy density,
 g_* : total number of relativistic
 degrees of freedom,
 $\xi = 2$ for fermions,
 $\xi = 1$ for bosons

- $B - L$ asymmetry in the SM sector

$$\Delta(B - L)_{\text{SM}} = \frac{45}{4\pi^2 g_* T} (\mu_{u_L} + \mu_{u_R} + \mu_{d_L} + \mu_{d_R} - \mu_{\nu_L} - \mu_{e_L} - \mu_{e_R}),$$


- η charge density

$$\Delta\eta = \frac{15}{4\pi^2 g_* T} (2\mu_{\psi_L} + 2\mu_{\psi_R} + \mu_{\chi_L} + \mu_{\chi_R} + \mu_{\eta_L} + \mu_{\eta_R}).$$


P. Fileviez Pérez, H. H. Patel, [arXiv:1311.6472 \[hep-ph\]](https://arxiv.org/abs/1311.6472)

Baryon and Dark Matter Asymmetries

$$B_f^{\text{SM}} \equiv \frac{15}{4\pi^2 g_* T} (12\mu_{u_L}) = \frac{32}{99} \Delta(B - L)_{\text{SM}} + \frac{(15 - 14B_2)}{198} \Delta\eta,$$



$$3(3\mu_{u_L} + \mu_{e_L}) + \mu_{\Psi_L} - \mu_{\Psi_R} = 0.$$



P. Fileviez Pérez, H. H. Patel, [arXiv:1311.6472 \[hep-ph\]](https://arxiv.org/abs/1311.6472)

Left-Right Symmetric Model

$$G_{LR} = SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$$
SM fields

- ▶ Connects neutrino masses and spontaneous parity violation.
 - ▶ Standard version uses hybrid version of type I and type II seesaw mechanism for neutrino masses.
- $Q_L \sim (\mathbf{2}, \mathbf{1}, 1/3)$
 $Q_R \sim (\mathbf{1}, \mathbf{2}, 1/3)$
 $\ell_L \sim (\mathbf{2}, \mathbf{1}, -1)$
 $\ell_R \sim (\mathbf{1}, \mathbf{2}, -1)$

Pati, Salam, PRD **10** (1974) 275, Mohapatra, Pati, PRD **11** (1975) 2558, Senjanovic, Mohapatra, PRD **12** (1975) 1502

- ▶ Promote gauge group

$$\Rightarrow SU(2)_L \otimes SU(2)_R \otimes U(1)_B \otimes U(1)_L$$

He, Rajpoot, PRD **41** (1990) 1636

Anomaly Cancellation: Type III Seesaw

- ▶ Anomalies that need to be cancelled:

$$\mathcal{A}_1 (SU(2)_L^2 \otimes U(1)_B) = 3/2$$

$$\mathcal{A}_2 (SU(2)_L^2 \otimes U(1)_L) = 3/2$$

$$\mathcal{A}_3 (SU(2)_R^2 \otimes U(1)_B) = -3/2$$

$$\mathcal{A}_4 (SU(2)_R^2 \otimes U(1)_L) = -3/2$$


- ▶ Simplest solution: type III seesaw fields

$$\rho_L \sim (\mathbf{3}, \mathbf{1}, -3/4, -3/4),$$

$$\rho_R \sim (\mathbf{1}, \mathbf{3}, -3/4, -3/4)$$

- ▶ Neutrino mass matrix:

$$\mathcal{M}_\nu^{3+2} = \begin{pmatrix} 0 & 0 & 0 & m_D^1 & m_D^2 \\ 0 & m_1 & 0 & m_D^3 & m_D^4 \\ 0 & 0 & m_2 & m_D^5 & m_D^6 \\ m_D^1 & m_D^3 & m_D^5 & 0 & 0 \\ m_D^2 & m_D^4 & m_D^6 & 0 & 0 \end{pmatrix}.$$



- ▶ Two light sterile neutrinos.

MD, P. Fileviez Pérez, M. Lindner, [arXiv:1306.0568 \[hep-ph\]](https://arxiv.org/abs/1306.0568) (PRD)