

Experimental particle. physics

esipap...
European School of Instrumentation
in Particle & Astroparticle Physics

3.

particle interactions
in particle detectors

*lots of material borrowed from the excellent course of H.-C. Schultz-Coulon
<http://www.kip.uni-heidelberg.de/~coulon/Lectures/Detectors/>*



How do we detect particles?

- In order to detect a particle, it must:
 - ✓ interact with the material of the detector
 - ✓ transfer energy in some recognizable fashion (signal)
- Detection of particles happens via their energy loss in the material they traverses



Charged particles	Ionization, Bremsstrahlung, Cherenkov, ...	multiple interactions
Photons	Photo/Compton effect, pair production	single interactions...
Hadrons	Nuclear interactions	multiple interactions
Neutrinos	Weak interactions	

Example of particle interactions



Ionization:



Pair production:



Compton scattering:




Energy loss by ionization: Bethe-Bloch formula

For now assume: $Mc^2 \gg m_e c^2$

i.e. energy loss for heavy charged particles
[dE/dx for electrons more difficult ...]

Interaction dominated
by elastic collisions with electrons ...



Bethe-Bloch Formula

$$-\left\langle \frac{dE}{dx} \right\rangle = K z^2 \frac{Z}{A} \frac{1}{\beta^2} \left[\frac{1}{2} \ln \frac{2m_e c^2 \beta^2 \gamma^2 T_{\max}}{I^2} - \beta^2 - \frac{\delta(\beta\gamma)}{2} \right]$$

$$\propto 1/\beta^2 \cdot \ln(\text{const} \cdot \beta^2 \gamma^2)$$

Bethe-Bloch formula for heavy particles

[see e.g. PDG 2010]

$$-\left\langle \frac{dE}{dx} \right\rangle = K z^2 \frac{Z}{A} \frac{1}{\beta^2} \left[\frac{1}{2} \ln \frac{2m_e c^2 \beta^2 \gamma^2 T_{\max}}{I^2} - \beta^2 - \frac{\delta(\beta\gamma)}{2} \right]$$

[· ρ]

 density

$$K = 4\pi N_A r_e^2 m_e c^2 = 0.307 \text{ MeV g}^{-1} \text{ cm}^2$$

$$T_{\max} = 2m_e c^2 \beta^2 \gamma^2 / (1 + 2\gamma m_e / M + (m_e / M)^2)$$

[Max. energy transfer in single collision]

$$N_A = 6.022 \cdot 10^{23}$$

[Avogadro's number]

$$r_e = e^2 / 4\pi\epsilon_0 m_e c^2 = 2.8 \text{ fm}$$

[Classical electron radius]

$$m_e = 511 \text{ keV}$$

[Electron mass]

$$\beta = v/c$$

[Velocity]

$$\gamma = (1 - \beta^2)^{-2}$$

[Lorentz factor]

z : Charge of incident particle

M : Mass of incident particle

Z : Charge number of medium

A : Atomic mass of medium

I : Mean excitation energy of medium


δ : Density correction [transv. extension of electric field]

Validity:

$$.05 < \beta\gamma < 500$$

$$M > m_\mu$$

Energy loss of pions in Cu



$\beta\gamma = 3-4$

Minimum ionizing particles (MIP): $\beta\gamma = 3-4$

dE/dx falls $\sim \beta^{-2}$; kinematic factor
[precise dependence: $\sim \beta^{-5/3}$]

dE/dx rises $\sim \ln(\beta\gamma)^2$; relativistic rise
[rel. extension of transversal E-field]

Saturation at large $(\beta\gamma)$ due to density effect (correction δ)
[polarization of medium]

Units: $\text{MeV g}^{-1} \text{cm}^2$

MIP loses ~ 13 MeV/cm
[density of copper: 8.94 g/cm^3]



Understanding Bethe-Bloch

1/β²-dependence:

Slower particles feel electric force of atomic electrons for longer time ...

Relativistic rise for βγ > 4:

High energy particle: transversal electric field increases due to Lorentz transform; $E_y \rightarrow \gamma E_y$. Thus interaction cross section increases ...



Corrections:

- low energy : shell corrections
- high energy : density corrections

Understanding Bethe-Bloch

Density correction:

Polarization effect ...
[density dependent]

→ Shielding of electrical field far from particle path; effectively cuts off the long range contribution ...

More relevant at high γ ...
[Increased range of electric field; larger b_{\max} ; ...]

For high energies:


$$\delta/2 \rightarrow \ln(\hbar\omega/I) + \ln \beta\gamma - 1/2$$

Shell correction:

Arises if particle velocity is close to orbital velocity of electrons, i.e. $\beta c \sim v_e$.

Assumption that electron is at rest breaks down ...

Capture process is possible ...



Density effect leads to saturation at high energy ...

Shell correction are in general small ...

Energy loss of (heavy) charged particles


Dependence on

Mass A
 Charge Z
 of target nucleus


Minimum ionization:

ca. $1 - 2 \text{ MeV/g cm}^{-2}$
 [H_2 : 4 MeV/g cm^{-2}]

$$-\left\langle \frac{dE}{dx} \right\rangle \sim \frac{Z}{A}$$



Identifying particles by dE/dx




Measured
energy loss

[ALICE TPC, 2009]

Bethe-Bloch

Remember:
 dE/dx depends on β !

ALICE TPC



Pb+Pb @ sqrt(s) = 2.76 ATeV

2010-11-08 11:30:46

Fill : 1482

Run : 137124


Event : 0x00000000D3BBE693

dE/dx fluctuations

Bethe-Bloch describes **mean** energy loss; measurement via energy loss ΔE in a material of thickness Δx with

$$\Delta E = \sum_{n=1}^N \delta E_n$$

N : number of collisions
 δE : energy loss in a single collision



Ionization loss δE
distributed statistically ...

so-called
Energy loss 'straggling'

Complicated problem ...

Thin absorbers: **Landau distribution**

Standard Gauss with mean energy loss E_0
+ tail towards high energies due to δ -electrons

see also Allison & Cobb
[Ann. Rev. Nucl. Part. Sci. 30 (1980) 253.]

Mean particle range

Integrate over energy loss
from E down to 0


$$R = \int_E^0 \frac{dE}{dE/dx}$$

Example:

Proton with $p = 1 \text{ GeV}$
Target: lead with $\rho = 11.34 \text{ g/cm}^3$

$$R/M = 200 \text{ g cm}^{-2} \text{ GeV}^{-1}$$

$$\rightarrow R = 200/11.34/1 \text{ cm} \sim 20 \text{ cm}$$



Energy loss of electrons

Bethe-Bloch formula needs modification

Incident and target electron have same mass m_e
Scattering of identical, indistinguishable particles

$$-\left\langle \frac{dE}{dx} \right\rangle_{\text{el.}} = K \frac{Z}{A} \frac{1}{\beta^2} \left[\ln \frac{m_e \beta^2 c^2 \gamma^2 T}{2I^2} + F(\gamma) \right]$$


[T: kinetic energy of electron]

$$W_{\text{max}} = \frac{1}{2}T$$

Remark: different energy loss for electrons and positrons at low energy as positrons are not identical with electrons; different treatment ...

Bremsstrahlung and Radiation Length

Bremsstrahlung arises if particles are accelerated in Coulomb field of nucleus



$$\frac{dE}{dx} = 4\alpha N_A \frac{z^2 Z^2}{A} \left(\frac{1}{4\pi\epsilon_0} \frac{e^2}{mc^2} \right)^2 E \ln \frac{183}{Z^{\frac{1}{3}}} \propto \frac{E}{m^2}$$

i.e. energy loss proportional to $1/m^2$ → main relevance for electrons ...
... or ultra-relativistic muons

Consider electrons:

$$\frac{dE}{dx} = 4\alpha N_A \frac{Z^2}{A} r_e^2 \cdot E \ln \frac{183}{Z^{\frac{1}{3}}}$$

$$\frac{dE}{dx} = \frac{E}{X_0} \quad \text{with} \quad X_0 = \frac{A}{4\alpha N_A Z^2 r_e^2 \ln \frac{183}{Z^{\frac{1}{3}}}}$$

[Radiation length in g/cm²]

$$\Rightarrow E = E_0 e^{-x/X_0}$$

After passage of one X_0 electron has lost all but $(1/e)^{\text{th}}$ of its energy

[i.e. 63%]

Critical Energy

Critical energy:

$$\left. \frac{dE}{dx}(E_c) \right|_{\text{Brems}} = \left. \frac{dE}{dx}(E_c) \right|_{\text{Ion}}$$

$$\left(\frac{dE}{dx} \right)_{\text{Tot}} = \left(\frac{dE}{dx} \right)_{\text{Ion}} + \left(\frac{dE}{dx} \right)_{\text{Brems}}$$


Approximation:

$$E_c^{\text{Gas}} = \frac{710 \text{ MeV}}{Z + 0.92}$$


$$E_c^{\text{Sol/Liq}} = \frac{610 \text{ MeV}}{Z + 1.24}$$

Example Copper:


$$E_c \approx 610/30 \text{ MeV} \approx 20 \text{ MeV}$$



Total Energy Loss of Electrons




Energy loss for muons



Cherenkov radiation

Particles moving in a medium with **speed larger than speed of light in that medium** will lose energy by emitting electromagnetic radiation

- ✓ Charged particle polarize medium generating an electrical dipole varying in time
- ✓ Every point in trajectory emits a spherical EM wave, waves constructively interfere...



Cherenkov radiation: radiators

Parameters of Typical Radiator

Medium	n	β_{thr}	$\theta_{\text{max}} [\beta=1]$	$N_{\text{ph}} [\text{eV}^{-1} \text{cm}^{-1}]$
Air	1.000283	0.9997	1.36	0.208
Isobutan	1.00127	0.9987	2.89	0.941
Water	1.33	0.752	41.2	160.8
Quartz	1.46	0.685	46.7	196.4

Note: Energy loss by Cherenkov radiation very small w.r.t. ionization (< 1%).

Example:

[Proton with $E_{\text{kin}}=1$ GeV passing through 1 cm water]

$$\beta = p/E \approx 0.875; \cos\theta_C = 1/n\beta = 0.859 \rightarrow \theta_C = 30.8^\circ$$

$$d^2N/(dE dx) = 370 \sin^2\theta_C \text{ eV}^{-1} \text{ cm}^{-1} \approx 100 \text{ eV}^{-1} \text{ cm}^{-1}$$

$$\begin{aligned} \rightarrow \Delta E_{\text{loss}} &= \langle E \rangle d^2N/(dE dx) \Delta E \Delta x \\ &= 2.5 \text{ eV} \cdot 100 \text{ eV}^{-1} \text{ cm}^{-1} \cdot 5 \text{ eV} \cdot 1 \text{ cm} = 1.25 \text{ keV} \end{aligned}$$

Visible light only!


[$E = 1 - 5$ eV; $\lambda = 300 - 600$ nm]

$$\Delta E_{\text{loss}} < 1.25 \text{ keV}$$

Identifying particles with Cherenkov radiation

Threshold detection:

Observation of Cherenkov radiation $\rightarrow \beta > \beta_{\text{thr}}$



Choose n_1, n_2 in such a way that for:

$$n_2 : \quad \beta_{\pi}, \beta_K > 1/n_2 \text{ and } \beta_p < 1/n_2$$


$$n_1 : \quad \beta_{\pi} > 1/n_1 \text{ and } \beta_K, \beta_p < 1/n_1$$

Light in C_1 and C_2 \rightarrow identified pion

Light in C_2 and not in C_1 \rightarrow identified kaon

Light neither in C_1 and C_2 \rightarrow identified proton

Cherenkov radiation: momentum dependence



Cherenkov angle
Number of photons

grows with β and reaches asymptotic value for $\beta = 1$
 $[\theta_{\max} = \arccos(1/n); N_{\infty} = x \cdot 370/\text{cm} (1 - 1/n^2)]$

LHCb RICH

Measurement of Cherenkov angle:



Use medium with known refractive index $n \rightarrow \beta$

Principle of:

RICH (Ring Imaging Cherenkov Counter)

DIRC (Detection of Internally Reflected Cherenkov Light)

DISC (special DIRC; e.g. Panda)



LHCb RICH

Transition radiation

Transition radiation occurs if a relativistic particle (**large γ**) passes the **boundaries between two media with different refraction indices...**

- ✓ Intensity of radiation is **logarithmically proportional to γ**

Angular distribution strongly forward peaked
[Interference; coherence condition]

$$\theta \leq 1/\gamma$$

Coherent radiation is generated only over a very small formation length

$$D = \gamma c / \omega_p$$

Plasma frequency
[from Drude model]

Volume element from which coherent radiation is emitted ...

$$V = \pi D \rho_{\max}^2$$

$\rho_{\max} = \gamma v / \omega$
[transversal range ...
... with large polarization]

Maximum energy of radiated photons limited by plasma frequency ...
[results from requiring $V \neq 0 \rightarrow \omega = \gamma \omega_p$]

$$E_{\max} = \gamma \hbar \omega_p$$

[X-Rays \rightarrow large γ !!]


Typical values: $\left. \begin{array}{l} \text{CH}_2: \hbar \omega_p = 20 \text{ eV}; \gamma = 10^3 \\ \text{[Air: } \hbar \omega_p = 0.7 \text{ eV]} \end{array} \right]$

$$D = 10 \mu\text{m}$$


[$d > D$: absorption dominates]

Identifying particles with transition radiation

Detection Principle:
[Electron-ID]



Detector Signal:




- Detector should be sensitive to $3 \leq E_\gamma \leq 30$ keV.
 - ✓ Gaseous detectors
- In gas $\sigma_{\text{photo effect}} \propto Z^5$
- Gases with high Z are required
 - ✓ e.g. Xenon ($Z=54$)


ATLAS Transition Radiation Tracker

Straw Tube Tracker
with interspace filled with foam

→ Tracking & transition radiation




Identifying particles with transition radiation



Interaction of photons with matter

Characteristic for interactions of photons with matter:

A single interaction removes photon from beam !



Possible Interactions

Photoelectric Effect
Compton Scattering
Pair Production

Rayleigh Scattering ($\gamma A \rightarrow \gamma A$; $A = \text{atom}$; coherent)
Thomson Scattering ($\gamma e \rightarrow \gamma e$; elastic scattering)
Photo Nuclear Absorption ($\gamma K \rightarrow pK/nK$)
Nuclear Resonance Scattering ($\gamma K \rightarrow K^* \rightarrow \gamma K$)
Delbruck Scattering ($\gamma K \rightarrow \gamma K$)
Hadron Pair production ($\gamma K \rightarrow h^+h^- K$)

$$dI = -\mu I dx$$

[μ : absorption coefficient]


depends on
 E, Z, ρ

→ Beer-Lambert law:

$$I(x) = I_0 e^{-\mu x}$$



with $\lambda = 1/\mu = 1/n\sigma$
[mean free path]

Interaction of photons with matter



Interaction of photons with matter

Photon Total Cross Sections



Pair production

Cross Section:
[for $E_\gamma \gg m_e c^2$]

$$\sigma_{\text{pair}} \approx \frac{7}{9} \underbrace{\left(4 \alpha r_e^2 Z^2 \ln \frac{183}{Z^{\frac{1}{3}}} \right)}_{A/N_A X_0}$$

[X_0 : radiation length]
[in cm or g/cm²]

Absorption coefficient:


$$\mu = n \sigma \quad [\text{with } n: \text{particle density}]$$

$$\mu = \rho \cdot N_A / A \sigma_{\text{pair}}$$

$$= 7/9 \frac{1}{X_0}$$

[where now X_0 is in cm]

$$I(x) = I_0 e^{-\mu x}$$



	ρ [g/cm ³]	X_0 [cm]
H ₂ [fl.]	0.071	865
C	2.27	18.8
Fe	7.87	1.76
Pb	11.35	0.56
Air	$1.2 \cdot 10^{-3}$	$30 \cdot 10^3$


Electromagnetic showers

Reminder:

Dominant processes
at high energies ...

Photons : Pair production

Electrons : Bremsstrahlung



Pair production:

$$\begin{aligned}\sigma_{\text{pair}} &\approx \frac{7}{9} \left(4 \alpha r_e^2 Z^2 \ln \frac{183}{Z^{1/3}} \right) \\ &= \frac{7}{9} \frac{A}{N_A X_0} \quad [X_0: \text{radiation length}] \\ &\quad \quad \quad [\text{in cm or g/cm}^2]\end{aligned}$$

Absorption
coefficient:

$$\mu = n\sigma = \rho \frac{N_A}{A} \cdot \sigma_{\text{pair}} = \frac{7}{9} \frac{\rho}{X_0}$$

Bremsstrahlung:

$$\frac{dE}{dx} = 4\alpha N_A \frac{Z^2}{A} r_e^2 \cdot E \ln \frac{183}{Z^{1/3}} = \frac{E}{X_0}$$

$$\rightarrow E = E_0 e^{-x/X_0}$$

After passage of one X_0 electron
has only $(1/e)^{\text{th}}$ of its primary energy ...
[i.e. 37%]

Electromagnetic showers

Further basics:


Critical Energy [see above]:

$$\left. \frac{dE}{dx}(E_c) \right|_{\text{Brems}} = \left. \frac{dE}{dx}(E_c) \right|_{\text{Ion}}$$

Approximations:

$$E_c^{\text{Gas}} = \frac{710 \text{ MeV}}{Z + 0.92} \quad \left[E_c^{\text{Sol/Liq}} = \frac{610 \text{ MeV}}{Z + 1.24} \right]$$

$$\left(\frac{dE}{dx} \right)_{\text{Brems}} / \left(\frac{dE}{dx} \right)_{\text{Ion}} \approx \frac{Z \cdot E}{800 \text{ MeV}}$$



with:

$$\left. \frac{dE}{dx} \right|_{\text{Brems}} = \frac{E}{X_0} \quad \& \quad \left. \frac{dE}{dx} \right|_{\text{Ion}} \approx \frac{E_c}{X_0} = \text{const.}$$

Transverse size of EM shower given by radiation length via Molière radius

[see also later]

$$R_M = \frac{21 \text{ MeV}}{E_c} X_0$$

R_M : Moliere radius
 E_c : Critical Energy [Rossi]
 X_0 : Radiation length

Electromagnetic showers



Typical values for X_0 , E_c and R_M of materials used in calorimeter

	X_0 [cm]	E_c [MeV]	R_M [cm]
Pb	0.56	7.2	1.6
Scintillator (Sz)	34.7	80	9.1
Fe	1.76	21	1.8
Ar (liquid)	14	31	9.5
BGO	1.12	10.1	2.3
Sz/Pb	3.1	12.6	5.2
PB glass (SF5)	2.4	11.8	4.3

A simple shower model

Simple shower model: [from Heitler]

Only two dominant interactions:
Pair production and Bremsstrahlung ...



Shower development governed by X_0 ...


After a distance X_0 electrons remain with
only $(1/e)^{\text{th}}$ of their primary energy ...

Photon produces e^+e^- -pair after $9/7X_0 \approx X_0$...

Assume:

$E > E_c$: no energy loss by ionization/excitation

$E < E_c$: energy loss only via ionization/excitation



Use
Simplification:

$$E_\gamma = E_e \approx E_0/2$$

[E_e loses half the energy]

$$E_e \approx E_0/2$$

[Energy shared by e^+/e^-]

... with initial particle energy E_0


A simple shower model

Simple shower model: [continued]

Shower characterized by:

- Number of particles in shower
- Location of shower maximum
- Longitudinal shower distribution
- Transverse shower distribution

Longitudinal components;
measured in radiation length ...
... use: $t = \frac{x}{X_0}$



Number of shower particles
after depth t :

$$N(t) = 2^t$$

Energy per particle
after depth t :

$$E = \frac{E_0}{N(t)} = E_0 \cdot 2^{-t}$$

$$\rightarrow t = \log_2(E_0/E)$$

Total number of shower particles
with energy E_1 :

$$N(E_0, E_1) = 2^{t_1} = 2^{\log_2(E_0/E_1)} = \frac{E_0}{E_1}$$

Number of shower particles
at shower maximum:

$$N(E_0, E_c) = N_{\max} = 2^{t_{\max}} = \frac{E_0}{E_c}$$

Shower maximum at:

$$t_{\max} \propto \ln(E_0/E_c) \propto E_0$$

A simple shower model

Simple shower model: [continued]

Longitudinal shower distribution increases only logarithmically with the primary energy of the incident particle ...

Some numbers: $E_c \approx 10$ MeV, $E_0 = 1$ GeV $\rightarrow t_{\max} = \ln 100 \approx 4.5$; $N_{\max} = 100$
 $E_0 = 100$ GeV $\rightarrow t_{\max} = \ln 10000 \approx 9.2$; $N_{\max} = 10000$

$$t_{\max} [X_0] \sim \ln \frac{E_0}{E_c}$$

EM shower longitudinal development

Longitudinal profile

Parametrization:
[Longo 1975]

$$\frac{dE}{dt} = E_0 t^\alpha e^{-\beta t}$$


α, β : free parameters

t^α : at small depth number of secondaries increases ...

$e^{-\beta t}$: at larger depth absorption dominates ...

Numbers for $E = 2$ GeV (approximate):

$$\alpha = 2, \beta = 0.5, t_{\max} = \alpha/\beta$$



More exact
[Longo 1985]

$$\frac{dE}{dt} = E_0 \cdot \beta \cdot \frac{(\beta t)^{\alpha-1} e^{-\beta t}}{\Gamma(\alpha)}$$

[Γ : Gamma function]


$$\rightarrow t_{\max} = \frac{\alpha - 1}{\beta} = \ln \left(\frac{E_0}{E_c} \right) + C_{e\gamma}$$

with:

$$C_{e\gamma} = -0.5 \quad [\gamma\text{-induced}]$$

$$C_{e\gamma} = -1.0 \quad [e\text{-induced}]$$

EM shower longitudinal development



*EGS = Electron Gamma Shower

EM shower transverse development

Transverse shower development ...

Opening angle

for bremsstrahlung and pair production

$$\langle \theta^2 \rangle \approx (m/E)^2 = 1/\gamma^2$$

Small contribution as $m_e/E_c = 0.05$

Multiple scattering

deflection angle in 2-dimensional plane ...


$$\langle \theta_k^2 \rangle = \sum_{m=1}^k \theta_m^2 = k \langle \theta^2 \rangle$$

$$\sqrt{\langle \theta^2 \rangle} \approx \frac{13.6 \text{ MeV}/c}{p} \sqrt{\frac{x}{X_0}} \quad [\beta = 1]$$

In 3-dimensions extra factor $\sqrt{2}$:

$$\sqrt{\langle \theta^2 \rangle_{3d}} \approx \frac{19.2 \text{ MeV}/c}{p} \sqrt{\frac{x}{X_0}} \quad [\beta = 1]$$

Multiple
coulomb scattering



Assuming the approximate range of electrons to be X_0 yields lateral extension: $R = \langle \theta \rangle \cdot X_0 \dots$

$$R_M = \langle \theta \rangle_{x=X_0} \cdot X_0 \approx \frac{21 \text{ MeV}}{E_C} X_0$$

Molière Radius;
characterizes lateral shower spread ...

EM shower transverse profile

Transverse profile

Parametrization:

$$\frac{dE}{dr} = \alpha e^{-r/R_M} + \beta e^{-r/\lambda_{\min}}$$

α, β : free parameters

R_M : Molière radius

λ_{\min} : range of low energetic photons ...

Inner part: coulomb scattering ...


Electrons and positrons move away from shower axis due to multiple scattering ...

Outer part: low energy photons ...


Photons (and electrons) produced in isotropic processes (Compton scattering, photo-electric effect) move away from shower axis; predominant beyond shower maximum, particularly in high-Z absorber media...

Shower gets wider at larger depth ...

energy deposit
[arbitrary unites]




EM shower profiles



Longitudinal and transversal shower profile for a 6 GeV electron in lead absorber ...

[left: linear scale; right: logarithmic scale]



EM showers in a nutshell

Radiation length:
$$X_0 = \frac{180A}{Z^2} \frac{\text{g}}{\text{cm}^2}$$

Critical energy:
$$E_c^{\text{Gas}} = \frac{710 \text{ MeV}}{Z + 0.92} \quad E_c^{\text{Sol/Liq}} = \frac{610 \text{ MeV}}{Z + 1.24}$$

Shower maximum:
$$t_{\text{max}} = \ln \frac{E}{E_c} - \begin{cases} 1.0 & e^- \text{ induced shower} \\ 0.5 & \gamma \text{ induced shower} \end{cases}$$


Longitudinal energy containment:
$$L(95\%) = t_{\text{max}} + 0.08Z + 9.6 [X_0]$$

Transverse Energy containment:
$$R(90\%) = R_M \quad R(95\%) = 2R_M \quad R_M = \frac{21 \text{ MeV}}{E_c} X_0$$

Hadronic showers

Shower development:

1. $p + \text{Nucleus} \rightarrow \text{Pions} + N^* + \dots$
2. Secondary particles ...
undergo further inelastic collisions until they fall below pion production threshold
3. Sequential decays ...
 $\pi^0 \rightarrow \gamma\gamma$: yields electromagnetic shower
 Fission fragments $\rightarrow \beta$ -decay, γ -decay
 Neutron capture \rightarrow fission
 Spallation ...



Mean number of secondaries: $\sim \ln E$

Typical transverse momentum: $p_t \sim 350 \text{ MeV}/c$

Substantial electromagnetic fraction

$$f_{em} \sim \ln E$$

[variations significant]

Cascade energy distribution:

[Example: 5 GeV proton in lead-scintillator calorimeter]

Ionization energy of charged particles (p, π, μ)	1980 MeV [40%]
Electromagnetic shower (π^0, η^0, e)	760 MeV [15%]
Neutrons	520 MeV [10%]
Photons from nuclear de-excitation	310 MeV [6%]
Non-detectable energy (nuclear binding, neutrinos)	1430 MeV [29%]
	<hr/>
	5000 MeV [29%]

Hadronic showers

Hadronic interaction:

Cross Section:

$$\sigma_{\text{tot}} = \sigma_{\text{el}} + \sigma_{\text{inel}}$$

For substantial energies
 σ_{inel} dominates:

$$\sigma_{\text{el}} \approx 10 \text{ mb}$$

$$\sigma_{\text{inel}} \propto A^{2/3} \text{ [geometrical cross section]}$$

$$\therefore \sigma_{\text{tot}} = \sigma_{\text{tot}}(pA) \approx \sigma_{\text{tot}}(pp) \cdot A^{2/3}$$

[σ_{tot} slightly grows with \sqrt{s}]

at high energies
 also diffractive contribution

Hadronic interaction length:


$$\lambda_{\text{int}} = \frac{1}{\sigma_{\text{tot}} \cdot n} = \frac{A}{\sigma_{pp} A^{2/3} \cdot N_A \rho} \sim A^{1/3} \quad \text{[for } \sqrt{s} \approx 1 - 100 \text{ GeV]}$$

$$\approx 35 \text{ g/cm}^2 \cdot A^{1/3}$$

which yields:

$$N(x) = N_0 \exp(-x/\lambda_{\text{int}})$$

Remark: In principle one should distinguish between collision length $\lambda_W \sim 1/\sigma_{\text{tot}}$ and interaction length $\lambda_{\text{int}} \sim 1/\sigma_{\text{inel}}$ where the latter considers inelastic processes only (absorption) ...



Total proton-proton cross section
 [similar for p+n in 1-100 GeV range]



Interaction length characterizes both,
 longitudinal and transverse profile of
 hadronic showers ...

Hadronic vs. EM showers

Comparison

hadronic vs. electromagnetic shower ...

[Simulated air showers]



Hadronic vs. EM showers

Hadronic vs. electromagnetic interaction length:

$$\left. \begin{aligned} X_0 &\sim \frac{A}{Z^2} \\ \lambda_{\text{int}} &\sim A^{1/3} \end{aligned} \right\} \rightarrow \frac{\lambda_{\text{int}}}{X_0} \sim A^{4/3}$$

$$\lambda_{\text{int}} \gg X_0$$

[$\lambda_{\text{int}}/X_0 > 30$ possible; see below]

Typical
Longitudinal size: $6 \dots 9 \lambda_{\text{int}}$ [EM: 15-20 X_0]
[95% containment]

Typical
Transverse size: one λ_{int} [EM: 2 R_M ; compact]
[95% containment]

Hadronic calorimeter need more depth
than electromagnetic calorimeter ...

Some numerical values for materials
typical used in hadron calorimeters

	λ_{int} [cm]	X_0 [cm]
Scint	79.4	42.2
LAr	83.7	14.0
Fe	16.8	1.76
Pb	17.1	0.56
U	10.5	0.32
C	38.1	18.8

Hadronic shower development

Hadronic shower development:
[estimate similar to e.m. case]

Depth (in units of λ_{int}):

$$t = \frac{x}{\lambda_{\text{int}}}$$

Energy in depth t :

$$E(t) = \frac{E}{\langle n \rangle^t} \quad \& \quad E(t_{\text{max}}) = E_{\text{thr}}$$

[with $E_{\text{thr}} \approx 290 \text{ MeV}$]

$$E_{\text{thr}} = \frac{E}{\langle n \rangle^{t_{\text{max}}}}$$

Shower maximum:

$$\langle n \rangle^{t_{\text{max}}} = \frac{E}{E_{\text{thr}}}$$

Number of particles lower by factor E_{thr}/E_c compared to e.m. shower ...

Intrinsic resolution: worse by factor $\sqrt{E_{\text{thr}}/E_c}$

$$t_{\text{max}} = \frac{\ln(E/E_{\text{thr}})}{\ln \langle n \rangle}$$

But:

Only rough estimate as ...

energy sharing between shower particles fluctuates strongly ...

part of the energy is not detectable (neutrinos, binding energy); partial compensation possible (n-capture & fission)

spatial distribution varies strongly; different range of e.g. π^\pm and π^0 ...

electromagnetic fraction, i.e. fraction of energy deposited by $\pi^0 \rightarrow \gamma\gamma$ increases with energy ...


$$f_{\text{em}} \approx f_{\pi^0} \sim \ln E / (1 \text{ GeV})$$

Explanation: charged hadron contribute to electromagnetic fraction via $\pi^+\pi^- \rightarrow \pi^0 n$; the opposite happens only rarely as π^0 travel only $0.2 \mu\text{m}$ before its decay ('one-way street') ...

At energies below 1 GeV hadrons lose their energy via ionization only ...

Thus: need Monte Carlo (GEISHA, CALOR, ...) to describe shower development correctly ...

Hadronic shower longitudinal development



Hadronic shower transverse development



Transverse shower profile

Typical transverse momenta of secondaries: $\langle p_t \rangle \simeq 350 \text{ MeV}/c \dots$

Lateral extend at shower maximum: $R_{95\%} \simeq \lambda_{\text{int}} \dots$

Electromagnetic component leads to relatively well-defined core: $R \simeq R_M \dots$

Exponential decay after shower maximum ...



Lateral profile for 300 GeV π^-

[target material ^{238}U]
[measured at depth $4 \lambda_{\text{int}}$]

More π^0 's and γ in core
Energetic neutrons and charged pions form a wider core
Thermal neutrons generate broad tail

Measurement from induced radioactivity:


^{99}Mo (fission): neutron induced ...
[energetic neutron component]

^{237}U : mainly produced via $^{238}\text{U}(\gamma, n)^{237}\text{U} \dots$
[electromagnetic component]

^{239}Np : from ^{239}U decay ...
[thermal neutrons]

Ordinate indicates decay rate of different radioactive nuclides ...

Detectors are imperfect...



- **Detection efficiency**

$$\varepsilon = \frac{N_{\text{detected}}}{N_{\text{incident}}} = M \cdot R \cdot D$$

- ✓ **M = P(entering active region)**
 - Upstream material, entrance windows, ...
- ✓ **R = P(generating signal)**
 - Interaction cross sections, response, fluctuations, ...
- ✓ **D = P(signal gets registered)**
 - Readout properties, thresholds, ...

- **Acceptance**

- ✓ **Instrumented/reactive region of the phase space (e.g. pseudorapidity, azimuthal angle, but also energy/momentum)**
 - dynamic range

Detectors are imperfect...

- **Dead Time**


- ✓ Delay between particle entrance in detector and signal acquisition, in which the detector is rendered inactive (detector + readout)
- ✓ Dead time efficiency

$$\varepsilon(T_d) = \frac{1}{N_{\text{detected}} T_d + 1}$$

- **Trigger**

- ✓ Detector cannot acquire signals continuously...

- **Response linearity**



- **Response resolution**

