

# SIX LOOP BETA FUNCTION IN $\varphi^4$ MODEL

Thursday 23<sup>rd</sup> July, 2015

M. Kompaniets

E. Panzer

# OVERVIEW

1. Status of the multiloop calculations in  $\varphi^4$  model
2. Two-point function
3. Four-point function
  - 3.1 IBP,  $R^*$
  - 3.2 Calculating graphs with subdivergences using HyperInt
  - 3.3 Preliminary results
4. Predictions for 6 loop term based on Kazakov, Tarasov, Shirkov (1979) paper
5. Borel resummation
6. Summary

STATUS OF THE MULTILoop  
CALCULATIONS IN  $\varphi^4$  MODEL

# STATUS OF THE MULTILoop CALCULATIONS IN $\varphi^4$ MODEL

$$S(\varphi) = - \int d\mathbf{x} \left( \frac{1}{2} \tau \varphi(\mathbf{x})^2 + \frac{1}{2} (\vec{\nabla} \varphi(\mathbf{x}))^2 + \frac{1}{24} g (\varphi(\mathbf{x})^2)^2 \right)$$

- critical exponents at 4-loop level:
  - **E. Brezin, J.C. LeGuillou and J. Zinn-Justin**, *Phys. Rev.*, D9 (1974) 1121.
  - **D.I. Kazakov, O.V. Tarasov and A.A. Vladimirov**, *Zh. Eksp. Teor. Fiz.*, 77 (1979) 1035.

---

<sup>1</sup>no subdivergences

# STATUS OF THE MULTILoop CALCULATIONS IN $\varphi^4$ MODEL

$$S(\varphi) = - \int d\mathbf{x} \left( \frac{1}{2} \tau \varphi(\mathbf{x})^2 + \frac{1}{2} (\vec{\nabla} \varphi(\mathbf{x}))^2 + \frac{1}{24} g (\varphi(\mathbf{x})^2)^2 \right)$$

- critical exponents at **4-loop level**:
  - **E. Brezin, J.C. LeGuillou and J. Zinn-Justin**, *Phys. Rev.*, D9 (1974) 1121.
  - **D.I. Kazakov, O.V. Tarasov and A.A. Vladimirov**, *Zh. Eksp. Teor. Fiz.*, 77 (1979) 1035.
- critical exponents at **5-loop level**
  - **K.G. Chetyrkin, A.L. Kataev, F.V. Tkachev** 1981 *Phys.Lett.* B 99 147; B 101 457(E)
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov**, 1983 *Phys. Lett.* B 132 351
  - **D.I. Kazakov** 1983 *Phys. Lett.* B 133 406; 1984 *Theor.Math.Phys.* 58 223-230
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov** 1986 *Preprint* INR P-0453, Moscow.
  - **H. Kleinert, J. Neu, V. Shulte-Frohlind, K.G. Chetyrkin, S.A. Larin** 1991 *Phys.Lett.* B 272 39; Erratum 1993 B 319, 545

---

<sup>1</sup>no subdivergences

# STATUS OF THE MULTILoop CALCULATIONS IN $\varphi^4$ MODEL

$$S(\varphi) = - \int d\mathbf{x} \left( \frac{1}{2} \tau \varphi(\mathbf{x})^2 + \frac{1}{2} (\vec{\nabla} \varphi(\mathbf{x}))^2 + \frac{1}{24} g (\varphi(\mathbf{x})^2)^2 \right)$$

- critical exponents at **4-loop level**:
  - **E. Brezin, J.C. LeGuillou and J. Zinn-Justin**, *Phys. Rev.*, D9 (1974) 1121.
  - **D.I. Kazakov, O.V. Tarasov and A.A. Vladimirov**, *Zh. Eksp. Teor. Fiz.*, 77 (1979) 1035.
- critical exponents at **5-loop level**
  - **K.G. Chetyrkin, A.L. Kataev, F.V. Tkachev** 1981 *Phys.Lett.* B 99 147; B 101 457(E)
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov**, 1983 *Phys. Lett.* B 132 351
  - **D.I. Kazakov** 1983 *Phys. Lett.* B 133 406; 1984 *Theor.Math.Phys.* 58 223-230
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov** 1986 *Preprint* INR P-0453, Moscow.
  - **H. Kleinert, J. Neu, V. Shulte-Frohlinde, K.G. Chetyrkin, S.A. Larin** 1991 *Phys.Lett.* B 272 39; Erratum 1993 B 319, 545
- critical exponent  $\eta$  (two-point function) at **6-loop level**
  - **D.V. Batkovich, K.G. Chetyrkin, M.V. Kompaniets** *in preparation*

---

<sup>1</sup>no subdivergences

# STATUS OF THE MULTILoop CALCULATIONS IN $\varphi^4$ MODEL

$$S(\varphi) = - \int d\mathbf{x} \left( \frac{1}{2} \tau \varphi(\mathbf{x})^2 + \frac{1}{2} (\vec{\nabla} \varphi(\mathbf{x}))^2 + \frac{1}{24} g (\varphi(\mathbf{x})^2)^2 \right)$$

- critical exponents at **4-loop level**:
  - **E. Brezin, J.C. LeGuillou and J. Zinn-Justin**, *Phys. Rev.*, D9 (1974) 1121.
  - **D.I. Kazakov, O.V. Tarasov and A.A. Vladimirov**, *Zh. Eksp. Teor. Fiz.*, 77 (1979) 1035.
- critical exponents at **5-loop level**
  - **K.G. Chetyrkin, A.L. Kataev, F.V. Tkachev** 1981 *Phys.Lett.* B 99 147; B 101 457(E)
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov**, 1983 *Phys. Lett.* B 132 351
  - **D.I. Kazakov** 1983 *Phys. Lett.* B 133 406; 1984 *Theor.Math.Phys.* 58 223-230
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov** 1986 *Preprint* INR P-0453, Moscow.
  - **H. Kleinert, J. Neu, V. Shulte-Frohlinde, K.G. Chetyrkin, S.A. Larin** 1991 *Phys.Lett.* B 272 39; Erratum 1993 B 319, 545
- critical exponent  $\eta$  (two-point function) at **6-loop level**
  - **D.V. Batkovich, K.G. Chetyrkin, M.V. Kompaniets** *in preparation*
- complete **6-loop** calculations of the anomalous dimensions of  $\varphi^4$  model (**this talk**)
  - **M.V. Kompaniets, E. Panzer** *in progress*

---

<sup>1</sup>no subdivergences

# STATUS OF THE MULTILoop CALCULATIONS IN $\varphi^4$ MODEL

$$S(\varphi) = - \int d\mathbf{x} \left( \frac{1}{2} \tau \varphi(\mathbf{x})^2 + \frac{1}{2} (\vec{\nabla} \varphi(\mathbf{x}))^2 + \frac{1}{24} g(\varphi(\mathbf{x})^2)^2 \right)$$

- critical exponents at **4-loop level**:
  - **E. Brezin, J.C. LeGuillou and J. Zinn-Justin**, *Phys. Rev.*, D9 (1974) 1121.
  - **D.I. Kazakov, O.V. Tarasov and A.A. Vladimirov**, *Zh. Eksp. Teor. Fiz.*, 77 (1979) 1035.
- critical exponents at **5-loop level**
  - **K.G. Chetyrkin, A.L. Kataev, F.V. Tkachev** 1981 *Phys.Lett.* B 99 147; B 101 457(E)
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov**, 1983 *Phys. Lett.* B 132 351
  - **D.I. Kazakov** 1983 *Phys. Lett.* B 133 406; 1984 *Theor.Math.Phys.* 58 223-230
  - **K.G. Chetyrkin, S.G. Gorishny, S.A. Larin and F.V. Tkachov** 1986 *Preprint* INR P-0453, Moscow.
  - **H. Kleinert, J. Neu, V. Shulte-Frohlinde, K.G. Chetyrkin, S.A. Larin** 1991 *Phys.Lett.* B 272 39; Erratum 1993 B 319, 545
- critical exponent  $\eta$  (two-point function) at **6-loop level**
  - **D.V. Batkovich, K.G. Chetyrkin, M.V. Kompaniets** *in preparation*
- complete **6-loop** calculations of the anomalous dimensions of  $\varphi^4$  model (**this talk**)
  - **M.V. Kompaniets, E. Panzer** *in progress*
- primitive<sup>1</sup> graphs up to **7-8 loops**
  - **D. J. Broadhurst and D. Kreimer**, *Int. J. Mod. Phys. C* 6 (Aug., 1995) 519–524 (**numerically, up to 7 loops**)
  - **O. Schnetz**, *Commun. Number Theory Phys.* 4 (2010), no. 1 1–47 (**analytically, up to 7 loops and almost all 8 loops**)

---

<sup>1</sup>no subdivergences



# TWO-POINT FUNCTION

# TWO-POINT FUNCTION

in collaboration with D.V. Batkovich and K.G. Chetyrkin


Number of graphs in  $\Gamma_2$   
(two-point 1PI Green function)

# of loops	1	2	3	4	5	6
# of graphs	0	1	1	4	11	<b>50</b>

Graph equals to  $G(1, 5\varepsilon)F(\varepsilon)$ , where  $G(1, 5\varepsilon) = -5/12 + \mathcal{O}(\varepsilon)$  and  $F(\varepsilon)$  is pole part of 5-loop subgraph.

It is not possible to calculate  $F(\varepsilon)$  using 4-loop IBP

Pole part of  $F(\varepsilon)$  can be reconstructed from the 5-loop counterterm.




## Theorem

(L)-loop counterterms can be calculated using  $R^*$  operation and (L-1)-loop IBP reduction

With 4-loop IBP reduction we can calculate **only 48 out of 50** 6-loop graphs

No IRR for this graph which allows to use 4-loop IBP

Calculated using transition to dual graph



## TWO-POINT FUNCTION, ANOMALOUS DIMENSION OF THE FIELD

$$\begin{aligned}\gamma_\varphi(u) = & \frac{u^2(n+2)}{36} - \left[8+n\right] \frac{u^3(n+2)}{432} + \left[500+90n-5n^2\right] \frac{u^4(n+2)}{5184} + \\ & + \left[-77056+8832\zeta_3-25344\zeta_4+(-22752+3072\zeta_3-5760\zeta_4)n+\right. \\ & + \left.(-296-288\zeta_3)n^2+(-39+48\zeta_3)n^3\right] \frac{u^5(n+2)}{186624} + \\ & + \left[1410544+297472\zeta_3+619776\zeta_4-833536\zeta_5-95232\zeta_3^2+1190400\zeta_6+\right. \\ & + \left.(549104+69888\zeta_3+215808\zeta_4-293632\zeta_5-28160\zeta_3^2+352000\zeta_6)n+\right. \\ & + \left.(30184+14976\zeta_3+15744\zeta_4-23680\zeta_5-1024\zeta_3^2+12800\zeta_6)n^2+\right. \\ & + \left.(-794+96\zeta_4)n^3+(-29-16\zeta_3+48\zeta_4)n^4\right] \frac{u^6(n+2)}{746496} + O(u^7)\end{aligned}$$

## TWO-POINT FUNCTION, ANOMALOUS DIMENSION OF THE FIELD

$$\begin{aligned}\gamma_\varphi(u) = & \frac{u^2(n+2)}{36} - \left[8+n\right] \frac{u^3(n+2)}{432} + \left[500+90n-5n^2\right] \frac{u^4(n+2)}{5184} + \\ & + \left[-77056+8832\zeta_3-25344\zeta_4+(-22752+3072\zeta_3-5760\zeta_4)n+\right. \\ & + \left.(-296-288\zeta_3)n^2+(-39+48\zeta_3)n^3\right] \frac{u^5(n+2)}{186624} + \\ & + \left[1410544+297472\zeta_3+619776\zeta_4-833536\zeta_5-95232\zeta_3^2+1190400\zeta_6+\right. \\ & + \left.(549104+69888\zeta_3+215808\zeta_4-293632\zeta_5-28160\zeta_3^2+352000\zeta_6)n+\right. \\ & + \left.(30184+14976\zeta_3+15744\zeta_4-23680\zeta_5-1024\zeta_3^2+12800\zeta_6)n^2+\right. \\ & + \left.(-794+96\zeta_4)n^3+(-29-16\zeta_3+48\zeta_4)n^4\right] \frac{u^6(n+2)}{746496} + O(u^7)\end{aligned}$$

No  $\zeta_7$  here

## TWO-POINT FUNCTION, CRITICAL EXPONENT $\eta$

$$N = 1, D = 4 - 2\varepsilon$$

$$\begin{aligned}\eta(\varepsilon) &= 2\gamma_\varphi(u_*) = \frac{2}{27}\varepsilon^2 + \frac{109}{729}\varepsilon^3 + \left(\frac{7217}{39366} - \frac{64}{243}\zeta_3\right)\varepsilon^4 + \\ &+ \left(\frac{321511}{2125764} - \frac{32}{81}\zeta_4 - \frac{1316}{2187}\zeta_3 + \frac{1280}{729}\zeta_5\right)\varepsilon^5 + \\ &+ \left(\frac{3421613}{38263752} - \frac{181462}{177147}\zeta_3 - \frac{658}{729}\zeta_4 + \frac{73232}{19683}\zeta_5 + \frac{2432}{2187}\zeta_3^2 + \frac{3200}{729}\zeta_6 - \frac{3136}{243}\zeta_7\right)\varepsilon^6 + O(\varepsilon^7) \\ &= 0.074074\varepsilon^2 + 0.149520\varepsilon^3 - 0.133260\varepsilon^4 + 0.821006\varepsilon^5 - \mathbf{5.201449}\varepsilon^6 + O(\varepsilon^7)\end{aligned}$$

$\zeta_7$  originates from  $u_*$  value

## TWO-POINT FUNCTION, COMPARISON WITH $1/N$ -EXPANSION

$1/N$ -expansion up to  $1/N^3$  using conformal bootstrap technique

A.N. Vasiliev, Yu.M. Pis'mak, J. Honkonen, *Theor. Math. Phys.*, 50,N 2, p127 (1982) <sup>2</sup>:

$$\eta_N = \frac{\eta_1(\varepsilon)}{N} + \frac{\eta_2(\varepsilon)}{N^2} + \frac{\eta_3(\varepsilon)}{N^3} + O\left(\frac{1}{N^4}\right) \quad \text{for any } \varepsilon,$$

We know  $\eta_N$  up to  $1/N^3$  for any  $\varepsilon$  and we know  $\eta_\varepsilon$  up to  $\varepsilon^6$  for any  $N$ .

$$\eta_N \text{ expansion by } \varepsilon = \eta_\varepsilon \text{ expansion by } 1/N$$

---

<sup>2</sup>This paper contains missprint in the second term of the r.h.s of eq. (22): the denominator must be  $3(2 - \mu)^3$  (for details see **Vasilev A. N.**, Quantum Field Renormalization Group in Critical Behavior Theory and Stochastic Dynamics, 2004 )

## TWO-POINT FUNCTION, COMPARISON WITH $1/N$ -EXPANSION

$1/N$ -expansion up to  $1/N^3$  using conformal bootstrap technique

A.N. Vasiliev, Yu.M. Pis'mak, J. Honkonen, *Theor. Math. Phys.*, 50,N 2, p127 (1982) <sup>2</sup>:

$$\eta_N = \frac{\eta_1(\varepsilon)}{N} + \frac{\eta_2(\varepsilon)}{N^2} + \frac{\eta_3(\varepsilon)}{N^3} + O\left(\frac{1}{N^4}\right) \quad \text{for any } \varepsilon,$$

We know  $\eta_N$  up to  $1/N^3$  for any  $\varepsilon$  and we know  $\eta_\varepsilon$  up to  $\varepsilon^6$  for any  $N$ .

$$\eta_N \text{ expansion by } \varepsilon = \eta_\varepsilon \text{ expansion by } 1/N$$

Three independent relations for 6 loop diagrams.

All of 6-loops self-energy diagrams give contribution to  $\eta_3$ .

---

<sup>2</sup>This paper contains missprint in the second term of the r.h.s of eq. (22): the denominator must be  $3(2 - \mu)^3$  (for details see **Vasilev A. N.**, Quantum Field Renormalization Group in Critical Behavior Theory and Stochastic Dynamics, 2004 )

## TWO-POINT FUNCTION, COMPARISON WITH $1/N$ -EXPANSION

$1/N$ -expansion up to  $1/N^3$  using conformal bootstrap technique

A.N. Vasiliev, Yu.M. Pis'mak, J. Honkonen, *Theor. Math. Phys.*, 50,N 2, p127 (1982) <sup>2</sup>:

$$\eta_N = \frac{\eta_1(\varepsilon)}{N} + \frac{\eta_2(\varepsilon)}{N^2} + \frac{\eta_3(\varepsilon)}{N^3} + O\left(\frac{1}{N^4}\right) \quad \text{for any } \varepsilon,$$

We know  $\eta_N$  up to  $1/N^3$  for any  $\varepsilon$  and we know  $\eta_\varepsilon$  up to  $\varepsilon^6$  for any  $N$ .

$$\eta_N \text{ expansion by } \varepsilon = \eta_\varepsilon \text{ expansion by } 1/N$$

Three independent relations for 6 loop diagrams.

All of 6-loops self-energy diagrams give contribution to  $\eta_3$ .

**We have indeed found FULL agreement to this 32 years old prediction!**

---

<sup>2</sup>This paper contains missprint in the second term of the r.h.s of eq. (22): the denominator must be  $3(2 - \mu)^3$  (for details see **Vasilev A. N.**, Quantum Field Renormalization Group in Critical Behavior Theory and Stochastic Dynamics, 2004 )



# FOUR-POINT FUNCTION

# FOUR-POINT FUNCTION

Number of graphs in  $\Gamma_4$   
(four-point 1PI Green function)

# of loops	1	2	3	4	5	6
# of graphs	1	2	8	26	124	627

- **factorized** – can be represented as product of two graphs  $\gamma = \gamma_1 \cdot \gamma_2$ ,  
 $KR'(\gamma) = -KR'(\gamma_1) \cdot KR'(\gamma_2)$
- **4-loop reducible** – graphs which contains 4-loop(or less) p-integral as subgraph and the rest part can be integrated using G-functions
- **primitive** – graphs with no subdivergences <sup>3</sup>
- **4-loop irreducible** – graphs cannot be calculated using 4-loop IBP and  $R^*$  <sup>4</sup>

factorized	124
4-loop reducible	481
primitive	10
4-loop irreducible	12

<sup>3</sup>O. Schnetz, *Commun. Number Theory Phys.* 4 (2010), no. 1 1–47, arXiv:0801.2856

<sup>4</sup>E. Panzer, *Comp. Phys. Comm.*, 188 (2015), pp. 148-166, arXiv:1403.3385

## 4-LOOP REDUCIBLE GRAPHS, IBP, $R^*$

- Infrared rearrangement(IRR) and  $R^*$ -operation implemented as *Python* library <sup>5</sup> on top of GraphState/Graphine library
- IBP reduction also implemented on *Python* with *mongodb* as cache. Reduction rules are generated by LiteRed<sup>6</sup>
- Master integral values are taken from Baikov/Chetyrkin paper<sup>7</sup> and Lee/Smirnov/Smirnov paper<sup>8</sup>  
  
Surprisingly that for 6-loop  $\Gamma_2$  graphs it is enough master integral values from Baikov/Chetyrkin paper, but for 6-loop  $\Gamma_4$  graphs one more term from Lee/Smirnov/Smirnov paper is required.
- Counterterms for graphs calculated with all possible rearrangements of external momenta (extensive consistency check for IBP and  $R^*$ )

---


<sup>5</sup>Batkovich D V, Kompaniets M V 2015 *J. Phys.: Conf. Ser.* 608 012068

<sup>6</sup>Lee R N, 2012 Presenting LiteRed: a tool for the loop integrals reduction (Preprint hep-ph/1212.2685)

<sup>7</sup>Baikov P A, Chetyrkin K G, 2010 *Nucl. Phys. B* 837 186

<sup>8</sup>Lee R N, Smirnov A V, Smirnov V A 2012 *Nucl. Phys. B* 856 95

# 4-LOOP IRREDUCIBLE GRAPHS



All graphs contain subdivergences (up to 4 subgraphs)

# HYPERINT (E.PANZER)

Many Feynman integrals are expressible via multiple polylogarithms (MPL)

$$\text{Li}_{n_1, \dots, n_d}(z_1, \dots, z_d) = \sum_{0 < k_1 < \dots < k_d} \frac{z_1^{k_1} \dots z_d^{k_d}}{k_1^{n_1} \dots k_d^{n_d}}$$

and there special values, like multiple zeta values (MZV)  $\zeta_{n_1, \dots, n_d} = \text{Li}_{n_1, \dots, n_d}(1, \dots, 1)$

MPL can be rewritten in terms of hyperlogarithms

$$G(\sigma_1, \dots, \sigma_w; z) = \int_0^z \frac{dz_1}{z_1 - \sigma_1} \int_0^{z_1} \frac{dz_2}{z_2 - \sigma_2} \dots \int_0^{z_{w-1}} \frac{dz_w}{z_w - \sigma_w}$$

Main idea: consider graph in alpha-representation and integrate out one variable after other

$$f_n = \int_0^\infty f_{n-1} d\alpha_n = \int_{(0, \infty)^n} f_0 d\alpha_1 \dots d\alpha_n, \quad f_0 = \frac{\psi^{sdd-D/2}}{\phi^{sdd}} \prod_e \alpha_e^{a_e-1}, \quad sdd = \sum_e a_e - DL/2$$

If on each step of integration function  $f_{n-1}$  can be represented in terms of hyperlogarithms in  $\alpha_n$  with rational prefactors

$$f_{n-1} = \sum_{\vec{\sigma}, \tau, k} \frac{G(\vec{\sigma}; \alpha_n)}{(\alpha_n - \tau)^k} \lambda_{\sigma, \tau, k}, \quad \text{with } \sigma \text{ and } \tau \text{ independent of } \alpha_n$$


graph is called *linearly reducible* and can be integrated analytically using HyperInt.

for details see **E.Panzer**, *Comp. Phys. Comm.*, 188 (2015), pp. 148-166,

**E.Panzer**, Feynman integrals and hyperlogarithms (PhD Thesis) arXiv:1506.07243 and references therein

# HYPERINT (E.PANZER), GRAPHS WITH SUBDIVERGENCES

- algorithm does not limited by some particular loop number
- some limitations on the graph combinatorial type (graph must be linearly reducible)  
At this moment all graphs we need to calculate are linearly reducible
- we need to pass to HyperInt integrand without subdivergences
- we need to find some combination of graphs which are free from subdivergences (on the integrand level), this combination except the target graph must contain graphs which are "easier" to calculate <sup>9</sup>



- For automated calculations we need automation for subtraction terms generation.  
(current implementation work only for graphs with 2-3 subdivergences)

---

<sup>9</sup>**Brown F., Kreimer D.** *Letters in Mathematical Physics* September 2013, Volume 103, Issue 9, pp 933-1007, arXiv:1112.1180

# PRELIMINARY RESULTS FOR BETA-FUNCTION

$N = 1, D = 4 - 2\varepsilon, \overline{MS}$ -scheme

$$\begin{aligned}
 \beta(u) &= u \left( -2\varepsilon + 3u - \frac{17}{3}u^2 + \left[ 145 + 96 \zeta_3 \right] \frac{u^3}{8} + \left[ -3499 - 3744 \zeta_3 + 864 \zeta_4 - 5760 \zeta_5 \right] \frac{u^4}{48} + \right. \\
 &+ \left[ 764621 + 1146960 \zeta_3 - 342432 \zeta_4 + 2274048 \zeta_5 + \right. \\
 &+ \left. 103680 \zeta_3^2 - 777600 \zeta_6 + 3048192 \zeta_7 \right] \frac{u^5}{2304} + \\
 &+ \left[ -94207135 - 187104720 \zeta_3 + 61160400 \zeta_4 - 367044480 \zeta_5 - 99970560 \zeta_3^2 + \right. \\
 &+ 192700800 \zeta_6 - 119771136 \zeta_{3,5} + 9331200 \zeta_3 \zeta_4 - 732983040 \zeta_7 - 270950400 \zeta_3 \zeta_5 + \\
 &+ \left. 609507072 \zeta_8 - 44236800 \zeta_3^3 - 903782400 \zeta_9 \right] \frac{u^6}{57600} + O(u^7) \Big) = \\
 &= u \left( -2\varepsilon + 3u - \frac{17}{3}u^2 + 32.54968u^3 - 271.60578u^4 + \right. \\
 &\quad \left. + 2848.56826u^5 - 35096.77397u^6 + O(u^7) \right)
 \end{aligned}$$

$$\zeta_{3,5} = \sum_{0 < n < m} \frac{1}{n^3 m^5} \approx 0.03770767298$$

PREDICTIONS FOR 6 LOOP TERM  
BASED ON KAZAKOV, TARASOV,  
SHIRKOV (1979) PAPER



# PREDICTIONS FOR 6 LOOP TERM

$$\bar{\beta}(u) = \frac{\beta(u) + 2\varepsilon u}{2} = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + 1424.28u^6 - 17548.38u^7 + O(u^8)$$

Borel transform: 
$$\beta^B(u) = \int_0^\infty \frac{dx}{u} e^{-x/u} \left(x \frac{\partial}{\partial x}\right)^b B(x), \quad B_n = \frac{\bar{\beta}_n}{n!n^b}$$

with consequent conformal mapping

$$B(x) = \sum_{n=2}^N x^n B_n \rightarrow B(x) = \left(\frac{x}{\omega}\right)^\nu \left(a_0 + a_1\omega + a_2\omega^2 + \dots + a_N\omega^N\right), \quad \omega(x) = \frac{\sqrt{1+x} - 1}{\sqrt{1+x} + 1}$$

coeffs  $a_i$  are fixed from the requirement to reproduce coefficients  $B_n$  known from loop expansion,  $\nu$  defines asymptotic of  $\beta(g)$  when  $g \rightarrow \infty$ , if asymptotic is not known  $\nu$  is defined from minimization of

$$\eta_{N+1} = 1 - \beta_{N+1}^B / \beta_N^B.$$

$$1.7 < \nu < 2.2$$

Expanding  $B(x)$  up to  $x^{N+1}$  one can get prediction for  $N + 1$  coefficient for beta function

for details see

**D. I. Kazakov, O. V. Tarasov, D. V. Shirkov**, (1979) "Analytic continuation of the results of perturbation theory for the model  $g\varphi^4$  to the region  $g \geq 1$ ", *Theoretical and Mathematical Physics*, Volume 38, Issue 1, pp 9-16

**Kazakov, D. I., Shirkov, D. V.** (1980), "Asymptotic Series of Quantum Field Theory and Their Summation." *Fortschr. Phys.*, 28: 465–499.

# PREDICTIONS FOR 6 LOOP TERM

$$\bar{\beta}(u) = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + 1424.28u^6 - 17548.38u^7 + O(u^8)$$

predictions based on 4,5,6 loop beta function ( $\nu = 2$ )<sup>10</sup>

$$\bar{\beta}^{P4}(u) = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + \mathbf{1404.30u^6} - \mathbf{16537.8u^7} + \mathbf{213452.8u^8} + O(u^9)$$

1.5%                      5.7%

---

<sup>10</sup>**D. I. Kazakov, O. V. Tarasov, D. V. Shirkov**, (1979), *Theoretical and Mathematical Physics*, Volume 38, Issue 1, pp 9-16

# PREDICTIONS FOR 6 LOOP TERM

$$\bar{\beta}(u) = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + 1424.28u^6 - 17548.38u^7 + O(u^8)$$

predictions based on 4,5,6 loop beta function ( $\nu = 2$ )<sup>11</sup>

$$\bar{\beta}^{P4}(u) = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + \mathbf{1404.30u^6} - \mathbf{16537.8u^7} + \mathbf{213452.8u^8} + O(u^9)$$


1.5%                      5.7%

$$\bar{\beta}^{P5}(u) = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + 1424.28u^6 - \mathbf{17150.62u^7} + \mathbf{226595.68u^8} + O(u^9)$$

2.3%!!

$$\bar{\beta}^{P6}(u) = \frac{3}{2}u^2 - \frac{17}{6}u^3 + 16.27u^4 - 135.80u^5 + 1424.28u^6 - 17548.38u^7 + \mathbf{242105.8u^8} + O(u^9)$$

prediction for 7 loop term



$\nu$  dependence of the relative error of the prediction for 6 loop term based on 3-6 loop beta function

<sup>11</sup>D. I. Kazakov, O. V. Tarasov, D. V. Shirkov, (1979), *Theoretical and Mathematical Physics*, Volume 38, Issue 1, pp 9-16

# PREDICTIONS FOR ANOMALOUS DIMENSION OF THE FIELD

$$\gamma_\varphi(u) = 0.0833u^2 - 0.0625u^3 + 0.3385u^4 - 1.9255u^5 + 14.383u^6 + O(u^7)$$

Predictions for  $\gamma_\varphi(u)$  based on 4,5,6 loop results ( $\nu = 3$ )

$$\gamma_\varphi^{P4}(u) = 0.0833u^2 - 0.0625u^3 + 0.3385u^4 - \underset{7\%}{2.0663u^5} + \underset{11\%}{16.068u^6} - 145.86u^7 + O(u^8)$$

# PREDICTIONS FOR ANOMALOUS DIMENSION OF THE FIELD

$$\gamma_\varphi(u) = 0.0833u^2 - 0.0625u^3 + 0.3385u^4 - 1.9255u^5 + 14.383u^6 + O(u^7)$$

Predictions for  $\gamma_\varphi(u)$  based on 4,5,6 loop results ( $\nu = 3$ )

$$\gamma_\varphi^{P4}(u) = 0.0833u^2 - 0.0625u^3 + 0.3385u^4 - \underset{7\%}{2.0663u^5} + \underset{11\%}{16.068u^6} - 145.86u^7 + O(u^8)$$

$$\gamma_\varphi^{P5}(u) = 0.0833u^2 - 0.0625u^3 + 0.3385u^4 - 1.9255u^5 + \underset{0.5\%!!}{14.316u^6} - 125.99u^7 + O(u^8)$$

$$\gamma_\varphi^{P6}(u) = 0.0833u^2 - 0.0625u^3 + 0.3385u^4 - 1.9255u^5 + 14.383u^6 - \underset{\text{prediction for 7 loop term}}{127.29u^7} + O(u^8)$$

# BOREL RESUMMATION

# ISING UNIVERSALITY CLASS ( $N = 1$ )

Critical exponents that can be measured in experiment

$$\alpha = \frac{2\Delta_\tau - d}{\Delta_\tau}, \quad \beta = \frac{\Delta_\phi}{\Delta_\tau}, \quad \gamma = \frac{d - 2\Delta_\phi}{\Delta_\tau}, \quad \delta = \frac{d - \Delta_\phi}{\Delta_\phi}, \quad \eta = 2\Delta_\phi - d + 2 = 2\gamma_\phi^* \quad (1)$$

where  $\Delta_\tau = 2 + \gamma_\tau^* = 1/\nu$ ,  $\Delta_\phi = d/2 - 1 + \gamma_\phi^*$

$D = 3$	$\alpha$	$\beta$	$\gamma$	$\delta$	$\nu$	$\eta$
$\epsilon$ -exp	0.1157	0.3255	1.2334	4.7899	0.6281	0.03629
HT & MC	0.110(1)	0.3265(3)	1.2372(5)	4.789(2)	0.6301	0.03601(4)
experiment	0.104-0.111	0.315-0.341	1.14-1.32		0.606-0.70	0.030-0.058

$D = 2$	$\alpha$	$\beta$	$\gamma$	$\delta$	$\nu$	$\eta$
$\epsilon$ -exp	0.1645	0.1092	1.6172	15.816	0.9178	0.2379
exact		0.125	1.75	15	1	0.25

HT, MC and experimental data are taken from **Pelissetto A., Vicari E.** "Critical Phenomena and Renormalization-Group Theory", *Phys.Rept.* 368:549-727,2002; arXiv:cond-mat/0012164

# SUMMARY



# SUMMARY

- 6 loop calculation for  $\varphi^4$  model are almost complete, beta function contains multiple zeta value  $\zeta_{3,5}$
- Automation for calculation of the graphs with subdivergences using HyperInt
- Additional crosschecks for beta-function using HyperInt
- Predictions of Kazakov/Tarasov/Shirkov for 6 loop terms gives only 2% error!
- Resumed critical exponents for 3D Ising model are in good agreement with both simulation and experimental data
- Resumed critical exponents for 2D Ising model are still far from exact solution