

Studying the Z boson resonance

– not with weak loops
– but with the SMATASY/ZFITTER-kit

Tord Riemann, Königs Wusterhausen, Germany

Thanks to: M. Grünewald + S. Riemann

talk held at “The High-Energy Physics KIT – HEPKIT2015”
5-8 October 2015, Karlsruhe Institute of Technology (KIT)

<https://indico.cern.ch/event/369827/overview>

Present interest in precision approaches to the Z boson

Belle-II

$\sqrt{s} \sim 10$ GeV

→ **Belle-II** will measure $10^9 \mu^+ \mu^-$ events

See e.g. T.Ferber [1]

Fcc-ee

$\sqrt{s} \sim M_Z$

→ **Fcc-ee** expects 10^{13} events at the Z resonance

See e.g. A. Freitas [2].

Much work on weak two-loop contributions to the Z resonance has been done by Hollik et al., Czakon et al., Freitas et al.

see [3] and many refs. therein.

Model-independent alternative: How to do?

→ Request by the Fcc-ee physics study group

→ **S-matrix approach** a la SMATASY/ZFITTER

See T.Riemann [4]

Outline

S-matrix approach to the Z line shape

- Developed as a model-independent analysis tool of $e^+e^- \rightarrow (\gamma, Z) \rightarrow f^+f^-$ **around the Z boson resonance**
- Aim: determinations of M_Z and Γ_Z
- $\rightarrow \sigma_T$: Leike/TR/Rose 1991 [5]
- $\rightarrow A_{FB,LR,pol}$: TR 1992 [6],
- \rightarrow SMATASY code: Kirsch/TR 1994 [7]
- First application: LEP/L3 1993 [8], also: Tristan/TOPAZ, VENUS, LEP/OPAL, ...

- 1 Introduction
- 2 Total cross sections
- 3 Asymmetries
- 4 Applications
- 5 SMATASY/ZFITTER
- 6 Summary

Introduction

The reaction

$$e^+e^- \rightarrow (\gamma, Z) \rightarrow f^+f^- + (n\gamma) \quad (1)$$

allows to study the Z boson, its mass M_Z , its width Γ_Z , its couplings, and potentially deviations from the Standard Model.

Need correct “model”

See experiences with *constant* and *s-dependent* Z width:

$$\frac{1}{[s - M_Z^2 + iM_Z \Gamma_Z(s)]} \quad \text{versus} \quad \frac{1}{[s - M_Z^2 + iM_Z \Gamma_Z]} \quad (2)$$

To a very good accuracy, it holds: $\Gamma_Z(s) \approx s/M_Z^2 \times \Gamma_Z$

see next slide, \rightarrow Bardin/Leike/Riemann/Schwartz 1988 [9]; also: Berends/Burgers/Hollik/v.Neerven 1988 [10]

Need correct unfolding ..

.. of *Realistic Observables* in order to get *Pseudo Observables*. \rightarrow e.g.: Borrelli/Consoli/Maiani/Sisto 1990 [11], Later: Bardin/Passarino 1999 [12], Bardin/Grünewald/Passarino 1999 [13], Passarino 2003 [14], Passarino 2013 [15] and refs. therein.

Lesson: The model influences numerical results




Fig. 1. Total cross sections σ_B , σ_{RC}^{QED} for the reactions $e^+e^- \rightarrow \mu^+\mu^- (\gamma)$ in Born approximation (left scale) and including $O(\alpha)$ QED corrections (right scale). Peaks of σ with energy-dependent width $\Gamma_Z(s)$ are shifted by 34 MeV to the left.

Total cross section for $e^+e^- \rightarrow \mu^+\mu^-$ production at LEP ...

... without (left) and with (right) QED corrections. Both sample data produced with an energy-dependent Z width.

The assumptions on the Z -propagator in the fit formulas influence the location of the peak, but not the “experimental errors”.

Fig.: from [9], license Number: 3557090997554.

	Born		Born + QED	
from fit: \rightarrow	M_Z	Γ_Z	M_Z	Γ_Z
$\Gamma_Z(s)$	93.000 \pm 0.013	2.498 \pm 0.009	93.000 \pm 0.016	2.498 \pm 0.011
Γ_Z	92.966 \pm 0.013	2.498 \pm 0.009	92.966 \pm 0.016	2.498 \pm 0.011

Introduction

Stuart 1991 [16], S-Matrix ansatz for $e^+e^- \rightarrow Z \rightarrow f^+f^-$

$$M = \frac{R}{s - s_0} + F(s), \quad s_0 = M_Z^2 - iM_Z\Gamma_Z \quad (3)$$

Allows to study:

- Mass M_Z and width $\Gamma_Z \rightarrow$ Leike/Riemann/Rose 1991 [5]
- How many independent degrees of freedom? \rightarrow Leike/Riemann/Rose 1991 [5], Kirsch/S.Riemann, L3 [17, 8]
- But also: How to define mass and width of the Z boson at higher orders of perturbation theory? \rightarrow Denner 2014 [18], Freitas 2014 [3] and Fcc-ee [2], Degrassi FCC-ee [19] and refs. therein

Total cross sections

There are immediate questions, from an experimental point:

- What about the photon exchange?
- What about QED corrections, e.g. the $2 \rightarrow 3$ part of the cross sections?
- What about asymmetries, besides σ_{tot} ?

We have to describe

$$e^+ e^- \longrightarrow (\gamma, Z) \longrightarrow f^+ f^- (\gamma) \quad (4)$$

Ansatz in the complex energy plane, for **four helicity matrix elements**:

$$\mathcal{M}^i(s) = \frac{R_\gamma^i}{s} + \frac{R_Z^i}{s - s_Z} + F^i(s), \quad i = 1, \dots, 4. \quad (5)$$

Beware: Eqn. (5) is mathematically not consistent → Böhm/Sato 2004 [20]

The poles of \mathcal{M} have complex residua R_Z and R_γ , the latter corresponding to the photon, and the background $F(s)$ is an analytic function without poles:

$$F^i(s) = \sum_{n=0}^{\infty} F_n^i (s - s_0)^n \quad (6)$$

Comment on the photon term (3 Feb 2015)

$$\begin{aligned}
 \frac{R_\gamma^i(s)}{s} &= \frac{\sum_{n=0}^{\infty} R_n^i(s-s_0)^n}{s} & (7) \\
 &= \frac{\sum_{n=0}^{\infty} R_n^i(s-s_0)^n}{s_0 - (s_0 - s)} \\
 &= \sum_{n=0}^{\infty} R_n^i(s-s_0)^n \frac{1}{s_0} \frac{1}{1 - \frac{s_0-s}{s_0}} \\
 &= \sum_{n=0}^{\infty} R_n^i(s-s_0)^n \frac{1}{s_0} \left[1 + \frac{s_0-s}{s_0} + \left(\frac{s_0-s}{s_0} \right)^2 \cdots \right]
 \end{aligned}$$

The term $R_\gamma^i(s)/s$ is part of the the background term $F(s)$.

- *It is useful to sum up a selected part of the photonic background of the Z resonance in order to take explicit notice of physically known pieces of the input expressions.*
- *Compare: It is useful to sum up a selected part of self-energy insertions in the propagators in order to derive the Breit-Wigner resonance form.*

Ansatz for realistic applications

The analysis of the Z line shape will be based here on the cross section

$$\sigma(s) = \sum_{i=1}^4 \sigma^i(s) = \frac{1}{4} \sum_{i=1}^4 s |\mathcal{M}^i(s)|^2, \quad (8)$$

where the sum must be performed over four helicity amplitudes with different residues R_Z^i and functions $F^i(s)$. The result is, with QED corrections folded in:

$$\sigma_T(s) = \frac{4}{3} \pi \alpha^2 \int \frac{ds'}{s} \left[\frac{r^\gamma}{s} + \frac{sR + (s - M_Z^2)J}{(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} + \dots \right] \rho_{ini} \left(\frac{s'}{s} \right). \quad (9)$$

The radiation connected with initial-final state interferences can be taken into account by an analogue formula to (9) with a slightly more complicated structure [21, 22]:

$$\sigma_{\text{int}}(s) = \int ds' \sigma(s, s') \rho_{\text{int}}(s'/s). \quad (10)$$

Some details

See [\[23\]](#)

ZFITTER

The Standard Model analysis tool for the Z resonance: ZFITTER (D. Bardin et al.)

- Complete electroweak radiative corrections;
- QED corrections by convolution:
with some $\sigma_0(s')$
beware: for initial-final state interferences with some $\sigma_0(s, s')$;
- **semi-analytical QED** integrations;
- free choice of $\sigma_0(s')$ by **user interfaces**;
- Standard Model interfaces: four weak form factors $\rho, \kappa_e, \kappa_f, \kappa_{ef}$.

ZFITTER is **well-tested**, **flexible**, **accurate** and **fast**.

References: CPC in 2001 [24] and in 2006 [25], also: CPC in 1990 [26]

- The actual Fortran package is v.6.44; version 6.42 is public in CPC program library, with CPC-licence.
- Beware: **Gfitter/GSM (2007-2011) is an illegal clone of ZFITTER**. The code is available at <http://zfitter-gfitter.desy.de/> and <http://fh.desy.de/projekte/gfitter01/Gfitter01.htm>.
For ZFITTER, see also: <http://sanc.jinr.ru/users/zfitter>

catalogue numbers: **ADMJ_v1_0 (2001), ADMJ_v2_0 (2006)**
<http://www.cpc.cs.qub.ac.uk/licence/licence.html>



Computer Physics Communications Program Library



Programs in Physics & Physical Chemistry

Disclaimer and Non-profit use Licence Agreement

The CPC non-profit use licence agreement is an agreement between the author(s) of a program distributed by the CPC Program Library and the person who acquires it. By acquiring the program the person is agreeing to be bound by the terms of this agreement.

1. This licence entitles the licensee (one person) and the licensee's research group to obtain a copy of the source or executable code and to use the acquired program for academic or non-profit use within a research group; or, it entitles the licensee (one company, organisation or computing centre) to install the program and allow access to the executable code to members of the licensee's organisation for academic or non-profit use. No user or site will re-distribute the source code or executable code to a third party in original or modified form without the written permission of the author.
2. Publications which result from using the acquired program will reference the article in Computer Physics Communications which describes the program.
3. This licence does not permit any commercial (profit-making or proprietary) use or re-licensing or re-distributions. Persons interested in for-profit use should contact the author.
4. To the extent permissible under applicable laws, no responsibility is assumed and is hereby disclaimed by Elsevier for any injury and/or damage to persons or property as a result of any actual or alleged libelous statements, infringement of intellectual property or privacy rights, or products liability, whether resulting from negligence or otherwise, including without limitation from any use or operation of any ideas, instructions, procedures, products or methods contained in the material therein. Access to this site is provided on an "as is" basis, and Elsevier does not warrant that the information or software contained herein is complete or accurate or free from error.

This information or software is provided by its creators or authors as a service to Elsevier subscribers on an "as is" basis, and if downloaded by the subscriber should be checked for defects or viruses before being used. Unless noted otherwise, the creators or authors retain copyright and other proprietary rights.

Born Asymmetries

On a Sunday in Summer 1992, I had a discussion with Luciano Maiani in the CERN library. He had doubt that, **for asymmetries**, an analogue to the model-independent ansatz for σ_{tot} might be usefully formulated, especially in view of the QED corrections.

I believed one can do that, and I followed the rule “The proof of the pudding is in the eating” [6]. The result:

$$A^{Born}(s) = A_0 + A_1 \left(\frac{s}{M_Z^2} - 1 \right) + A_2 \left(\frac{s}{M_Z^2} - 1 \right)^2 + \dots \quad (11)$$

$$A_{FB} = \frac{\sigma_{FB}}{\sigma_T}, \quad A_{pol} = \frac{\sigma_{pol}}{\sigma_T}. \quad (12)$$

The A_{FB} and A_{pol} are helicity combinations as also σ_T is, i.e. have also S-matrix ansatzes.

The parameters in $A^{Born}(s)$ are in [QED-]Born approximation:

$$A_0 = \frac{R_A}{R_T}, \quad (13)$$

and

$$A_1 = \left[\frac{J_A}{R_A} - \frac{J_T}{R_T} \right] A_0. \quad (14)$$

Some details

See [23]

QED corrections for asymmetries

QED corrections to asymmetries lead to few simple correction factors [6, 7]:

$$A_{LR}^{QED}(s) = A_{0,LR}^{Born} + c_{1,T}(s) A_{1,LR}^{Born} \left(\frac{s}{M_Z^2} - 1 \right) + \dots \quad (15)$$

$$A_{FB}^{QED}(s) = c_{0,FB}(s) A_{0,FB}^{Born} + c_{1,FB}(s) A_{1,FB}^{Born} \left(\frac{s}{M_Z^2} - 1 \right) + \dots \quad (16)$$

The A_0 and A_1 are constant, and the same as in Born approximation.
The QED corrections are contained in the model-independent factor $C(s)$.

$$c_{0,FB}(s) = \frac{C_{FB}^R}{C_T^R}, \quad c_{0,T}(s) = 1 \quad (17)$$

$$c_{1,A}(s) = c_{0,A} \frac{C_T^J}{C_T^R} \quad (18)$$

Sample QED factor

$$C_{T,FB}^R(s) = \int dk \rho_{T,FB}(s'/s) \frac{s'R}{sR} \frac{(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2}{(s' - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \quad (19)$$

Sketch of derivation of the expression for $A_{FB}(s)$ with QED corr's

See [23]

QED corrections for asymmetries




Figure 1 : The forward-backward asymmetry for the process $e^+e^- \rightarrow \mu^+\mu^-$ near the Z boson peak. From Kirsch/Riemann 1994 [7], license Number: 3557090997554.

Applications

In Leike/S.Riemann/Riemann 1992 [27] correlations are discussed.

For the Z **peak position** s_{peak} , one may derive the relation:

$$\Delta\sqrt{s_{peak}} = \Delta M_Z + \frac{1}{4} \frac{\Gamma_Z^2}{M_Z} \Delta \left(\frac{J_T}{R_T} \right) + \dots \quad (20)$$

between an uncertainty in M_Z and an uncertainty in the γZ interference. The latter also influences A_1 .

Similarly, for a **hypothetical heavy gauge boson** Z' , the effects from its virtual exchange transform after a partial fraction decomposition into simple shifts of the γZ interferences [27]:

$$\Delta \left(\frac{J_T}{R_T} \right) = -2 \frac{g'^2}{g^2} \frac{M_{Z'}^2}{M_{Z'}^2 - M_Z^2} \frac{(a_e a'_e + v_e v'_e)(a_f a'_f + v_f v'_f)}{(a_e^2 + v_e^2)(a_f^2 + v_f^2)}, \quad (21)$$

Correlations: L3 at LEP1, 1993

Table 1

Results of the S matrix fit to total cross-sections and forward-backward asymmetries: (a) all parameters except the photon exchange are left free; (b) in addition the γZ interference terms are fixed to the Standard Model expectation.

Parameter	Case (a)	Case (b)
\overline{m}_Z (GeV)	91.152 ± 0.015	91.160 ± 0.010
$\overline{\Gamma}_Z$ (GeV)	2.494 ± 0.012	2.492 ± 0.012
$r_{\text{tot}}^{\text{lep}}$	0.141 ± 0.002	0.140 ± 0.002
$j_{\text{tot}}^{\text{lep}}$	0.032 ± 0.064	fixed to 0.0058
$r_{\text{fb}}^{\text{lep}}$	0.004 ± 0.001	0.004 ± 0.001
$j_{\text{fb}}^{\text{lep}}$	0.674 ± 0.087	0.675 ± 0.087
$r_{\text{tot}}^{\text{had}}$	2.859 ± 0.030	2.855 ± 0.029
$j_{\text{tot}}^{\text{had}}$	0.720 ± 0.700	fixed to 0.219

Table 2

Results of the S matrix fit to total cross-sections, forward-backward asymmetries and τ polarization: (a) all parameters except the photon exchange are left free; (b) in addition the hadronic γZ interference terms for the total cross-section are fixed to the Standard Model expectation.

Parameter	Case (a)	Case (b)
\overline{m}_Z (GeV)	91.155 ± 0.013	91.160 ± 0.010
$\overline{\Gamma}_Z$ (GeV)	2.494 ± 0.012	2.492 ± 0.012
$R_Z^{\text{lep}0}$	0.429 ± 0.012	0.429 ± 0.012
$R_Z^{\text{lep}1}$	-0.370 ± 0.003	-0.370 ± 0.003
$R_Z^{\text{lep}2}$	0.323 ± 0.016	0.323 ± 0.016
$r_{\text{tot}}^{\text{had}}$	2.860 ± 0.030	2.856 ± 0.029
$j_{\text{tot}}^{\text{had}}$	0.620 ± 0.620	fixed to 0.219
m_Z (GeV)	91.189 ± 0.013	91.194 ± 0.010
Γ_Z (GeV)	2.495 ± 0.012	2.493 ± 0.012
$\widehat{g}_n^{\text{lep}}$	-0.037 ± 0.010	-0.037 ± 0.010
$\widehat{g}_a^{\text{lep}}$	-0.4991 ± 0.0019	-0.4988 ± 0.0019
$\sin^2 \widehat{\theta}_W$	0.2317 ± 0.0037	0.2316 ± 0.0037

From: L3-Collaboration, "An S matrix analysis of the Z resonance", Phys. Lett. B315 (1993) 494-502.

See also **LEPEWWG et al., Phys. Rept. 2006**, section 2 [28] and **LEPEWWG et al., Phys. Rept. 2013**, App. A [29]

Correlations: LEP1 + LEP2

A complete analysis of the LEP-2 data in terms of J_{had}^{tot} is lacking.
But see K. Sachs, 2003 [30].

Including more measurements from LEP1 solves this problem, reducing the correlation. The final result of $M_Z = 91\,186.9 \pm 2.3 \text{ MeV}$ ⁸ is in very good agreement with the result of the standard lineshape fit $M_Z = 91\,187.6 \pm 2.1 \text{ MeV}$ ⁹ with only slightly increased error.




Figure 2: Correlation between the mass of the Z and J_{had}^{tot} . Results are shown for LEP1 data only and for a combined fit to LEP1 and LEP2 data. The yellow band indicates the 1σ error from the 9 parameter fit.

Figure 2 : K. Sachs, “Standard model at LEP II”, talk held at Moriond 2001, fig. 2 [30]

Correlations: LEP1 + TRISTAN

LEP experiments use cross-section and forward-backward asymmetry results from $\sqrt{s} \sim M_Z$ and LEP II. OPAL and L3 have reported preliminary results which are given in Table 1, and are compared to the value obtained by VENUS [9] using data at $\sqrt{s} \sim 60$ GeV and preliminary LEP I S-Matrix results. The results are consistent with each other, and with the SM prediction $j_{had}^{tot} = 0.22$.

Expt	Data	j_{had}^{tot}
L3:	LEP I + LEP II	0.30 ± 0.10
OPAL:	LEP I + LEP II	0.21 ± 0.12
VENUS:	VENUS + LEP I	0.20 ± 0.08

Table 1: Measurements of j_{had}^{tot}

Figure 3 : P. Holt, "Fermion pair production above the Z^0 resonance", talk held at HEP 2001, table 1 [31]

Fortran programs: ZPOLE - ZUSMAT - SMATASY/ZFITTER

ZPOLE – The stand-alone Fortran test package (Leike/Riemann, v.0.5, July 1991) is available on request.

It was used for the numerics of [5].

ZUSMAT ... the S-Matrix interface of older ZFITTER versions.

ZUSMAT was used for analysing the total cross sections, but could not treat asymmetries.

SMATASY/ZFITTER – With interface package **SMATASY** one has the full functionality of **ZFITTER** corrections [24, 25, 32].

The actual Fortran program for the S-matrix Z line shape approach:

M. Grünewald, S. Kirsch, T. Riemann 1994 [7]

SMATASY v.6.42.01 = SMATA642 (2 June 2005)

available at <https://gruenew.web.cern.ch/gruenew/smatasy.html>

ADBG_v1_0 (1995) <http://www.cpc.cs.qub.ac.uk/licence/licence.html>



Computer Physics Communications Program Library



Programs in Physics & Physical Chemistry

Disclaimer and Non-profit use Licence Agreement

The CPC non-profit use licence agreement is an agreement between the author(s) of a program distributed by the CPC Program Library and the person who acquires it. By acquiring the program the person is agreeing to be bound by the terms of this agreement.

1. This licence entitles the licensee (one person) and the licensee's research group to obtain a copy of the source or executable code and to use the acquired program for academic or non-profit use within a research group; or, it entitles the licensee (one company, organisation or computing centre) to install the program and allow access to the executable code to members of the licensee's organisation for academic or non-profit use. No user or site will re-distribute the source code or executable code to a third party in original or modified form without the written permission of the author.
2. Publications which result from using the acquired program will reference the article in Computer Physics Communications which describes the program.
3. This licence does not permit any commercial (profit-making or proprietary) use or re-licensing or re-distributions. Persons interested in for-profit use should contact the author.
4. To the extent permissible under applicable laws, no responsibility is assumed and is hereby disclaimed by Elsevier for any injury and/or damage to persons or property as a result of any actual or alleged libelous statements, infringement of intellectual property or privacy rights, or products liability, whether resulting from negligence or otherwise, including without limitation from any use or operation of any ideas, instructions, procedures, products or methods contained in the material therein. Access to this site is provided on an "as is" basis, and Elsevier does not warrant that the information or software contained herein is complete or accurate or free from error.
This information or software is provided by its creators or authors as a service to Elsevier subscribers on an "as is" basis, and if downloaded by the subscriber should be checked for defects or viruses before being used. Unless noted otherwise, the creators or authors retain copyright and other proprietary rights.

Good Scientific Practice – when using software ...

There is practically no literature on that. Software is never mentioned.

The general rules of international academic basic research apply. They include:

- Attribution of a scientific achievement to those who made it.
- Often by a proper citation.
- Other proper attributions are possible.

A proper attribution informs on:

- What was done?
- Who did it?
- How important is it?

In case of ethical or legal problems, questions have to be answered, in this order:

- On facts: What are the initial facts?
- On rules: Are there rules, are they violated?
- On sanctions: in case. Are there sanctions foreseen? By whom?

Reference [33]:

F. Carminati (CERN), D. Perret-Gallix (LAPP), T. Riemann (DESY)

“Summary of the ACAT 2013 round table discussion: Open-source, knowledge sharing and scientific collaboration”

arXiv.1407.0540, DESY-14-079, in proceedings of ACAT 2013

Summary

- The S-matrix approach is absolutely independent of the Standard Model approach.
- The degrees of freedom for σ_{tot} are, at minimum:

M_Z

Γ_Z

R – the residue of the Z resonance, *per scattering channel*

J – the value of the γZ interference, *per scattering channel*

- So we have at least **four degrees of freedom**.
This deserves at least **five data points** as a function of s .

See also: M.Grünewald, S.Kirsch 1993 [34].

- **Asymmetries** may be described as well as σ_{tot} .
- For an exact numerical analysis of data, an **accurate description of QED** corrections is mandatory.
This has been realised by combining SMATASY with ZFITTER.
- With so much more statistics at the Fcc-ee compared to LEP-1 and LEP-2:

The S-matrix approach might gain at the Fcc-ee even more interest as an alternative to the Standard Model approach.

References |

- [1] T. Ferber, Towards First Physics at Belle II. Talk at Spring Conference of DPG, 9-13 March 2015, Wuppertal, Germany.
http://www.staff.uni-giessen.de/~gd1472/belle/dpg2015_torbenferber.pdf.
- [2] A. Freitas, About projected theory uncertainties. Talk at the 2015 Pisa Fcc-ee meeting,
<https://agenda.infn.it/conferenceOtherViews.py?view=standard&confId=8830>.
- [3] A. Freitas, Higher-order electroweak corrections to the partial widths and branching ratios of the Z boson, JHEP 1404 (2014) 070.
[arXiv:1401.2447](https://arxiv.org/abs/1401.2447), [doi:10.1007/JHEP04\(2014\)070](https://doi.org/10.1007/JHEP04(2014)070).
- [4] T. Riemann, S-matrix approach to the Z line shape - A reminiscence. Prospects? Talk at the 2015 Pisa Fcc-ee meeting,
<https://agenda.infn.it/getFile.py/access?contribId=6&sessionId=7&resId=0&materialId=slides&confId=8830>.
- [5] A. Leike, T. Riemann, J. Rose, S matrix approach to the Z line shape, Phys. Lett. B273 (1991) 513–518.
[arXiv:hep-ph/9508390](https://arxiv.org/abs/hep-ph/9508390), [doi:10.1016/0370-2693\(91\)90307-C](https://doi.org/10.1016/0370-2693(91)90307-C).
- [6] T. Riemann, Cross-section asymmetries around the Z peak, Phys. Lett. B293 (1992) 451–456.
[arXiv:hep-ph/9506382](https://arxiv.org/abs/hep-ph/9506382), [doi:10.1016/0370-2693\(92\)90911-M](https://doi.org/10.1016/0370-2693(92)90911-M).
- [7] S. Kirsch, T. Riemann, SMATASY: A program for the model independent description of the Z resonance, Comput. Phys. Commun. 88 (1995) 89–108.
[arXiv:hep-ph/9408365](https://arxiv.org/abs/hep-ph/9408365), [doi:10.1016/0010-4655\(95\)00016-9](https://doi.org/10.1016/0010-4655(95)00016-9).
- [8] L3 collab., O. Adriani, et al., An S matrix analysis of the Z resonance, Phys. Lett. B315 (1993) 494–502.
[doi:10.1016/0370-2693\(93\)91646-5](https://doi.org/10.1016/0370-2693(93)91646-5).
- [9] D. Y. Bardin, A. Leike, T. Riemann, M. Sachwitz, Energy Dependent Width Effects in e^+e^- Annihilation Near the Z Boson Pole, Phys. Lett. B206 (1988) 539–542.
[doi:10.1016/0370-2693\(88\)91625-5](https://doi.org/10.1016/0370-2693(88)91625-5).
- [10] F. A. Berends, G. Burgers, W. Hollik, W. van Neerven, The Standard Z Peak, Phys. Lett. B203 (1988) 177.
[doi:10.1016/0370-2693\(88\)91593-6](https://doi.org/10.1016/0370-2693(88)91593-6).
- [11] A. Borrelli, M. Consoli, L. Maiani, R. Sisto, Model Independent Analysis of the Z Line Shape in e^+e^- Annihilation, Nucl. Phys. B333 (1990) 357.
[doi:10.1016/0550-3213\(90\)90042-C](https://doi.org/10.1016/0550-3213(90)90042-C).

References II

- [12] D. Y. Bardin, G. Passarino, The standard model in the making: Precision study of the electroweak interactions, International series of monographs on physics, 104 (Oxford University Press, 1999).
http://www.amazon.de/Standard-Model-Making-Interactions-International/dp/019850280X/ref=sr_1_1?ie=UTF8&qid=1422902184&sr=8-1&keywords=bardin+passarino.
- [13] D. Bardin, M. Grünewald, G. Passarino, Precision calculation project report.
[arXiv:hep-ph/9902452](http://arxiv.org/abs/hep-ph/9902452).
- [14] G. Passarino, Pseudo versus realistic observables: All that theories can tell us is how the world could be, talk at 'Workshop on Electroweak Precision Data and the Higgs Mass', DESY, Zeuthen, Feb. 28 - March 1, 2003 137–146.
<http://www--library.desy.de/preparch/desy/proc//proc03--01/14.ps.gz>.
- [15] G. Passarino, Higgs CAT, Eur. Phys. J. C74 (2014) 2866.
[arXiv:1312.2397](http://arxiv.org/abs/1312.2397), doi:10.1140/epjc/s10052-014-2866-7.
- [16] R. G. Stuart, Gauge invariance, analyticity and physical observables at the Z^0 resonance, Phys. Lett. B262 (1991) 113–119.
[doi:10.1016/0370-2693\(91\)90653-8](https://doi.org/10.1016/0370-2693(91)90653-8).
- [17] S. Kirsch, S. Riemann, A Combined Fit to the L3 Data Using the S-Matrix Approach (First Results) , L3 note 1233, 1992.
<http://13.web.cern.ch/13/note/notes1992.html>.
- [18] A. Denner, J.-N. Lang, The Complex-Mass Scheme and Unitarity in perturbative Quantum Field Theory, Eur. Phys. J. C75 (8) (2015) 377.
[arXiv:1406.6280](http://arxiv.org/abs/1406.6280), doi:10.1140/epjc/s10052-015-3579-2.
- [19] G. Degrossi, Precision observables in the Standard Model: a reexamination. Talk at the 2015 Pisa Fcc-ee meeting,
<https://agenda.infn.it/conferenceOtherViews.py?view=standard&confId=8830>.
- [20] A. R. Böhm, Y. Sato, Relativistic resonances: Their masses, widths, lifetimes, superposition, and causal evolution, Phys. Rev. D71 (2005) 085018.
[arXiv:hep-ph/0412106](http://arxiv.org/abs/hep-ph/0412106), doi:10.1103/PhysRevD.71.085018.
- [21] D. Y. Bardin, M. S. Bilenky, A. Chizhov, A. Sazonov, O. Fedorenko, T. Riemann, M. Sachwitz, Analytic approach to the complete set of QED corrections to fermion pair production in e^+e^- annihilation, Nucl. Phys. B351 (1991) 1–48.
[arXiv:hep-ph/9801208](http://arxiv.org/abs/hep-ph/9801208), doi:10.1016/0550-3213(91)90080-H.

References III

- [22] D. Y. Bardin, M. S. Bilenky, A. Chizhov, A. Sazonov, Y. Sedykh, T. Riemann, M. Sachwitz, The convolution integral for the forward - backward asymmetry in e^+e^- annihilation, Phys. Lett. B229 (1989) 405.
doi:[10.1016/0370-2693\(89\)90428-0](https://doi.org/10.1016/0370-2693(89)90428-0).
- [23] T. Riemann, S-matrix approach to the Z line shape. Talk at CALC2015, JINR, Dubna, Russia, July 2015,
https://indico.cern.ch/event/368497/session/7/contribution/59/attachments/1134075/1622013/02_riemann-tord-calc2015.pdf.
- [24] D. Bardin, M. Bilenky, P. Christova, M. Jack, L. Kalinovskaya, A. Olchevski, S. Riemann, T. Riemann, ZFITTER v.6.21: A semi-analytical program for fermion pair production in e^+e^- annihilation, Comput. Phys. Commun. 133 (2001) 229–395.
arXiv:[hep-ph/9908433](https://arxiv.org/abs/hep-ph/9908433), doi:[10.1016/S0010-4655\(00\)00152-1](https://doi.org/10.1016/S0010-4655(00)00152-1).
- [25] A. Arbuzov, M. Awramik, M. Czakon, A. Freitas, M. Grünewald, K. Mönig, S. Riemann, T. Riemann, ZFITTER: A Semi-analytical program for fermion pair production in e^+e^- annihilation, from version 6.21 to version 6.42, Comput. Phys. Commun. 174 (2006) 728–758.
arXiv:[hep-ph/0507146](https://arxiv.org/abs/hep-ph/0507146), doi:[10.1016/j.cpc.2005.12.009](https://doi.org/10.1016/j.cpc.2005.12.009).
- [26] D. Y. Bardin, M. S. Bilenky, T. Riemann, M. Sachwitz, H. Vogt, P. C. Christova, DIZET: A program package for the calculation of electroweak one loop corrections for the process $e^+e^- \rightarrow f^+f^-$ around the Z^0 peak, Comput. Phys. Commun. 59 (1990) 303–312.
doi:[10.1016/0010-4655\(90\)90179-5](https://doi.org/10.1016/0010-4655(90)90179-5).
- [27] A. Leike, S. Riemann, T. Riemann, Z Z-prime mixing and radiative corrections at LEP-1, Phys. Lett. B291 (1992) 187–194.
arXiv:[hep-ph/9507436](https://arxiv.org/abs/hep-ph/9507436), doi:[10.1016/0370-2693\(92\)90142-0](https://doi.org/10.1016/0370-2693(92)90142-0).
- [28] ALEPH collab., DELPHI collab., L3 collab., OPAL collab., SLD Collaboration, LEP Electroweak Working Group, SLD Electroweak Group, SLD Heavy Flavour Group, S. Schael, et al., Precision electroweak measurements on the Z resonance, Phys. Rept. 427 (2006) 257–454.
arXiv:[hep-ex/0509008](https://arxiv.org/abs/hep-ex/0509008), doi:[10.1016/j.physrep.2005.12.006](https://doi.org/10.1016/j.physrep.2005.12.006).
- [29] ALEPH collab., DELPHI collab., L3 collab., OPAL collab., LEP Electroweak Working Group, S. Schael, et al., Electroweak Measurements in Electron-Positron Collisions at W-Boson-Pair Energies at LEP, Phys. Rept. 532 (2013) 119–244.
arXiv:[1302.3415](https://arxiv.org/abs/1302.3415), doi:[10.1016/j.physrep.2013.07.004](https://doi.org/10.1016/j.physrep.2013.07.004).
- [30] K. Sachs, Standard model at LEP II, Proc. of XXXVIII Rencontres de Moriond: Electroweak Interactions and Unified Theories, Les Arcs, March 15-22, 2003.
arXiv:[hep-ex/0307009](https://arxiv.org/abs/hep-ex/0307009).

References IV

- [31] P. Holt, Fermion pair production above the Z^0 resonance, PoS HEP2001 (2001) 115.
http://pos.sissa.it/archive/conferences/007/115/hep2001_115.pdf.
- [32] A. Akhundov, A. Arbuzov, S. Riemann, T. Riemann, The ZFITTER project, Phys. Part. Nucl. 45 (3) (2014) 529–549.
[arXiv:1302.1395](https://arxiv.org/abs/1302.1395), [doi:10.1134/S1063779614030022](https://doi.org/10.1134/S1063779614030022).
- [33] F. Carminati, D. Perret-Gallix, T. Riemann, Summary of the ACAT 2013 round table discussion: Open-source, knowledge sharing and scientific collaboration, J. Phys. Conf. Ser. 523 (2014) 012066.
[doi:10.1088/1742-6596/523/1/012066](https://doi.org/10.1088/1742-6596/523/1/012066).
- [34] M. Grünewald, S. Kirsch, A Possible modification of a LEP energy scan for an improved determination of Z boson parameters, submitted to: Z. Phys. C, KEK scan http://www-lib.kek.jp/cgi-bin/img_index?9402241.