

hep-ph/0507244, 0603030, 0606169, **0611204**

# The effect of small $x$ resummations in global parton fits

Chris White, Nikhef


Small  $x$  meeting - Tuesday 8th July

### Overview

What effect do small  $x$  resummations have on a global parton fit?

- ▶ Solution of BFKL solution at NLL order with running coupling.
- ▶ Implementation of impact factors for light and heavy flavours.
- ▶ Global fit with resummed splitting and coefficient functions.

# The High Energy Problem



- ▶ Coefficient and splitting functions for the proton structure functions unstable at low  $x$  due to terms
$$\sim x^{-1} \bar{\alpha}_S^n \log^m(1/x), \\ m \leq n - 1.$$

- ▶ Divergence due to t-channel gluon exchange at LL order, with some quark mixing at NLL order.
- ▶ Must resum the gluon 4-point function by solving the **BFKL** equation.
- ▶ Relate gluon to structure functions using the  $k_T$  factorisation formula ([Collins & Ellis](#); [Catani, Ciafaloni & Hautmann](#)).

## Running coupling solution of BFKL equation

- Mellin moments:

$$f(\gamma, N) = \int_0^\infty (k^2)^{-\gamma-1} \int_0^1 dx x^N f(x, k^2)$$

- Substitute LO running coupling into BFKL equation (Collins & Kwiecinski):

$$\begin{aligned} \frac{d^2 f(\gamma, N)}{d\gamma^2} &= \frac{d^2 f_I(\gamma, Q_0^2)}{d\gamma^2} - \frac{1}{\bar{\beta}_0 N} \frac{d(\chi_0(\gamma) f(\gamma, N))}{d\gamma} \\ &\quad + \frac{\pi}{3\bar{\beta}_0^2 N} \chi_1(\gamma) f(\gamma, N), \end{aligned}$$

with  $\bar{\beta}_0 = 3/(\pi\beta_0)$ .

- Solve with ansatz:

$$f(N, \gamma) = \exp\left(-\frac{\chi_1(\gamma)}{\bar{\beta}_0 N}\right) \int_\gamma^\infty A(\tilde{\gamma}) \exp\left(\frac{\chi_1(\tilde{\gamma})}{\bar{\beta}_0 N}\right) d\tilde{\gamma}$$

(Ciafaloni & Colferai).

- ▶ Can shift lower limit  $\gamma \rightarrow 0$  up to power-suppressed corrections (Thorne).
- ▶ Gluon factorises:

$$\mathcal{G}(N, t) = \mathcal{G}_E(N, t)\mathcal{G}_I(Q_0^2, N)$$

$$(t = \log Q^2/\Lambda^2).$$

- ▶ Perturbative piece:

$$\mathcal{G}_E^1(N, t) = \frac{1}{2\pi i} \int_{1/2-i\infty}^{1/2+i\infty} \frac{f^{\beta_0}}{\gamma} \exp [\gamma t - X_1(\gamma, N)/(\bar{\beta}_0 N)] d\gamma$$

with:

$$X_1(\gamma, N) = \int_{\frac{1}{2}}^{\gamma} \left[ \chi_0(\tilde{\gamma}) + N \frac{\chi_1(\tilde{\gamma})}{\chi_0(\tilde{\gamma})} \right] d\tilde{\gamma}.$$


- ▶ Similarly, get structure functions:

$$\mathcal{F}_E^1(N, t) = \frac{1}{2\pi i} \int_{1/2-i\infty}^{1/2+i\infty} \frac{h(\gamma, N)f^{\beta_0}}{\gamma} \exp [\gamma t - X_1(\gamma, N)/(\bar{\beta}_0 N)] d\gamma$$

- ▶ If impact factors known, can disentangle all resummed coefficient and splitting functions (within a particular factorisation scheme).
- ▶ However, NLL impact factors  $h(\gamma, N)$  not known. Work in progress ([Bartels, Colferai, Gieseke & Kyrieleis](#)).
- ▶ Instead LL factors with exact gluon kinematics have been calculated ([Bialas, Navelet & Peschanski; White, Peschanski & Thorne](#)).
- ▶ These provide a very good estimate to the full NLL impact factors ([White & Thorne](#)).
- ▶ Can use these to calculate all the NLL resummed coefficient and splitting functions in the DIS scheme.
- ▶ Finally, combine resummed results with NLO DGLAP:


$$P^{tot.} = P^{NLL} + P^{NLO} - \left[ P^{NLL(0)} + P^{NLL(1)} \right]$$

## Results for Splitting Functions



- ▶ Results shown at  $n_f = 4$ ,  $t = 6$ .
- ▶ Running coupling suppresses low  $x$  divergence.
- ▶ NLL kernel and impact factor effects lead to even more suppression.
- ▶ Main feature is a dip below the NLO DGLAP result.

## Comparison with Alternative Approaches



- ▶ Can compare  $P_{gg}$  with ABF and CCSS groups (not latest results)...
- ▶ Dip common to all approaches at NLL.

## Implementation

- ▶ Have calculated all splitting functions (using approx. impact factors).
- ▶ Also have resummed heavy flavour coefficient functions and matrix elements.
- ▶ These are needed for a consistent (general mass variable flavour) scheme at NLL order.
- ▶ Have defined such a scheme ( $\text{DIS}(\chi)$ ) up to NLL+NLO order.
- ▶ Heavy flavours are important!
- ▶ Have all necessary ingredients for a global parton fit...



## A Global Parton Fit

- ▶ Have undertaken a global parton fit at NLL + NLO order in the DIS scheme.
- ▶ Resummed fit gives an overall fit quality  $\chi^2 = 2249$  for 2181 data points.
- ▶ Compare NLO DIS scheme  $\chi^2 = 2352$  and  $\overline{\text{MS}}$  scheme  $\chi^2 = 2307$ .
- ▶ A previous LL resummed fit gave  $\chi^2 = 2336$ , with significant momentum conservation violation.
- ▶ Resummation seems to decrease tension between data sets.



## Effects of Resummation in a Global Fit



- ▶ Recall the NLL splitting functions dip below their NLO counterparts.
- ▶ Allows increased gluon at both large and small  $x$ .
- ▶ Thus can improve fit to Tevatron jet data without compromising fit the HERA data.
- ▶ Gluon is qualitatively different - leads to more stable  $F_L$ .
- ▶ Correct turnover in reduced cross-section at high  $y$ .

## Gluon Distribution




- ▶ Gluons differ for  $x \lesssim 10^{-2}$ .
- ▶ NLL resummed gluon positive and growing at small  $x$ !
- ▶ Not true at fixed order.
- ▶ Positive gluon avoids negative structure functions.

Results -  $F_2$ 


Results -  $F_L$ 

- ▶ Data currently unable to distinguish between predictions.
- ▶ But fixed order looks unstable.

Charm contribution to  $F_2$ 

- ▶ Clearly important at small  $x$ .
- ▶ Thus, consistent treatment of heavy flavours necessary.

## Reduced Cross-Section




- ▶ Turnover required by data at low  $x$  (high  $y$ ) - NLO fails.
- ▶ Resummation helps!  
Interesting to compare with NNLO.


### Conclusions

- ▶ Have undertaken a global parton fit including NLL BFKL resummations.
- ▶ Qualitative changes in gluon behaviour, and some quantitative improvement in the fit.
- ▶ Have to go to NLL in the resummation to achieve sensible results.
- ▶ Dip in evolution (also ABF, CCSS) important.
- ▶ So is correct treatment of heavy flavours.

# Longitudinal Structure Function



- ▶ Clearly see perturbative instability in fixed order results.
- ▶ This is cured by the resummation.

Results -  $F_2^c$ 

- ▶ Resummed fit performs better for small  $x$  data - note slope as  $Q^2$  increases.
- ▶ Fit is also improved over the whole range of  $x$ .
- ▶ Resummed  $F_2^c$  at lower end of range allowed by data.