

Low and High Energy Modeling in Geant4

Hadronic Shower Simulation Workshop
FNAL, 6-8 September 2006
Dennis Wright
(on behalf of Geant4 Collaboration)

Overview

- Quark-Gluon String Model
- Bertini Cascade
- Binary Cascade
- CHIPS

Origin of the QGS (Quark-Gluon String) Model


- Author: H-P. Wellisch, M. Komagorov
- Most code unique to Geant4
 - guidance from Dubna QGS model (N.S. Amelin)
 - fragmentation code based on pre-existing FORTRAN

Applicability


- Incident p, n, π, K
- Also for high energy γ when CHIPS model is connected
- $\sim 20 \text{ GeV} < E < 50 \text{ TeV}$
- Model handles:
 - Selection of collision partners
 - Splitting of nucleons into quarks and diquarks
 - Formation and excitation of quark-gluon string
 - String hadronization
- Damaged nucleus remains. Another Geant4 model must be added for nuclear fragmentation and de-excitation
 - pre-compound model, or CHIPS for nuclear fragmentation

Quark Gluon String Model

- Two or more strings may be stretched between partons within hadrons
 - Strings from cut cylindrical Pomerons
- Parton interaction leads to color coupling of valence quarks
 - sea quarks included too
- Partons connected by quark gluon strings, which hadronize



Quark Gluon String Model Algorithm



- Build up 3-dimensional model of nucleus
- Large γ -factor collapses nucleus to 2 dimensions
- Calculate impact parameter with all nucleons
- Calculate hadron-nucleon collision probabilities
 - based on quasi-eikonal model, using Gaussian density distributions for hadrons and nucleons
- Sample number of strings exchanged in each collision
- Unitarity cut, string formation and decay

The Nuclear Model

- Nucleon momenta are sampled assuming Fermi gas model
- Nuclear density
 - harmonic oscillator shape for $A < 17$
 - Woods-Saxon for others
- Sampling is done in a correlated manner:
 - local phase-space densities are constrained by Pauli principle
 - sum of all nucleon momenta must equal zero

Collision Criterion

- In the Regge-Gribov approach, the probability of an inelastic collision with nucleon i can be written as

$$p_i(b_i, s) = (1/c)(1 - \exp[-2u(b_i, s)]) = \sum_{n=1}^{\infty} p_{i(n)}(b_i, s)$$

- where

$$p_{i(n)}(b_i, s) = (1/c) \exp[-2u(b_i, s)] \frac{[2u(b_i, s)]^n}{n!}$$

is the probability of finding n cut pomerons in the collision

$$u(b_i, s) = \frac{z(s)}{2} \exp(b_i^2/4L(s))$$

is the eikonal amplitude for hadron-nucleon elastic scattering with pomeron exchange

Pomeron Parameters

- The functions $z(s)$ and $L(s)$ contain the pomeron parameters:
 - fitted to N-N, π -N, K-N collision data (elastic, total, single diffraction cross sections)
 - pomeron trajectory: $\alpha_P' = 0.25 \text{ GeV}^{-2}$, $\alpha_P(0) = 1.0808$ for π , K, 0.9808 for N
- Other parameters:
 - energy scale $s_0 = 3.0 \text{ GeV}^2$ for N, 1.5 GeV^2 for π , 2.3 GeV^2 for K
 - Pomeron-hadron vertex parameters also included:
 - coupling: $\gamma_P^N = 6.56 \text{ GeV}^{-2}$
 - radius of interaction: $R_P^{2N} = 3.56 \text{ GeV}^{-2}$

Diffraction Dissociation

- Need to sample the probability of diffraction
 - get it from difference of total and inelastic collision probabilities

$$p_{ij}^{diff}(b_{ij}, s) = \frac{c-1}{c} (p_{ij}^{tot}(b_{ij}, s) - p_{ij}(b_{ij}, s))$$

- where c is the “shower enhancement” coefficient
 - $c = 1.4$ for nucleons, 1.8 for pions
- Splitting off diffraction probabilities with parameter c follows method of Baker 1976

String Formation


- Cutting the pomeron yields two strings
- String formation is done by parton exchange (Capella 94, Kaidalov 82)
 - for each participating hadron, parton densities are sampled
 - requires quark structure function of hadron
 - parton pairs combined into color singlets
 - sea quarks included with $u:d:s = 1: 1: 0.27$

Longitudinal String Fragmentation


- String extends between constituents
- Break string by inserting q-qbar pair according to
 - $u : d : s : qq = 1 : 1 : 0.27 : 0.1$
- At break -> new string + hadron
- Gaussian P_t , $\langle P_t^2 \rangle = 0.5 \text{ GeV}$
- Created hadron gets longitudinal momentum from sampling QGSM fragmentation functions
 - Lund functions also available

QGSM - Results

$\pi^- \text{Mg} \rightarrow \pi^+ X$, $P_{\text{lab}} 320 \text{ GeV}/c$







$$\text{Rapidity } \eta = \frac{1}{2} \ln \left(\frac{E + p_z}{E - p_z} \right)$$



$$P_t^2 [\text{GeV}^2]$$

Cascade Modeling Concept



Geant4 Bertini Cascade: Origin

- A re-engineered version of the INUCL code of N. Stepanov (ITEP)
- Employs many of the standard INC methods developed by Bertini (1968)
 - using free particle-particle collisions within cascade
 - step-like nuclear density
- Similar methods used in many different intra-nuclear transport codes

Applicability of the Bertini Cascade

- inelastic scattering of p , n , π , K , Λ , Σ , Ξ
- incident energies: $0 < E < 10 \text{ GeV}$
 - upper limit determined by lack of partial final state cross sections and the end of the cascade validity region
 - lower limit due to inclusion of internal nuclear de-excitation models
- in principle, can be extended to:
 - anti-baryons
 - ion-ion collisions

Bertini Cascade Model

- The Bertini model is a classical cascade:
 - it is a solution to the Boltzmann equation on average
 - no scattering matrix calculated
- Core code:
 - elementary particle collider: uses free cross sections to generate secondaries
 - cascade in nuclear medium
 - pre-equilibrium and equilibrium decay of residual nucleus
 - nucleus modelled as three concentric spheres of different densities; density constant within sphere

Bertini Cascade Modeling Sequence (1)

- Nuclear entry point sampled over projected area of nucleus
- Incident particle is transported in nuclear medium
 - mean free path from total particle-particle cross sections
 - nucleus modeled as 3 concentric, constant-density shells
 - nucleons have Fermi gas momentum distribution
 - Pauli exclusion invoked
- Projectile interacts with a single nucleon
 - hadron-nucleon interactions based on free cross sections and angular distributions
 - pions can be absorbed on quasi-deuteron


Bertini Cascade Modeling Sequence (2)

- Each secondary from initial interaction is propagated in nuclear potential until it interacts or leaves nucleus
 - can have reflection from density shell boundaries
 - currently no Coulomb barrier
- As cascade collisions occur, exciton states are built up, leading to equilibrated nucleus
 - selection rules for p-h state formation: $\Delta p = 0, \pm 1,$
 $\Delta h = 0, \pm 1, \Delta n = 0, \pm 2$
- Model uses its own exciton routine based on that of Griffin
 - Kalbach matrix elements used
 - level densities parametrized vs. Z and A

Bertini Cascade Modeling Sequence (3)

- Cascade ends and exciton model takes over when secondary KE drops below 20% of its original value or 7 X nuclear binding energy
- Nuclear evaporation follows for most nuclei
 - emission continues as long as excitation is large enough to remove a neutron or α
 - γ emission below 0.1 MeV
- For light, highly excited nuclei, Fermi breakup
- Fission also possible

Validation of the Bertini Cascade



Origin and Applicability of the Binary Cascade

- H.P. Wellisch and G. Folger (CERN)
- Based in part on Amelin's kinetic model
- Incident p, n
 - $0 < E < \sim 3 \text{ GeV}$
- light ions
 - $0 < E < \sim 3 \text{ GeV/A}$
- π
 - $0 < E < \sim 1.5 \text{ GeV}$

Binary Cascade

- Hybrid between classical cascade and full QMD model
- Detailed model of nucleus
 - nucleons placed in space according to nuclear density
 - nucleon momentum according to Fermi gas model
- Nucleon momentum taken into account when evaluating cross sections, collision probability
- Collective effect of nucleus on participant nucleons described by optical potential
 - numerically integrate equation of motion

Binary Cascade Modeling (1)

- Nucleon-nucleon scattering (t-channel) resonance excitation cross-sections are derived from p-p scattering using isospin invariance, and the corresponding Clebsch-Gordan coefficients
 - elastic N-N scattering included
- Meson-nucleon inelastic (except true absorption) scattering modelled as s-channel resonance excitation. Breit-Wigner form used for cross section.
- Resonances may interact or decay
 - nominal PDG branching ratios used for resonance decay
 - masses sampled from Breit-Wigner form

Binary Cascade Modeling (2)

- Calculate imaginary part of the R-matrix using free 2-body cross-sections from experimental data and parametrizations
- For resonance re-scattering, the solution of an in-medium BUU equation is used.
 - The Binary Cascade at present takes the following strong resonances into account:
 - The delta resonances with masses 1232, 1600, 1620, 1700, 1900, 1905, 1910, 1920, 1930, and 1950 MeV
 - Excited nucleons with masses 1440, 1520, 1535, 1650, 1675, 1680, 1700, 1710, 1720, 1900, 1990, 2090, 2190, 2220, and 2250 MeV

Binary Cascade Modeling (3)


- Nucleon-nucleon elastic scattering angular distributions taken from Arndt phase shift analysis of experimental data
- Pauli blocking implemented in its classical form
 - finals state nucleons occupy only states above Fermi momentum
- True pion absorption is modeled as s-wave absorption on quasi-deuteron
- Coulomb barrier taken into account for charged hadrons

Binary Cascade Modeling (4)

- Cascade stops when mean energy of all scattered particles is below A-dependent cut
 - varies from 18 to 9 MeV
 - if primary below 45 MeV, no cascade, just precompound
- When cascade stops, the properties of the residual exciton system and nucleus are evaluated, and passed to a pre-equilibrium decay code for nuclear de-excitation

Binary Cascade - results

p Pb -> n X



Chiral Invariant Phase Space (CHIPS)

- Origin: M.V. Kosov (CERN, ITEP)

- Manual for the CHIPS event generator, KEK internal report 2000-17, Feb. 2001 H/R.

- Use:

- capture of negatively charged hadrons at rest
 - anti-baryon nuclear interactions
 - gamma- and lepto-nuclear reactions
 - back end (nuclear fragmentation part) of QGSC model


CHIPS Fundamental Concepts

- **Quasmon**: an ensemble of massless partons uniformly distributed in invariant phase space
 - a 3D bubble of quark-parton plasma
 - can be any excited hadron system or ground state hadron
- **Critical temperature T_c** : model parameter which relates the quasmon mass to the number of its partons:
 - $M_Q^2 = 4n(n-1)T_c^2 \Rightarrow M_Q \sim 2nT_c$
 - $T_c = 180 - 200 \text{ MeV}$
- **Quark fusion hadronization**: two quark-partons may combine to form an on-mass-shell hadron
- **Quark exchange hadronization**: quarks from quasmon and neighbouring nucleon may trade places

CHIPS Applications

- u,d,s quarks treated symmetrically (all massless)
 - model can produce kaons, but s suppression parameter is needed, η suppression parameter also required
 - real s-quark mass is taken into account by using masses of strange hadrons
- CHIPS is a universal method for fragmentation of excited nuclei (containing quasmons).
- Unique, initial interactions were developed for:
 - interactions at rest such as π^- capture, $p\bar{p}$ annihilation
 - gamma- and lepto-nuclear reactions
 - hadron-nuclear interaction in-flight are in progress
- Anti-proton annihilation on p and π^- capture at rest in a nucleus illustrate two CHIPS modelling sequences

Modeling Sequence for Proton – antiproton Annihilation (1)



Modeling Sequence for Proton - antiproton Annihilation (2)

- anti-proton and proton form a quasmon in vacuum
 - no quark exchange with neighboring nucleons
 - $n = M/2T_c$ quark-partons uniformly distributed over phase space with spectrum $dW/kdk \propto (1 - 2k/M)^{n-3}$
- quark fusion occurs
 - calculate probability of two quark-partons in the quasmon to combine to produce effective mass of outgoing hadron:
 - sample k in 3 dimensions
 - second quark momentum q from spectrum of $n-1$ quarks
 - integrate over vector q with mass shell constraint for outgoing hadron


Modeling Sequence for Proton - antiproton Annihilation (3)

- determine type of final state hadron to be produced
 - probability that hadron of given spin and quark content is produced: $P = (2s_h + 1) z^{N-3} C_Q$
 - C_Q is the number of ways a hadron h can be made from the choice of quarks in the quasmon
 - z^{N-3} is a kinematic factor from the previous momentum selection
- first hadron is produced, escapes quasmon
- randomly sample residual quasmon mass, based on original mass M and emitted hadron mass


Modeling Sequence for Proton - antiproton Annihilation (4)

- Repeat quark fusion with reduced quasmon mass and quark-parton content
- hadronization process ends when minimum quasmon mass m_{\min} is reached
 - m_{\min} is determined by quasmon quark content at final step
 - depending on quark content, final quasmon decays to two hadrons or a hadron and a resonance
 - kaon multiplicity regulated by the s-suppression parameter ($s/u = 0.1$)
 - η/η' suppression regulated by η -suppression parameter (0.3)

Validation of CHIPS for Proton Anti-Proton Annihilation



Modeling Sequence for π^- Capture at Rest in a Nucleus (1)



Modeling Sequence for π^- Capture at Rest in a Nucleus (2)

- pion captures on a subset or **cluster** of nucleons
 - resulting quasmon has a large mass, many partons
 - capture probability is proportional to number of clusters in nucleus
 - 3 clusterization parameters determine number of clusters
- both quark exchange and quark fusion occurs
 - only quarks and diquarks can fuse
 - mesons cannot be produced, so quark-anti-quark cannot fuse as in vacuum case (p-pbar)
 - because q-qbar fusion is suppressed, quarks in quasmon exchange with neighboring nucleon or cluster
 - produces correlation of final state hadrons

Modeling Sequence for π^- Capture at Rest in a Nucleus (3)

- some final state hadrons escape nucleus, others are stopped by Coulomb barrier or by over-barrier reflection
- as in vacuum, hadronization continues until quasimon mass reaches lower limit m_{\min}
 - in nuclear matter, at this point nuclear evaporation begins
 - if residual nucleus is far from stability, a fast emission of p, n, α is made to avoid short-lived isotopes

Known Problems and Improvements (1)

■ QGS:

- gaussian sampling of p_T too simple => incorrect diffraction, not enough π^- suppression in p scattering
- internal cross sections being improved

■ Medium energy (~10 GeV - 60 GeV):

- too low for QGS, HEP models
- too high for cascade, LEP models
- improved parametrized model being developed

■ Cascades:

- no Coulomb barrier in Bertini

Known Problems and Improvements (2)

■ CHIPS:

- originally designed only as final state generator, not intended for projectile interaction with nucleus
- extension planned for inelastic scattering
- neutrino scattering recently added

Backup Slides



String Formation



- Cutting the pomeron yields two strings
- String formation is done by parton exchange (Capella 94, Kaidalov 82)
 - for each participating hadron h , parton densities are sampled

$$f^h(x_1, x_2, \dots, x_{2n-1}, x_{2n}) = f_0 \prod_{i=1}^{2n} u_{p_i}^h(x_i) \delta(1 - \sum_{i=1}^{2n} x_i)$$

- parton pairs combined to form color singlets
- u is quark structure function of hadron h
- sea quarks included with $u:d:s = 1:1:0.27$



QGS Model

Pion and proton scattering



QGS Model



K⁺ Scattering from Au (π^+ inclusive)



Solid dots: J.J.Whitmore et.al., Z.Phys.C62(1994)199

QGS Model

K⁺ Scattering from Au (π^- inclusive)



Solid dots: J.J.Whitmore et.al., Z.Phys.C62(1994)199

Chiral Invariant Phase Space (CHIPS)

- Hadron spectra reflect spectra of quark-partons within quasmon

- 1-D quark exchange:

$$k + M = q + E, \quad k = p - q \quad \Rightarrow \quad k = (E - M + p)/2$$

- 1-D quark fusion:

$$k + q = E, \quad k - q = p \quad \Rightarrow \quad k = (E + p)/2$$


Currently Implemented Mechanisms (1)

- Negative meson captured by nucleon or nucleon cluster:
 - $dE_{\pi} = m_{\pi}$, $dE_K = m_K + m_N - m_{\Lambda}$
- Negative hyperon captured by nucleon or nucleon cluster:
 - $dE_{\Sigma^-} = m_{\Sigma^-} - m_{\Lambda}$, $dE_{\Xi^-} = m_{\Xi^-} + m_N - 2m_{\Lambda}$, $dE_{\Omega} = m_{\Omega} + 2m_N - 3m_{\Lambda}$
- Nuclear capture of anti-baryon:
 - annihilation happens on nuclear periphery
 - 4π explosion of mesons irradiates residual nucleus
 - secondary mesons interacting with residual nucleus create more quasimons in nuclear matter
 - large excitation: $dE = m_{\text{antibaryon}} + m_N$


Currently Implemented Mechanisms (2)

- In photo-nuclear reactions γ is absorbed by a quark-parton
 - $dE_\gamma = E_\gamma$
- In back-end of string-hadronization (QGSC model) soft part of string is absorbed:
 - $dE_{\text{QGSC}} = 1 \text{ GeV/fm}$
- lepto-nuclear reactions γ^* , W^* are absorbed by quark-parton:
 - $dE_l = E_{\gamma^*}$, $\cos(\theta_k) = (2k/v - Q^2)/2kq$, $Q^2 = q^2 - v^2$
 - with $k < M/2$, if $q - v < m_N$, virtual γ cannot be captured by one nucleon


P-pbar Annihilation into Two Body Final States



Validation of CHIPS Model for Pion Capture at Rest on Tantalum



Neutrons from C on C at 290 MeV/c



$p+Ta=h+X$, 400 GeV, 90°

