String-Math 2016, Collège de France, Paris



Contribution ID: 34

Type: not specified

## Umbral symmetry groups and K3 CFTs

Friday 1 July 2016 15:30 (25 minutes)

Umbral moonshine is a connection between mock modular forms and discrete symmetry groups which arise as automorphisms of the Niemeier lattices, the 24-dimensional unimodular lattices labeled by their ADE root systems. The first example of Umbral moonshine was original discovered by Eguchi, Ooguri, Tachikawa when expanding the elliptic genus of a K3 surface into N=4 characters and seeing coefficients which are sums of irreducible representations of the sporadic group M\_{24}. In recent work with Miranda Cheng, we have suggested that all umbral symmetry groups may have some relation to K3 CFTs. On the other hand, a theorem of Gaberdiel, Hohenegger, and Volpato has shown that symmetries of K3 CFTs can naturally be classified by elements of the group Conway\_0, the automorphism group of the Leech lattice, which preserve a four-plane. In this talk we briefly review the aforementioned results, and discuss the extension of this theorem a) to singular CFTs, in which case the role of Conway\_0 is naturally replaced by certain umbral groups, and b) when one takes worldsheet parity into account. Based on [ArXiv:1406.0619, 1512.04942], and work to appear with Miranda Cheng and Roberto Volpato

## **Summary**

**Presenter:** HARRISON, Sarah (Harvard) **Session Classification:** Plenary session