



Contribution ID: 15

Type: **not specified**

## Towards a Maximum Likelihood estimate of the dwarf spheroidal galaxies $J$ factor

Wednesday 23 September 2015 08:55 (25 minutes)

The dwarf spheroidal satellite galaxies of the Milky Way appear to be the most dark matter (DM) dominated objects in the near Universe. Their very low astrophysical background in the  $\gamma$ -ray energy range makes them ideal targets for DM indirect detection, which can be achieved by searching for its decay or annihilation signals. The latter approach requires the calculation of the  $J$ -factor, which quantifies the amount of DM along the line of observation. This quantity has been previously derived with bayesian techniques, thereby subjecting the results to the effects of priors. We report here the development of a new fully frequentist approach based on the Maximum Likelihood procedure, which thus allows to build the profile likelihood for  $J$  and from it the calculation of confidence intervals. The new results are in most cases consistent with the previously derived ones, being compatible to the  $1\text{-}\sigma$  level; their uncertainties are also compatible and consistently scale with the dataset size, being smaller for the larger samples; the largest discrepancies are restricted to the systems with the smallest dataset. We also present possible improvements and extensions to this technique.

**Authors:** Mr CHIAPPO, Andrea (The Oskar Klein Center, Stockholm University); Prof. CONRAD, Jan (Stockholm University); COHEN-TANUGI, Johann (Université de Montpellier, CNRS/IN2P3); STRIGARI, Louis (Stanford University)

**Presenter:** Mr CHIAPPO, Andrea (The Oskar Klein Center, Stockholm University)

**Session Classification:** Dark Matter density