

Recasting LHC analyses with MADANALYSIS 5

Fuks Benjamin

IPHC - U. Strasbourg

With E. Conte & the PAD team
(S. Kraml *et al.*, K. Mawatari & K. de Causmaecker, *etc.*)

Joint LPC & MC4BSM *Data Challenge* @ Fermilab, USA

21 May 2015

Outline

1. MADANALYSIS 5 in a nutshell
2. Reinterpretation of LHC analyses with MADANALYSIS 5 and DELPHES
3. The PAD (public analysis database) and examples
4. Summary

MADANALYSIS 5 in a nutshell

[Conte, BF, Serret (CPC '13); Conte, Dumont, BF, Wymant (EPJC '14)]

◆ What is MADANALYSIS 5?


- ❖ A framework for **phenomenological analyses**
 - ★ Prospective studies
 - ★ **Reinterpretation studies**
- ❖ **Any level of sophistication**: partonic, hadronic, detector, reconstructed
- ❖ **Several input** format: STDHEP, HEPMC, LHE, LHCO, ROOT (from DELPHES)
- ❖ User-friendly, flexible and fast
- ❖ **Interfaces** to other HEP packages (fast detector simulation, jet clustering, etc.)

◆ Two modules

- ❖ A **PYTHON** command line interface (interactive)
- ❖ A **C++/ROOT** core module, SAMPLEANALYZER

MADANALYSIS 5: normal mode

[Conte, BF, Serret (CPC '13); Conte, Dumont, BF, Wymant (EPJC '14)]



MADANALYSIS 5: expert mode

[Conte, BF, Serret (CPC '13); Conte, Dumont, BF, Wymant (EPJC '14)]

User inputs

Event samples

Job

Execution

SAF files

Code skeleton to be generated from the PYTHON interface

Libraries

SAMPLEANALYZER
FASTJET *
DELPHES 3 **

To be installed from the PYTHON interface

* [Cacciari, Salam (PLB '06)]
** [de Favareau et al (JHEP'14)]

Reimplementing new physics LHC searches (I)

- ◆ Many search channels are investigated by CMS and ATLAS
 - ❖ Limits set on popular new physics theories and/or simplified models
 - ❖ There are plethora of new physics realizations that deserve to be studied
 - ★ The simplified model approach is **not sufficient** (e.g., different topologies)
 - ★ Need for implementations of LHC analyses in **public tools**
 - ★ Need to rely on a **public detector simulation**

- ◆ The MADANALYSIS 5 way
 - ❖ Use of the expert mode of the program as a framework
 - ❖ Use of DELPHES 3 for detector simulation

Reimplementing new physics LHC searches (2)

◆ Validation of the reimplementations

- ❖ **Built-in differences:** DELPHES versus ATLAS and CMS detector simulations
- ❖ Comparison of cut-flows, kinematical distributions for specific benchmarks
- ❖ Aiming for a **20%-30% agreement**

◆ Complications: incomplete experimental documentation

- ❖ **But... the material is better and better!**
- ❖ Loss of months in exchanges with ATLAS and CMS
- ❖ Sometimes dead ends

[See the forthcoming discussion]

Detector simulation with DELPHES and MADANALYSIS (I)

[Dumont, BF, Kraml et al. (EPJC '15)]

◆ Fast detector simulation with MADANALYSIS 5

- ♣ MADANALYSIS 5 has been interfaced to DELPHES 3
- ♣ Starts from events at the hadron level and produces ROOT files (DELPHES)
- ♣ DELPHES is modular ➤ MADANALYSIS 5 includes extra modules (DELPHES-MA5Tune)
 - ★ Extra information on lepton isolation
 - ★ Track information
 - ★ Smaller output ROOT files (DELPHES)
 - ★ This version of DELPHES can be executed from MADANALYSIS 5

Detector simulation with DELPHES and MADANALYSIS (2)

[Dumont, BF, Kraml et al. (EPJC '15)]

◆ Running DELPHES in MADANALYSIS 5

- ❖ Running of DELPHES via the MADANALYSIS 5 interpreter (in the reco mode)
- ❖ Choice of ATLAS or CMS; pile-up can be included
- ❖ Different cards could be necessary for different analyses
- ❖ The ROOT output file is stored

◆ Future developments

- ❖ In collaboration with the CHECKMATE team (common analysis implementations)
- ❖ Use of the standard DELPHES with a DELPHES-recasting card

The expert mode of MADANALYSIS 5 (I)


[Conte, Dumont, BF, Wymant (EPJC '14)]

◆ Main features (enable an easy recast of cut-based LHC analyses)

- ❖ Support for **multiple sub-analyses** (signal and control regions)
- ❖ New ready-to-use observables (M_{T2} , M_{T2W} , etc.)
- ❖ New optimized handling of cuts and histograms

◆ Handling cuts and histograms

- ❖ Naive approach **not efficient** (see cut #4 for instance)



```
count the event in region D
if (condition 3)
{
  count the event in region C
  if (condition 4)
  {
    count the event in region A
  }
}
if (condition 4)
{
  count the event in region B
}
```

- ❖ A **more efficient** algorithm has been implemented
 - ★ Each cut condition is only evaluated once
 - ★ It is applied to all surviving regions **simultaneously**
- ❖ Similar treatment for histograms

The expert mode of MADANALYSIS 5 (2)

[Conte, Dumont, BF, Wymant (EPJC '14)]

◆ MADANALYSIS 5 is used without its PYTHON interface

- ❖ More freedom in the **observables** (only some of them can be called from the PYTHON console)
- ❖ **Complicated cuts** can be implemented
- ❖ More suitable for **large numbers of events** (using several cores)

◆ The expert mode is developer-friendly

The analysis is a C++ class

The SAMPLEANALYZER internal data format

- ★ Readers for LHE, STDHEP, HEPMC, LHCO and DELPHES
- ★ Many classes and methods for particle and object properties
- ★ Specific methods for histograms and cuts
- ★ etc.

Services

- ★ Physics observables (transverse variables, object identification, isolation)
- ★ Streamers
- ★ Exceptions
- ★ etc.

Interfaces

- ★ FASTJET
- ★ DELPHES 3
- ★ New DELPHES modules

The new DELPHES modules (linked to SAMPLEANALYZER):

- ★ isolation
- ★ tracks
- ★ output files

Scripts

- ★ Compilation
- ★ Linking
- ★ Analysis skeleton generator

The Physics Analysis Database (PAD) of MADANALYSIS

[Dumont, BF, Kraml et al. (EPJC '15)]

- ◆ A database with MADANALYSIS 5 implementations of LHC analyses has been initiated
- ❖ <http://madanalysis.irmp.ucl.ac.be/wiki/PhysicsAnalysisDatabase>
- ❖ Easy to install (*install PAD*)
- ❖ Seven analyses are public and validated

ATLAS analyses, 8 TeV

Analysis	Short Description	Implemented by	Code	Validation note	Status
ATLAS-SUSY-2013-05 (published)	stop/sbottom search: 0 leptons + 2 b-jets	G. Chalons	Inspire	PDF (figures)	done
ATLAS-SUSY-2013-11 (published)	EWK-inos, 2 leptons + MET	B. Dumont	Inspire	PDF (source)	done
ATLAS-HIGG-2013-03 (published)	ZH->ll+invisible	B. Dumont	Inspire	PDF (source)	done
ATLAS-EXOT-2014-06 (published)	mono-photons + MET	D. Barducci	Inspire	PDF	done

[Delphes card](#) for ATLAS-SUSY-2013-05
[Delphes card](#) for ATLAS-SUSY-2013-11

DELPHES cards

Code from INSPIRE

Validation notes
(cutflows, distributions, etc.)

CMS analyses, 8 TeV

Analysis	Short Description	Implemented by	Code	Validation note	Status
CMS-SUS-13-011 (published)	stop search in the single lepton mode	B. Dumont, B. Fuks, C. Wymant	Inspire [1]	PDF (source)	done
CMS-SUS-13-012 (published)	gluino/squark search in jet multiplicity and missing energy	S. Bein, D. Sengupta	Inspire	PDF (source)	done
CMS-SUS-13-016 (PAS)	search for gluinos using OS dileptons and b-jets	D. Sengupta, S. Kulkarni	Inspire	PDF (source)	done

[Delphes card](#) for these analyses

DELPHES cards

MADANALYSIS 5 analyses on INSPIRE

[Dumont, BF, Kraml et al. (EPJC '15)]

- ◆ Implementation of LHC analyses can be uploaded on INSPIRE
- ❖ DOI are assigned: can be cited, searched for, etc.

Information Citations (3) Files Files are versioned, can be downloaded

MadAnalysis 5 implementation of CMS-SUS-13-011: search for stops in the single lepton final state at 8 TeV

DOI and citations
 Dumont, Beranger (LPSC, Grenoble); Fuks, Benjamin (CERN); Wymant, Chris (Annecy, LAPTH)

Cite as: (2014) authors, <http://doi.org/10.7484/INSPIREHEP.DATA.LR5T.2RR3>

Description: This is the MadAnalysis 5 implementation of the CMS search for top-squark pair production in the single lepton final state with 19.5/fb at 8 TeV, to be used for re-interpretation studies. The C++ code contains extensive comments and can thus easily be used as a template for implementing other analyses.

Note: This analysis requires MINUIT libraries. Therefore, the line <LIBFLAGS += -lMinuit> should be added to the Makefile of the Build/ directory before compilation. More information how to use this code as well as a detailed validation summary are available at <http://madanalysis.irmp.ucl.ac.be/wiki/PhysicsAnalysisDatabase>

Cite as: Dumont, B., Fuks, B., Wymant, C. (2014) MadAnalysis 5 implementation of CMS-SUS-13-011: search for stops in the single lepton final state at 8 TeV. doi: [10.7484/INSPIREHEP.DATA.LR5T.2RR3](http://doi.org/10.7484/INSPIREHEP.DATA.LR5T.2RR3)

This dataset complements the following publication:
[Toward a public analysis database for LHC new physics searches using MADANALYSIS 5](#)

Record added 2014-06-19, last modified 2014-07-17


CMS-SUS-13-011

[Dumont, BF, Wymant (2014)]

- ◆ CMS search for stops in the single lepton channel
 - ♣ Benchmark: stop of 650 GeV and neutralino of 50 GeV

◆ Validation with public material from CMS

Cut	MADANALYSIS 5	CMS
At least one lepton, four jets and 100 GeV of missing transverse energy	31.4	29.7
At least one b -tagged jet	27.1	25.2
No extra loosely-isolated lepton or track	22.5	21.0
No hadronic tau	22.0	20.6
Angular separation between the missing momentum and the two hardest jets	18.9	17.8
Hadronic top quark reconstruction	12.7	11.9
The transverse mass M_T (defined in the text) is larger than 120 GeV	10.4	9.6
At least 300 GeV of missing transverse energy and $M_{T2}^W > 200$ GeV	5.1	4.2




CMS results can be reproduced at the 20%-30% level

ATLAS-SUS-2013-05

[Chalons (2014)]

- ◆ ATLAS search for stops and sbottoms in the zero lepton plus 2 b-jets channel
- ♣ Benchmarks: sbottom/neutralino (300/200 GeV) and stop/chargedino/neutralino (250/155/50 GeV)


◆ Validation with public ATLAS material



ATLAS limits can be reproduced
at the 50 GeV level

◆ Setting limits

- ♣ CLs: from observed and expected numbers
- ♣ Best expected signal region is employed
- Conservative



Reinterpretation and constraining new physics

[Belanger, Dumont, Goudelis, Herrmann, Kraml, Sengupta (2015)]

◆ ATLAS searches in dilepton + missing energy channel

- ♣ ATLAS-SUSY-2013-11: chargino, neutralino and slepton searches [10.7484/INSPIREHEP.DATA.HLMR.T56W.2]
- ♣ ATLAS-HIGG-2013-03: Z (to two leptons) + H (invisible) [10.7484/INSPIREHEP.DATA.RT3V.9PJK]

◆ Constraining the inert doublet model

- ♣ One SM Higgs + non-mixing extra scalars
- ♣ The dilepton + missing energy probe

$$q\bar{q} \rightarrow Z \rightarrow A^0 H^0 \rightarrow Z^{(*)} H^0 H^0 \rightarrow \ell^+ \ell^- H^0 H^0,$$


$$q\bar{q} \rightarrow Z \rightarrow H^\pm H^\mp \rightarrow W^{\pm(*)} H^0 W^{\mp(*)} H^0 \\ \rightarrow \nu \ell^+ H^0 \nu \ell^- H^0,$$

$$q\bar{q} \rightarrow Z \rightarrow Z h^{(*)} \rightarrow \ell^+ \ell^- H^0 H^0,$$

$$q\bar{q} \rightarrow Z \rightarrow Z H^0 H^0 \rightarrow \ell^+ \ell^- H^0 H^0.$$

◆ Constraints in the model parameter space

- ♣ Regions with $\mu < 1$ are excluded (95% CL)



Summary

◆ MADANALYSIS 5 in a nutshell

- ❖ A **unique** framework for collider phenomenology at all levels (parton, hadron, reconstructed)
- ❖ **User-friendly** by means of its PYTHON interface (normal mode)
- ❖ **Flexible** thanks to its C++ kernel (expert mode)
- ❖ **Interfaced** to several other HEP packages (DELPHES, FASTJET)

◆ MADANALYSIS 5 and LHC analyses

- ❖ The expert mode has been extended to facilitate the implementation of LHC analyses
- ❖ **Seven analyses** are validated and public
 - ATLAS and CMS results are recovered at the 20-30% level
- ❖ **O(20) analyses are being validated** (i.e., will become public soon)
- ❖ An **interface with CHECKMATE** is being developed

◆ Please use and contribute to the database

- ❖ <http://launchpad.net/madanalysis5>
- ❖ <http://madanalysis.irmp.ucl.ac.be/wiki/PhysicsAnalysisDatabase>