

# Experimental Summary and Outlook

---

H. Jung (DESY, University Antwerp)  
[hannes.jung@desy.de](mailto:hannes.jung@desy.de)

# Experimental Summary and Outlook

---

H. Jung (DESY, University Antwerp)  
[hannes.jung@desy.de](mailto:hannes.jung@desy.de)

How to prepare a good summary talk ?

# Experimental Summary and Outlook

---

H. Jung (DESY, University Antwerp)  
[hannes.jung@desy.de](mailto:hannes.jung@desy.de)

How to prepare a good summary talk ?

Look at previous summary talks

# Experimental Summary and Outlook

---

H. Jung (DESY, University Antwerp)  
[hannes.jung@desy.de](mailto:hannes.jung@desy.de)

How to prepare a good summary talk ?

Look at previous summary talks

- Too many individual talks, so cannot cover everything !

# Experimental Summary and Outlook

---

H. Jung (DESY, University Antwerp)  
[hannes.jung@desy.de](mailto:hannes.jung@desy.de)

How to prepare a good summary talk ?

Look at previous summary talks

- Too many individual talks, so cannot cover everything !
  - Make selection ?
    - what should be left out ?

# Experimental Summary and Outlook

---

H. Jung (DESY, University Antwerp)  
[hannes.jung@desy.de](mailto:hannes.jung@desy.de)

How to prepare a good summary talk ?

Look at previous summary talks

- Too many individual talks, so cannot cover everything !
  - Make selection ?
    - what should be left out ?
    - instead of covering everything, I'll ask what I do not understand

# Soft QCD and Multiparton Dynamics

# Contributions

---


- Soft QCD and multi-parton interactions, FIELD, Richard Dryden
- Soft QCD measurements from the LHC, BRUNI, Alessia
- Overview of MC tuning to LHC data, SIEGERT, Frank
- QCD in Heavy Ion Collisions, MORSCH, Andreas
- Hard Diffraction at HERA, BRITZGER, Daniel
- Results on total pp cross sections, diffractive and exclusive final states from ATLAS, ARRATIA MUNOZ, Miguel Ignacio
- Results on minimum bias interactions, underlying event and particle production from ATLAS, CAIRO, Valentina
- Associated quarkonium production at ATLAS as a new probe of QCD, BERTSCHE, David Edwin
- Exclusive processes at HERA, BRUNI, Alessia
- Small-x Physics at the LHeC, ARMESTO PEREZ, Nestor
- Di-photon and photon-hadron correlations at the LHC, REZAEIAN, Amir
- Data-driven approaches to pile-up treatment at the LHC, VAN HAEVERMAET, Hans
- Searches for resonant and non-resonant new phenomena in CMS, YOO, Hwi Dong
- Searches for resonant and non-resonant new phenomena from ATLAS, ELLIOT, Alison
- Precision calculations for squark and gluino production at threshold, BORSCHENSKY, Christoph
- Hadronization processes in neutrino interactions for oscillation physics, KATORI, Teppei



# Total x-section in pp

J. Butterworth, Overview, plenary  
 A. Bruni, Soft QCD, plenary  
 M. Arratia (ATLAS), Total pp xsections  
 Multiparton Dynamics

## Results: Total cross-section




$$\sigma = 73.1 \pm 0.9(\text{exp.}) \pm 6.6(\text{lumi}) \pm 3.8(\text{extrapol.}) \text{ mb}$$


- First measurement of  $\sigma_{pp}$  at  $\sqrt{s} = 13 \text{ TeV}$
- Important consistency check, needed for all calculations and simulations, also for pile-up !

Measurement with different methods:

- at 13 TeV:  
detecting final states





- at 8 TeV:  
using ALPFA and elastic slope



28 / 29

# Minimum Bias measurements



- Very essential measurements for experiments
  - basic x-check of MC prediction needed for pileup simulation
  - big success of MC tuning: no extra tune needed for Min Bias !

# Apropos pileup ....

- Many QCD processes are distorted by high pileup at high luminosity



arXiv 1508.07811

> Extract signal without relying on Monte Carlos

> From mixed sample can extract true signal successfully

> Advantages:

- works in high  $N_{PU}$  regime
- no data at low pile-up needed
- no Monte Carlo needed




# Parton Density Functions

# Contributions

---

- Status of PDFs, ALEKHIN, Sergey
- Proton structure and hard QCD at HERA, SARKAR, Amanda
- QCD at the LHeC, NEWMAN, Paul Richard
- MMHT14 PDFs: updates and outlook, HARLAND-LANG, Lucian
- The CT14 Global Analysis of QCD, STUMP, Daniel
- On the impact of lepton PDFs, CARRAZZA, Stefano
- Self Organizing Maps Parameterization of Parton Distribution Functions, SIMONETTA, Iuti
- Measurements of  $W$  charge asymmetry, PENNING, Bjoern
- HERAFitter project and its related studies, SAPRONOV, Andrey
- Studies of PDF sensitivity relevant for ATLAS measurements, HANNA, Remie
- Jet measurements,  $\alpha_s$  and PDF results from CMS, RABBERTZ, Klaus
- Recent developments in the fast reproduction of QCD calculations with the APPLgrid project, SUTTON, Mark
- fastNLO v2 Developments, RABBERTZ, Klaus
- Nuclear PDFs in eA collisions at the LHeC, ARMESTO PEREZ, Nestor
- Proton PDFs at the LHeC, GWENLAN, Claire
- Parton Distributions for the LHC Run II, ROTTOLI, Luca
- Reduction strategies for the combination of PDF sets, ROJO CHACON, Juan
- New PDF4LHC recommendations for parton distribution uncertainties for the LHC run-II era, DE ROECK, Albert
- New CTEQ-Jefferson Lab (CJ15) analysis of parton distributions, MELNITCHOUK, Wally
- Impact of heavy-flavour production cross sections measured by the LHCb experiment on parton distribution functions at low  $x$ , SARKAR, Amanda
- Forward charm production and the small- $x$  gluon PDF, GAULD, Rhorry
- A Variable Flavour Number Scheme with Intrinsic Charm, BONVINI, Marco

## 6) Incl. jet, di-jet and 3-jet cross-sections JHEP 02 (2015) 153 - JHEP05(2014)059 4.5 fb<sup>-1</sup> of 7 TeV data - Eur. Phys. J. C (2015) 75



- Inclusive jet cross section** : the predictions CT10, MSTW2008, NNPDF 2.1 (left), HERA1.5, ABM11 (right) are generally consistent with the measured cross-sections, except in a few bins (HERA, ABM at low rapidity).

# Jet measurements: constrain PDFs

K. Rabbertz (CMS), Jets  
Hard QCD & PDFs


Agreement with predictions of QCD over many orders of magnitude in cross section and beyond 2 TeV in jet  $p_T$

Constrains PDFs  
"Harder" gluon at high  $x$  compared to DIS


$$\frac{d^2\sigma}{dp_T dy} \propto \alpha_s^2$$

Data vs. NLO+PS @ EW corrections  
→ impact visible in dijet angular obs.

anti- $k_T$ ,  $R=0.7$ , 7 TeV, 2011




CMS, EPJC 75 (2015) 288.



# Tools for calculations: **herafitter**, applgrid, fastnlo, ...

- HERAFitter is an open-source tool for QCD analyses in lepton-hadron and hadron-hadron

A. Saprosov HERAFitter  
PDF



30 public results obtained using HERAFitter from the beginning of the project

- Perform QCD analysis of proton structure using experimental data
- Extract and improve precision of the PDFs
- Assess impact of the new data on PDF determination
- Check experimental data consistency
- Test different theoretical approaches to the proton structure description.





# Tools for calculations: **herafitter**, applgrid, fastnlo, ...

- HERAFitter is an open-source tool for QCD analyses in lepton-hadron and hadron-hadron

A. Saprosov HERAFitter PDF


The Tevatron W and Z data exhibit significant impact on valence quarks relative to HERA-I data alone. Especially on d-quark.



# Tools for calculations: herafitter, **applgrid**, fastnlo, ...

- Applgrid is a **fully open source package** to build a library of C++ utility classes for performing fast (N)NLO convolutions with PDFs
- includes now:
  - MCFM, NLOjet++ for jet prod.
  - All fixed order NLO processes in aMC@NLO, using the aMCfast interface, arXiv:1406.7693
  - Sherpa for fixed NLO processes using MCgrid, arxiv:1312.4460
  - Sherpa interface is available



- Written by Del Debbio, Hartland and Schumann [arXiv:1312.4460](http://arxiv.org/abs/1312.4460). <http://mcgrid.hepforge.org>
- Available for all fixed order NLO processes using the generic PDF decomposition



- MCgrid works as a Rivet plugin using the HepMC event record


M Sutton - Recent developments with APPLgrid

M. Sutton (ATLAS), applgrid,  
Hard QCD & PDF




# Tools for calculations: herafitter, applgrid, **fastnlo**, ...

- fastNLO is a **fully open source package** to build a library of C++ utility classes for performing fast (N)NLO convolutions with PDFs
- includes now: NLOjet++ for jet production, DIFFTop, BlackHat
- Sherpa for fixed NLO processes using MCgrid2




MCgrid, L. Del Debbi et al., CPC185, 2011

K. Rabbertz (CMS),  
fastNLO  
Hard QCD & PDFs




# Tools for calculations: herafitter, applgrid, **fastnlo**, ...

- fastNLO is a **fully open source package** to build a library of C++ utility classes for performing fast (N)NLO convolutions with PDFs
- includes now: NLOjet++ for jet production, DIFFTop, BlackHat
- Sherpa for fixed NLO processes using MCgrid2



K. Rabbertz (CMS),  
fastNLO  
Hard QCD & PDFs



Move towards a common interface for both applgrid and fastNLO with the different QCD calculations !

# New PDF4LHC recommendations

A. DeRoeck, PDF4LHC, PDF  
J. Rojo, Reduction Strategies, PDF

## Basically two kinds of usage:

- For assessment of PDF uncertainties in comparing predictions to theory (top xsection, jets, W/Z xsections)
  - **Use individual PDF sets (CT, NNPDF, MMHT, HERAPDF, ABM, JR, CJ ...)**
- For assessment of PDF uncertainties as in searches, discoveries, acceptance measurements (e.g. Higgs, SUSY, ... )
  - **Use PDF4LHC prescription**
    - PDF4LHC15\_mc: a Monte Carlo PDF set with  $N_{\text{rep}} = 100$  replicas
    - PDF4LHC15\_30: a symmetric Hessian PDF set with  $N_{\text{eig}} = 30$  eigenvectors
    - PDF4LHC15\_100: a symmetric Hessian PDF set with  $N_{\text{eig}} = 100$  eigenvectors
    - These are available at NLO and NNLO and for standard  $\alpha_s = 0.118$  and  $\delta\alpha_s = \pm 0.0015$  variations
    - Available in LHAPDF repository
  - **Which ones to use when: Part of the recommendation being documented eg**
    - \_MC set: contains also non-gaussian effects. Important for searches at high masses. ..
    - \_30 set: very good precision and probably usable for most experimental needs
    - \_100 superior precision, eg for TH Higgs cross section uncertainty determinations...

# Hard QCD

# Contributions

---

- LHC Experimental Overview and Prospects, BUTTERWORTH, Jonathan
- Theoretical Developments in Hard QCD, Predictions GLOVER, Nigel
- Recent QCD results from the LHC, ARCIDIACONO, Roberta
- Frontiers of QCD: new insights from SCE WAALEWIJN, Wouter
- Recent developments in QCD MCs SOPER, Davison
- Multiboson production at the LHC (theory) VON MANTEUFFEL, Andreas
- Multiboson production at the LHC ROVELLI, Chiara Ilaria
- EW corrections at high energy, DENNER, Ansgar
- Developments in QCD analytic resummation, LAENEN, Eric
- QCD in Higgs and BSM, TOMPKINS, Lauren Alexandra
- Measurements of jet and photon production in pp collisions with the ATLAS detector, DE LA TORRE, Hector
- Measurement of the Inclusive Isolated Prompt Photon production cross section in ppbar collisions at  $\sqrt{s}=1.96\text{TeV}$  using the full CDF dataset, LUCA, Alessandra
- Numerical multi-loop calculations based on the program SecDec, HEINRICH, Gudrun
- Five-point two-loop master integrals in QCD, LO PRESTI, Nicola Adriano
- Color and Kinematic Decomposition for QCD Amplitudes, OCHIROV, Alexander
- Studies of PDF sensitivity relevant for ATLAS measurements, HANNA, Remie
- Jet measurements,  $\alpha_s$  and PDF results from CMS, RABBERTZ, Klaus
- Subleading Logarithms in High Energy Jets, BROOKS, Helen
- Recent developments in the fast reproduction of QCD calculations with the APPLgrid project, SUTTON, Mark
- fastNLO v2 Developments, RABBERTZ, Klaus

# Contributions

---


- Higgs measurements from CMS, ZGHICHE, Amina
- Higgs measurements from ATLAS, LIU, Bo
- NNLO cross sections using the jettiness method, GIELE, Walter
- Higgs + jet at NNLO QCD: Fiducial cross sections, SCHULZE, Markus Christian
- Approximate N3LO: Higgs and more, BONVINI, Marco
- Analytic NNNLO corrections to the total Higgs production cross section in the  $qq'$ -Channel, HASSELHUHN, Alexander
- Fully differential decay rate of a standard model Higgs boson into a b-quark pair at NNLO accuracy, TROCSANYI, Zoltan Laszlo
- Fully differential VBF Higgs production at NNLO, DREYER, Frederic Alexandre
- Higgs production in gluon fusion in association with jets, GREINER, Nicolas
- Electroweak and QCD corrections with MadGraph5\_aMC@NLO: top quark pair production in association with heavy bosons, SHAO, Huasheng
- Vector boson plus jets measurements with the ATLAS detector, TANNENWALD, Benjamin
- Automated NLO QCD+EW corrections for V+multijet production, LINDERT, Jonas
- Z boson production in association with a jet at NNLO QCD, MORGAN, Thomas
- Loop-tree duality and its application to NLO computations, SBORLINI, German
- Measurements of Drell--Yan transverse momentum, lepton azimuthal decorrelation and angular distributions with the ATLAS detector, LIN, Tai-Hua
- W/Z results from CMS, LAWHORN, Jay
- NNLOPS Simulations of Higgs and Vector Boson Production, HAMILTON, Keith
- Combining parton showers with NNLO matrix elements, LI, Ye
- Recent electroweak results from ATLAS, BAAS, Alessandra
- Diboson production at NNLO, GRAZZINI, Massimiliano
- Automated NNLL+NLO Resummation for Jet-Veto Cross Sections, ROTHEN, Lorena



# Jet measurements


J. Butterworth, Overview, plenary  
 R. Arcidiacono, Recent results, plenary  
 H. de la Torre (ATLAS), Measurements of jets  
 Hard QCD

- ▶ Sensitivity to resonances (as a function of dijet mass) decaying into two jets.
- ▶ High dijet-mass region constrains gluon PDF.
- ▶  $P_T^{jet1} \geq 100$  GeV,  $P_T^{jet2} \geq 50$  GeV
- ▶  $|y^{jet}| < 3.0$  and  $|y^*| < 3.0$
- ▶  $y^* = |y^{jet1} - y^{jet2}|/2$
- ▶ NLO pQCD: NLOJET++.
- ▶ Several PDF sets.
- ▶ NLO is corrected for non-perturbative and EW effects.
  - ▶ NPC obtained with PYTHIA/HERWIG with various tunes.
- ▶ Data unfolded to particle-level using an IDS method.




# Dijet measurements: azimuthal correlations

$\Delta\phi_{jj}$  in bins of  $p_{T1}$  for  $p_T > 100$  GeV,  $p_{T1} > 200$  GeV,  $|y_1|, |y_2| < 2.5$   
 - dijet LO configuration is always  $\Delta\phi_{jj} = \pi$   
 - deviations through multijet production



**Comparison to fixed-order PQCD**  
 → need multijet NLO  
 Sherpa + BlackHat → 4-jet NLO  
 Njet → 5-jet NLO to be checked  
 Comparison to LO ME+PS and multijet ME+PS  
 → good general description



CMS-PAS-SMP-14-015 (2015).



# Dijets for searches




H. Yoo, Searches in CMS  
 A. Elliot Searches from ATLAS  
 Multiparton Dynamics

- 13 TeV data are already analyzed




# Diboson production

H. Yoo, Searches in CMS  
 A. Elliot Searches from ATLAS  
 Multiparton Dynamics



# Higgs +jet measurements

L. Tompkins, QCD in Higgs and BSM, plenary  
 A.Zghiche, Higgs from CMS, Hard QCD  
 B.Liu Higgs from ATLAS, Hard QCD  
 K.Hamilton, NNLOPS, Hard QCD



- Jet bins low for all MCs ; HNNLOPS expectedly better in 0-jet bin
- “ ... poorest agreement with data ... in the inclusive and exclusive 1-jet bins, with p-values ranging between 0.1% and 3.6%.”
- Normalised lead jet  $p_T$  agreeing well between data & all MCs

# Heavy Quarks

# Contributions

---

- Heavy flavour theory developments, DE FREITAS, Abilio
- Heavy quark production results, NEEDHAM, Matthew David
- Top quark production at the LHC, CUEVAS MAESTRO, Javier
- Heavy flavour production with the ATLAS experiment, BJERGAARD, David Martin
- Quarkonia results in heavy ions from CMS, KAMIN, Jason Adrian
- Heavy quark spectroscopy, PAPPAGALLO, Marco
- Spectroscopy and decay properties with b-hadrons at the ATLAS experiment, TOMS, Konstantin
- LHCb early measurements focusing on B and Charm production, NEEDHAM, Matthew David
- $t\bar{t}$  + isolated photon production at NLO accuracy matched with parton shower, TROCSANYI, Zoltan Laszlo
- On NLL soft gluon corrections to the  $t\bar{t}$ -Higgs boson production at the LHC, MOTYKA, Leszek
- Top quark pair production measurements using the ATLAS detector at the LHC, YAMAUCHI, Katsuya
- Top quark pair properties using the ATLAS detector at the LHC, LEVY, Mark
- Measurement of t-channel single top quark production in pp collisions, ROCKER, Steffen
- 3-loop heavy flavor non-singlet contributions to different observables, BLUEMLEIN, Johannes
- Decay rate of the SM Higgs boson to bottom quarks at  $O(\alpha^2)$ , MIHAILA, Luminita
- Heavy quark production asymmetries at LHCb, GAULD, Rhorry Graham
- Searches for top/bottom partners and new phenomena in top/bottom quark pair signatures in CMS, MARGAROLI, Fabrizio
- Single Top quark production cross section and properties using the ATLAS detector at the LHC, PEDRAZA LOPEZ, Sebastian
- Heavy flavour theory developments, DE FREITAS, Abilio
- Heavy quark production results, NEEDHAM, Matthew David

# Contributions

---



- LHCb CP Violation in B decays, GRILLO, Lucia
- LHCb rare decays, CLIFF, Harry Victor
- LHCb exotics, PAPPAGALLO, Marco
- Measurements of heavy-flavour production in pp and p-Pb collisions with ALICE at the LHC, KIM, Minjung
- Higgs production in association with top quarks in CMS, MARGAROLI, Fabrizio
- Recent  $V$ +charm/beauty jets measurements from DZero, PRICE, Darren
- Measurements of heavy flavour production in association with  $W$  and  $Z$  bosons with the ATLAS detector, LAW, Alexander
- Vector boson production in association with jets and heavy flavor quarks from CMS, LUETIC, Jelena



# W+heavy quarks

A. Law, heavy flavor with W/Z ATLAS  
 J. Luetic, Vector boson + jets, CMS  
 D. Price, V+charm/bottom D0  
 Heavy Quarks

## W+b



Even larger difference observed in W+c in D0

# Rare decays

FCNC decays proceed via loops in the SM

- Highly suppressed (hence rare)
- Can receive significant modifications from NP

In this talk:

$b \rightarrow s$  transitions

$$B^0 \rightarrow K^{*0} \mu^+ \mu^- \quad \text{LHCb-CONF-2015-002}$$

$$B_s^0 \rightarrow \phi^0 \mu^+ \mu^- \quad \text{arXiv:1506.08777}$$

$$\Lambda_b^0 \rightarrow \Lambda^0 \mu^+ \mu^- \quad \text{JHEP 06 (2015) 115}$$



$b \rightarrow d$  transition

$$B^+ \rightarrow \pi^+ \mu^+ \mu^- \quad \text{Preliminary, LHCb-PAPER-2015-035 in prep.}$$

Search for dark bosons using

$$B^0 \rightarrow K^{*0} \chi^0 (\rightarrow \mu^+ \mu^-) \quad \text{arXiv:1508.04094}$$


All use Run I  $3\text{fb}^{-1}$  data set.



# Rare decays


Parameterise the decay in terms of three angles ( $\theta_l$ ,  $\theta_K$ ,  $\phi$ ) and  $q^2 = m_{\mu\mu}$

$$\frac{1}{d(\Gamma + \bar{\Gamma})/dq^2} \frac{d^3(\Gamma + \bar{\Gamma})}{d\Omega} \Big|_P = \frac{9}{32\pi} \left[ \frac{3}{4}(1 - F_L) \sin^2 \theta_K + F_L \cos^2 \theta_K \right. \\ \left. + \frac{1}{4}(1 - F_L) \sin^2 \theta_K \cos 2\theta_l \right. \\ \left. - F_L \cos^2 \theta_K \cos 2\theta_l + S_3 \sin^2 \theta_K \sin^2 \theta_l \cos 2\phi \right. \\ \left. + S_4 \sin 2\theta_K \sin 2\theta_l \cos \phi + S_5 \sin 2\theta_K \sin \theta_l \cos \phi \right. \\ \left. + \frac{4}{3} A_{FB} \sin^2 \theta_K \cos \theta_l + S_7 \sin 2\theta_K \sin \theta_l \sin \phi \right. \\ \left. + S_8 \sin 2\theta_K \sin 2\theta_l \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_l \sin 2\phi \right].$$



Can also calculate observables that have less form factor dependence e.g.

$$P'_{4,5} = S_{4,5} / \sqrt{F_L(1 - F_L)} \quad A_T^{(2)} = S_3 / (1 - F_L)$$






- Deviation from SM at  $2.9\sigma$  in  $[4.0-6.0]$  and  $[6.0-8.0]$   $\text{GeV}^2$
- Tension with SM at  $3.7\sigma$
- Possible explanations:  $Z'$ , leptoquark or charm-loops.

# Charmed Pentaquarks

## AMPLITUDE ANALYSIS OF $\Lambda_b \rightarrow J/\psi K p$

26k  $\Lambda_b^0$  candidates  
Background  $\sim 5.4\%$

[LHCb: PRL 115, 07201 (2015)]




Unexpected narrow peak in  $m(J/\psi p)$ !

- Efficiency flat over the “Dalitz” plot
- Cross checks:
  - ✓ Veto  $B_s \rightarrow J/\psi K K$  &  $B^0 \rightarrow J/\psi K \pi$  after swapping the mass hypothesis of the  $\Lambda_b$  daughters:  $p \leftrightarrow K$  or  $K \leftrightarrow \pi$
  - ✓ Clone and ghost tracks carefully removed
  - ✓ Exclude  $\Xi_b$  decays
- Could it be a reflection of the interfering  $\Lambda^*$ 's  $\rightarrow p K^-$ 
  - ✓ 6D unbinned maximum likelihood fit ( $m_{Kp}, \theta_{\Lambda_b}, \theta_{\Lambda^*}, \phi_K, \theta_\psi, \phi_\mu$ )


# ADDING $P_c \rightarrow J/\psi p$ AMPLITUDES

[LHCb: PRL 115, 07201 (2015)]


Extended  $\Lambda^*$  model



Extended  $\Lambda^*$  model + 1  $P_c^+$



Reduced  $\Lambda^*$  model + 2  $P_c^+$ 's



## Extended $\Lambda^*$ model:

- ...the fit projection can reproduce the peaking structure in  $J/\psi p$
- Adding non-resonant term,  $\Sigma^*$ 's or extra unknown  $\Lambda^*$ 's doesn't help

## Extended $\Lambda^*$ model + 1 Pentaquark decaying to $J/\psi p$



- Try all  $J^P$  of  $P_c^+$  up to  $7/2^\pm$
- Best fit has  $J^P = 5/2^\pm$ . Still not a good fit

## Reduced $\Lambda^*$ model + 2 Pentaquarks decaying to $J/\psi p$

- Obtain good fits even with the reduced  $\Lambda^*$  model
- Best fit has  $J^P = (3/2^-, 5/2^+)$ , also  $(3/2^+, 5/2^-)$  &  $(5/2^+, 3/2^-)$  are preferred

# Top measurements: single top

J. Maestro, Top quark, plenary  
 K. Yamauchi Top quark pair ATLAS  
 M. Levy Top quark pair ATLAS detector  
 S. Rocker Measurement of single top  
 Heavy Quarks



# Questions ...


Questions ...

Multiparton Dynamics




# What goes into description of $dn/d\eta$ ?

- basic process:  $gg \rightarrow gg \dots$
- parton densities at small  $x$  and small scales
- non-diffractive, diffractive processes (single diffractive, double diffractive dissociation)




- multiparton interactions:
  - w/o MPI  $dn/d\eta$  NOT described !



# The underlying event



- Measure particle density and average  $p_t$  in region transverse to trigger object (jet, Z...)

V. Cairo (ATLAS), Min Bias & UE  
 A. Bruni, Soft QCD, plenary  
 R. Field, Soft QCD, plenary





- more activity than predicted from parton shower (higher orders) and hadronization observed

➔ introduce MPI:



# Underlying event measurements at 13 TeV

V. Cairo (ATLAS), Min Bias & UE



- None of the models describes the initial rise well
- From 10 GeV quite good description for the UE Tunes
- EPOS 15% off in the plateau

September 1-5, 2015


V. Cairo

17

# Multi-parton interaction — why ?

- The basic problem for  $p_t \rightarrow 0$

T. Sjöstrand and P. Skands. Multiple interactions and the structure of beam remnants. JHEP, 03:053, 2004.




# Multi-parton interaction — why ?

- The basic problem for  $p_t \rightarrow 0$

T. Sjöstrand and P. Skands. Multiple interactions and the structure of beam remnants. JHEP, 03:053, 2004.


- Can we measure this ?



# Multi-parton interaction — why ?


- The basic problem for  $p_t \rightarrow 0$

T. Sjöstrand and P. Skands. Multiple interactions and the structure of beam remnants. JHEP, 03:053, 2004.



- Can we measure this ?


arXiv:1507.00233



# Multi-parton interaction — why ?

- The basic problem for  $p_t \rightarrow 0$

T. Sjöstrand and P. Skands. Multiple interactions and the structure of beam remnants. JHEP, 03:053, 2004.




- Taming of x-section with:

$$\frac{\alpha_s^2(p_{\perp 0}^2 + p_{\perp}^2)}{\alpha_s^2(p_{\perp}^2)} \frac{p_{\perp}^4}{(p_{\perp 0}^2 + p_{\perp}^2)^2}$$

- determine  $p_{t0}$  from tunes

- Can we measure this ?

arXiv:1507.00233




# Multiparton interactions - double parton scattering



- Quarkonium+W as a clean probe for DPS

D. Bertsche (ATLAS), Associated quarkonium Multiparton Dynamics

## J/ψ+W results: azimuthal opening angle



DPS processes are expected to give a flat azimuthal opening angle ( $\Delta\phi$ ) distribution



Possible indication of DPS and SPS J/ψ+W production.




- Very clean process for DPS (in principle)
- What about theory predictions for
  - SinglePartonScattering, DoublePartonScattering ?



# Multiparton interaction and diffraction

D. Britzger (H1), Hard diffraction at HERA  
Multiparton Dynamics

- Hard diffraction and diffractive parton densities measured at HERA, assuming hard scattering factorization in DIS
- Multiparton interaction can destroy diffractive processes
  - gap - survival probability
- Not at all implemented in any MC simulation
- Factorization tests with dijets at HERA in  $\gamma p$  and DIS



# Questions in Soft QCD & Multiparton Dynamics

---

- Do we have really (really !!!) evidence for MPI and/or DPS ?
  - all what we measure is more activity than predicted within a certain framework:
    - collinear factorization and collinear PDFs
- How can diffraction be consistently included into calculations/simulations ?
  - if all comes from high parton densities, they must be directly related
  - could have significant effects on hadronization parameters, tuning etc.
- In UE tuning, a significant fraction of ColorReconnection is needed:
  - how can color-rearrangement be understood if we believe QCD should work down to small scales ?
    - similar issue arises in Quarkonium production.


# Questions ...

# PDFs

# HERA 1 & 2 — combined data

M. Cooper Sakar, Proton structure, plenary


## H1 and ZEUS



M.

- Combine HERA1 & 2 constrains PDFs at large  $x$


## H1 and ZEUS



see also parallel talks by:  
 HARLAND-LANG, Lucian  
 STUMP, Daniel  
 MELNITCHOUK, Wally  
 ISARKAR, Amanda

# The small x side


- news from precision HERA data fits



- This has relation to min bias, UE, pileup at LHC

M. Cooper Sakar, Proton structure, plenary  
 P. Newman, LHeC, plenary  
 C. Gwenlan, PDFs at LHeC, PDF  
 N. Armesto, smallx physics LHeC, Multiparton  
 N. Armesto, Nuclear PDFs LHeC, PDF

- how to address the small x side ?
  - LHeC, FCeC ?




# Needs to go beyond collinear PDFs ?

- at small  $x$ , transverse momentum dependent (TMD) pdfs are needed, in CCFM etc.
- at larger  $x$ , TMD are used & useful to describe DY, Higgs  $q_T$  spectrum, for example in SCET and other approaches.
- MC generators with parton showers generate effectively TMDs, even when using in NLO matched calculations.

## POWHEG Reshuffling

From P.Nason: TOPLHCWG, 22 May 2014, <https://indico.cern.ch/event/301787>

The cause: **MOMENTUM RESHUFFLING**



**ISR shower** throws off shell the incoming gluon. In order to conserve 4-momentum, the final state is boosted. The mass  $m_{t\bar{t}}$  is preserved

**FSR shower** changes the mass of final state partons. In order to conserve 4-momentum, the final state momenta are rescaled.

# Questions ...

## hard QCD

# Kinematic effects in NLO + PS


- Parton shower effects in Jet distributions

S. Dooling, et al  
Phys.Rev., D87:094009, 2013.

$$K^{\text{NP}} = N_{\text{NLO-MC}}^{(\text{ps+mpi+had})} / N_{\text{NLO-MC}}^{(\text{ps})}$$

$$K^{\text{PS}} = N_{\text{NLO-MC}}^{(\text{ps})} / N_{\text{NLO-MC}}^{(0)}$$

- Sizable effects from Parton Shower
- Not necessarily covered by PDF and scale uncertainties
- How to estimate uncertainties ?



- Top quark pt-spectrum - reweighting
  - most effect comes from parton shower





# Threshold resummation

- Activities in large x threshold resummation:

$$\frac{1}{1-x}$$

E. Laenen, Developments, plenary  
 L. Rottoli, Parton Distributions, PDFs  
 H. Yoo, Searches CMS  
 A. Elliot Searches ATLAS  
 Multiparton Dynamics



NNPDF3.0 DIS+DY+Top,  $Q^2=10^4 \text{ GeV}^2$



- To observe large x resummation, need a different scenario: define  $x' = \frac{m_{\mu\mu}}{m_{vis}}$  with visible mass accessible in detector.
- towards measurements at kinematic limit !

# Higgs differential x-section measurements

L. Tompkins, QCD in Higgs and BSM, plenary  
 A. Zghiche, Higgs from CMS, Hard QCD  
 B. Liu Higgs from ATLAS, Hard QCD




- Naively one would expect to see a maximum at  $\sim 20$  GeV, a drop at small  $p_T$ .
  - it is hidden in the acceptance cuts ... MC describes it, but ?
- Discussion issue: should experimental results be fully corrected, or can they even be at Detector level since we anyway compare to MC-Rivet etc ?
  - provide full resolution matrix, without loss of Infos ? **Need discussion in community**

Questions ...

Quarkonium Production


# Are color-octet states in Quarkonium needed ?

Figures by Pietro Faccioli



CMS Phys.Lett. B727 (2013) 101

- color-octet states introduced to describe  $p_t$  spectrum in collinear factorization.
- in  $kt$  - factorization the contribution from color-octet is much smaller (if needed at all)
- How are color-octet states in Quarkonium production connected to color-reconnection “needed” in UE tunes ?
  - it’s both soft gluons, but .... ?



# What can we conclude ?

---


- Huge progress is made on both experimental and theory/pheno side to understand, what is measured at LHC.
- LHC run 2 has just started, and we already have seen impressive new results, just after a few month of data taking.
- Many very interesting and challenging results will come out in the near future, which will occupy all our resources, from the experimental and theoretical side.

# What can we conclude ?

---

- Huge progress is made on both experimental and theory/pheno side to understand, what is measured at LHC.
- LHC run 2 has just started, and we already have seen impressive new results, just after a few month of data taking.
- Many very interesting and challenging results will come out in the near future, which will occupy all our resources, from the experimental and theoretical side.
- In a bigger picture, apart from precision measurements and precision calculations, **for me**, two fundamental questions remain:
  - high energy behavior of QCD
  - color structure of QCD — color confinement

# Instead of a summary



A very colorful time  
is ahead of us !



But this is not the  
end...

# Thank so much, local organizers !

---

- Thanks for hosting this very nice, interesting and stimulating conference at Queen Mary !

# Thank so much, local organizers !

---

- Thanks for hosting this very nice, interesting and stimulating conference at Queen Mary !
- Thanks so much for the excellent organization

# Thank so much, local organizers !

---

- Thanks for hosting this very nice, interesting and stimulating conference at Queen Mary !
- Thanks so much for the excellent organization
  - except the weather !!!!

# Thank so much, local organizers !

---

- Thanks for hosting this very nice, interesting and stimulating conference at Queen Mary !
- Thanks so much for the excellent organization
  - except the weather !!!!
- Thanks so much for the great reception and dinner and excursion

# Thank so much, local organizers !

---

- Thanks for hosting this very nice, interesting and stimulating conference at Queen Mary !
- Thanks so much for the excellent organization
  - except the weather !!!!
- Thanks so much for the great reception and dinner and excursion
- Thanks very much to the conveners for preparing and chairing very good, lively and fascinating sessions:
  - Hard QCD: Evelin Meoni / Joao Pires
  - PDFs: Jun Gao / Ringaile Placakyte / Lucian Harland-Lang
  - Multi-parton dynamics: James Robinson / Nestor Armesto / Leszek Motyka
  - Heavy Quarks: Fred Olness / Philip Ilten

# Thank so much, local organizers !

---

- Thanks for hosting this very nice, interesting and stimulating conference at Queen Mary !
- Thanks so much for the excellent organization
  - except the weather !!!!
- Thanks so much for the great reception and dinner and excursion
- Thanks very much to the conveners for preparing and chairing very good, lively and fascinating sessions:
  - Hard QCD: Evelin Meoni / Joao Pires
  - PDFs: Jun Gao / Ringaile Placakyte / Lucian Harland-Lang
  - Multi-parton dynamics: James Robinson / Nestor Armesto / Leszek Motyka
  - Heavy Quarks: Fred Olness / Philip Ilten
- Thanks to all the speakers in the parallel and plenary sessions for their excellent talks

# And many, many thanks to ...

---

... the Local organizing team:

- Eram Rizvi
  - Agni Bethani
  - Marcella Bona
  - Teppei Katori
  - Dan Traynor
- with the students and helpers
  - Jazmina Vaca Ortiz
  - Lewis Armitage
  - Lewis Millward
  - Tamsin Nooney
  - Tom Stevenson
  - Andres Ramirez Morales
  - Ryan Hill
  - Aston Dawsmith
  - Kathryn Coldham

Many Thanks!