



University of Michigan
Medical School

A Hypercellular Component of Glioblastoma Identified by High b- Value Diffusion Weighted Imaging

Y. Cao, D. Wahl, P. Pramanik, M. Kim, T.
Lawrence, H. Parmar

*Departments of Radiation Oncology, Radiology, Neurology and
Biomedical Engineering*

The University of Michigan, Ann Arbor, MI, USA

Supported in part by US NIH RO1NS064973



Introduction



- Standard Tx of GBM is resection, followed by concurrent RT (60 Gy in 30 fx) and TMZ, and 6-12 cycles of TMZ
- The majority of the patients fails locally.



Introduction

- NRG BN001 trial aims to determine whether intensified radiation therapy with concurrent TMZ followed by adjuvant TMZ improve OS?
 - ✓ It is important to stratify the patients based upon radiosensitivity using genetic biomarkers
 - ✓ **Where to boost is matter**
 - ✓ Is the contrast enhanced GTV the effective target to be boosted?



Post Gd T1



T2 FLAIR

Hypercellularity of GBM Identified by High b-value Diffusion Imaging

- Microscopic mobility of water in the cellular environment
 - ✓ Sensitive to cellular density, water content and organization of the cellular and extracellular space
- Preclinical studies in 1990s: ADC is inversely correlated with cellularity
- **Unpredictable ADC in GBM**
A **single voxel in GBM** is consisted of tumor cells (low ADC), edema (high ADC), micro-necrosis (high ADC), normal tissue (normal ADC)





Purpose

- We aim to suppress MR signals from edema, micro-necrosis and even normal tissue to reveal hypercellular (HC) tumor by increasing diffusion weighting (high b-value).
- MR signals decay exponentially with diffusion coefficients and diffusion weighting (b-value).



Patients


- 44 patients (age: 23-79 years) with GBM
 - ✓ 35 patients with newly diagnosed GBM
 - Received concurrent RT+TMZ, followed by adjuvant TMZ
 - Radiation planning based upon conventional MRI (median dose of 60 Gy in 30 fx)
- MRI scans prior to RT but post-resection on a 3T scanner
 - ✓ Conventional post-Gd T1W and T2 FLAIR images
 - ✓ DWI with $b=0, 1000, (2000)$ and 3000 s/mm^2 (~ 5 min)
- Conventional MRI at routine f/u and recurrence

Questions


- ✓ Spatial relationship between **FLAIR AV** and hypercellularity volume (**HCV**) identified by DW imaging at $b=3000$
- ✓ Non-enhanced HCV
- ✓ Relationship between **HCV** and patterns of failure
- ✓ Prognostic values of **HCV** for PSF and OS




Hypercellularity




Post-Gd T1WI




FLAIR




Conventional ADC



DWI b=1000



DWI b=2000



DWI b=3000

Red: post-Gd GTV

Green: FLAIR AV




Yellow: b3000 HCV

Blue: normal-appearing tissue





University of Michigan
Medical School

HCV is 10 times smaller than FLAIR AV!



Red: post-Gd GTV
Cyan: b3000 HCV
Green: FLAIR AV

Nonenhanced HCV




Red: post-Gd GTV
Cyan: b3000 HCV

77% (27) patients had non-enhanced HCVs > 1 cc
23% (8) patients had non-enhanced HCVs > 5 cc




University of Michigan
Medical School


Radiation Dose Coverage



Post-Gd T1WI



FLAIR



DWI b=3000

Pink: 95PDV

Red: post-Gd GTV

Green: FLAIR AV






Yellow: b3000 HCV

A small number of patients (n=3) had 1 cc or greater HCV that was not covered by 95PDV



Pattern of Failure

Uni



b=3000 DWI
Pre-RT

b=3000 DWI
End of RT

Post-T1W Image







19 months post-RT

b=3000 DWI





University of Michigan
Medical School

Cross-Comparison







HCV vs Pattern of Failure



Red: post-Gd GTV
Green: FLAIR AV









Cyan: b3000 HCV
Blue: normal-appearing tissue





HCV vs Pattern of Failure

University of Michigan
Medical School




Red: post-Gd GTV
Green: FLAIR AV
Yellow: HCV
Cyan: reGTV_Gd



University of Michigan
Medical School

Early Detection of Progression



Pre-RT

Week 3
During RT

Week 6
During RT

1M post RT

Progression and Survival

➤ 35 Patients

✓ Progression

- ✓ 28 patients had progression clinically

 - ✓ Median PSF of 12 months

- ✓ 6 had no progression and 1 dead without progression

✓ Survival

- ✓ 22 patients were dead

 - ✓ Median OS of 17.5 months

- ✓ 13 were alive with a median followup of 20 months

Prediction for PSF and OS

- The HCV pre-RT predicted PSF and (p<0.003) and OS (p<0.05, univariate Cox regression)
- Considering MGMT status in a multivariate Cox regression model, the HCV pre-RT was still a significant predictor for OS (p<0.002)

Discussions and Conclusion


- The HCV identified by high b-value diffusion imaging is aggressive.
- Using FLAIR AV to define non-enhanced tumor could result in high-risk of radiation toxicity.
- High b-value DW imaging could provide a means to extract hypercellularity components of GB, and define radiation boosting target and assess response.
- A phase II trial is in preparation.




University of Michigan
Medical School

Results: Dose Coverage & HCV Changes

Pre-RT



Last Week RT




Blue: 95PDV of 72 Gy




MRI Characteristics and Changes

University of Michigan
Medical School

Post Gd
T1WI



T2
FLAIR



RT + TMZ

TMZ

Evastine/Drug

drugs + re-RT

Pre-RT

4M post-RT

6M post-RT


16M post-RT




MRI Characteristics and Changes

University of Michigan
Medical School

Post Gd
T1WI



T2
FLAIR



Tumor?

RT + TMZ

TMZ

Evastine/Drug

drugs + re-RT

Pre-RT

4M post-RT

6M post-RT

16M post-RT