


Radiation therapy and anti HPV treatments

Eric Deutsch

Gustave Roussy Cancer Campus




HPV



Le Papillomavirus humain (HPV)

- More than 98% cervix cancer, (IARC 2007).
- 85% of squamous cell anal cancer, (Daling JR, Cancer 2004).
- 50% of penile cancer, (Hernandez BY, Cancer. 2008).
- 33-72% of oropharyngeal cancers (Kreimer AR, Cancer Epid Biomarkers, 2005).



→ 26,000 cancers each year, 18,000 females
and 8,000 males (WU, JAMA 2012).

HPV & cancer : Clinical impact?



- Prevention
(primary vaccination, secondary pap smear..)




HPV & cancer : Clinical impact?

- Prevention (primary vaccination, secondary pap smear..)
- Biomarker : predictive and prognostic

HPV as a biomarker in head and neck cancers :




HPV negative :
Response less
likely





Radiation sensitizer

HPV Positive :
Response likely
Optimise radiotherapy
Decrease late effects



OS by UICC/AJCC TNM Stage (7th edition)




HPV-positive	Events / Total	3 years	5 years	<i>p</i> value	HPV-negative	Events / Total	3 years	5 years	<i>p</i> value
Total	164 / 573	80% (77-84)	74% (70-78)		Total	161 / 237	49% (42-55)	38% (32-45)	
Stage I	1 / 8	88% (67-100)	88% (67-100)		Stage I	4 / 8	88% (67-100)	70% (42-100)	
Stage II	5 / 25	87% (75-100)	78% (62-97)		Stage II	16 / 31	67% (53-86)	58% (42-80)	
Stage III	27 / 79	81% (72-90)	71% (62-83)		Stage III	24 / 38	63% (49-80)	50% (36-70)	
Stage IV	131 / 461	80% (76-84)	74% (70-78)	0.560	Stage IV	117 / 160	39% (32-48)	30% (23-39)	0.004

Courtesy Brian O'Sullivan

Proposed Anatomic Stage Grouping

Recursive Partitioning Analysis (RPA) model:

- Based on an optimized binary partition of ordinal T- or N-categories
- Derived *stage* groups with minimal hazard difference within same stage



- Better performance than alternative AHR Model
- Consistent with current de-intensification trial criteria designed on the basis of the disease behaviour by T and N criteria (eg. NRG HN 02)
- Better “Grid” than AHR Model

Huang, Wei, O’Sullivan et al JCO (accepted)

Courtesy Brian O’Sullivan


HPV & cancer :

Clinical impact?

- Prevention (primary vaccination, secondary pap smear..)
- Biomarker : predictive and prognostic
- Development of tailored therapies :
pharmacological agents?
therapeutic vaccines?

Cidofovir :

mechanism of action the antiviral agent







■ (S)-9-(3-hydroxy-2-phosphonylmethoxypropyl) Cytosine: [(S)-HPMPC]

■ Acyclic Phosphonate nucleosid

■ Half-life *in vivo* : 48 h


- Johnson JA, Antimicrob Agents Chemother 1999;43(5):1198-205.
- Andrei G, Oncol Res 1998;10(10):523-31.
- Andrei G, Oncol Res 1998;10(10):533-9.

Me180 cell line



Cidofovir : translational approach

Cidofovir



Oncogene

2001,2002

Lancet Oncol

2003

Rad Res

2006

Plos One

2009

« HPV-RX » : antiviral agent + chemoradiation phase I trial

J_0 → DLT

S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	
CDV	CDV		CDV		CDV		CDV		CDV	
			External beam 45 Grays (fraction hebdo : 1,8 Grays)						Brachy 15 Gy	
			chemotherapy C Paraplatine® AUC=2.5						chemo C	

**Tumor
biopsy**



C= Weekly Carboplatin, AUC=2,5


CDV=Weekly Cidofovir

HPV vaccines

Two HPV vaccines are currently on the market: Gardasil (Merck) and Cervarix (Glaxo).

Both vaccines are based on recombinant **L1 viral protein**, assembled in virus-like particles (VLPs) and protect **against high-risk HPV 16 and 18**, responsible of around 60-70% of cervical cancer.

Gardasil also protects **from low risk HPV 6 and 11**, responsible of the majority of genital warts.




Partridge EE, Med Ed


BOTH ARE PROPHYLACTIC VACCINES ONLY!!

HPV-E7 vaccine

Targets antigen presenting cells.
Shigatoxine receptor
expressed on dendritic cells







Nature Reviews | Immunology





Johannes, Gene Ther 2005

Combination treatment by local radiation and therapeutic STxB-E7 vaccine elicits an effective antitumor response.






Adjuvant effect? Threshold dose?




A





B



Fractionated irradiation?




CD8+ cells into the tumor?





CD8+ lymphocytes linked with response?

CD8+
lymphocytes
inversely
correlate with
the tumor size




Do CD8+ cells play a role?






Mecanism of synergy ?

Any impact on immune check-points?






- Synergy between immune check-point blockers and IR blocker reported to be fractionation dependant (Formenti, JNCI 2013).
- PD1+, TIM3 negative lymphocytes infiltrates « favorable » in HPV positive tumor context (Badoual Can Res 2013).



PD1, CTLA4 and Tim3 expression on CD8⁺ T cells not modulated by combination treatment.



Duration of anti tumor response?



Long term local or systemic immune response?







Increased vascular permeability upon combined treatment


Miles Assay





Evans blue tail injection




Effect of treatments on tumor vessels




aSMA staining
(pericytes marker)



Effect of treatments on tumor vessels




NG2 staining
(pericytes marker)




ICAM-1 staining increased by combined treatment.


Control




Vaccine







IR




Vaccine + IR



CD31 staining



Synergy between anti-angiogenic and immunotherapy already suggested...



...similar case for radiotherapy??

Conclusions

- Combined HPV vaccine/irradiation therapy promotes tumor regression and greatly improve survival.
- Cytotoxic T-cell response against HPV antigen is required for anti-tumor efficacy and is greatly induced by combined therapy.
- Effective combined therapy protects from recurrences and metastases, through induction of T cell memory.
- Combined therapy promotes tumor vessel remodeling and increases perfusion

Are all vaccines even?

Recently, two vaccines targeting HPV antigens have been tested in association with radiotherapy in murine models (Wu, 2014, Draghiciu 2014). However, though HPV-associated cancers are located in mucosal sites, both studies were conducted using subcutaneous tumors.

Recent data indicate that the site of tumor implantation affects the response to cancer vaccine and the homing of induced CD8⁺ cells (Sandoval, Sci Trans Med 2013).

No back to back comparison available.

Acknowledgments

C Haie
P Morice
C Lhomme
P Pautier

HEGP, Paris :
Eric Tartour

Curie, Paris :
Suzie Scholl
Ludger Johanes

INSERM U1030
“Molecular radiotherapy”

Michele Mondini
JL Perfettini

INSERM U981

J Adam
E Louvet

-
FP7 RAIDS