

Extraction of the Strong Coupling α_s Through W-boson Hadronic Decays

[Workshop on high-precision α_s measurements: From LHC to FCC-ee]

Matej Srebre¹, David d'Enterria

CERN


matej.srebre@cern.ch, dde@cern.ch

October 2015

¹University of Ljubljana, currently master's student at LMU Munich

Motivation

Extraction of α_s through comparison of various experimental observables to different perturbative QCD predictions.



- 1 Hadronic τ decays: $R_\tau = \frac{\Gamma(\tau^- \rightarrow \nu_\tau + \text{hadrons})}{\Gamma(\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e)} = S_{\text{EW}} N_C (1 + \sum_{n=1}^4 c_n (\frac{\alpha_s}{\pi})^n + \mathcal{O}(\alpha_s^5) + \delta_{np})$ (N³LO)
- 2 Lattice QCD: Various short-distance quantities: $K^{\text{NP}} = K^{\text{PT}} = \sum_{i=0}^n c_i \alpha_s^i$ (NNLO)
- 3 Hadronic Z decays: $R_Z = \frac{\Gamma(Z \rightarrow \text{hadrons})}{\Gamma(Z \rightarrow \ell)}$ = $R_Z^{\text{EW}} N_C (1 + \sum_{n=1}^4 c_n (\frac{\alpha_s}{\pi})^n + \mathcal{O}(\alpha_s^5) + \delta_m + \delta_{np})$ (N³LO)
- 4 $e^+e^- \rightarrow q\bar{q}(g)$: Event-shapes, jet rates: $\frac{1}{\sigma} \frac{d\sigma}{dY} = \frac{dA}{dY} \hat{\alpha}_s + \frac{dB}{dY} \hat{\alpha}_s^2 + \frac{dC}{dY} \hat{\alpha}_s^3$ (NNLO)
- 5 $e^\pm p \rightarrow \text{hadrons}$ (PDF): $\sigma(\text{jet}), \frac{\partial}{\partial \ln Q^2} D_i^h(x, Q^2) = \sum_j \int_x^1 \frac{dz}{z} \frac{\alpha_s}{4\pi} P_{ji}(\frac{x}{z}, Q^2) D_j^h(z, Q^2)$ (NLO, NNLO)
- 6 $pp, p\bar{p} \rightarrow t\bar{t}, \text{jets}$ (NNLO, NLO)
- 7 Hadronic W decays:
 - What is the theoretical and experimental status?
 - Can we extract α_s through hadronic W decays? With which precision?

Theoretical Calculation of $\Gamma_W(\text{hadronic})$

The hadronic W-boson decay width has not been used so far for α_s extraction because:

- a complete N³LO/NNLO formula with all computed corrections [1] was not available until recently [2] (albeit with a few approximations),
- the 2% relative experimental uncertainty on $\Gamma_W(\text{hadronic})$ was significantly large compared to 0.1% of $\Gamma_Z(\text{hadronic})$.

We recalculated $\Gamma_W(\text{hadronic})$ through implementation in *MATHEMATICA* the $\mathcal{O}(\alpha_s^4)$ or N³LO formula using [2]:

$$\Gamma_W(\text{hadronic}) = \sum_{i=0}^4 \Gamma_{\text{QCD}}^{(i)} + \Gamma_{\text{EW}}^{(1)} + \Gamma_{\text{Mixed}}^{(2)} = \frac{\sqrt{2}G_F m_W^3}{4\pi} \left[\sum_{i,j} |V_{i,j}|^2 \right] \left(1 + \sum_{k=1}^4 \left(\frac{\alpha_s}{\pi} \right)^k + \delta_{\text{EW}} + \delta_{\text{Mixed}} \right),$$

where

- $\Gamma_{\text{QCD}}^{(k)}$ is the leading order decay width and QCD corrections of order $\mathcal{O}(\alpha_s^k)$ and $k = 1, \dots, 4$,
- $\Gamma_{\text{EW}}^{(1)}$ electroweak corrections of order $\mathcal{O}(\alpha)$,
- $\Gamma_{\text{Mixed}}^{(2)}$ mixed corrections of order $\mathcal{O}(\alpha\alpha_s)$.

[1] - A. Denner, B. Kniehl, J. Kühn, K. Chetyrkin, ...

[2] - D. Kara, Nucl. Phys. B 877, 3 (2013)

Improvements with Respect to Previous Γ_W (hadronic) Calculations

In our calculations we carry out the following improvements compared to previous works:

- 1 implement **finite quark masses** in the dominant Γ_W (hadronic) terms: Born and first-order QCD corrections,
- 2 use **NNLO α_s running instead of LO** (between m_W and m_Z),
- 3 use **current PDG world average values** for parameters of the Standard Model (α_{QED} , G_F , m_q , m_ℓ , m_W , m_Z , m_H , *CKM matrix elements* $|V_{i,j}|$),
- 4 determination of associated **theoretical** and **parametric** uncertainties.

Hadronic W-boson Decay Width - Numerical Results

Previous state-of-the-art [2]: $\Gamma_W(\text{hadronic}) = (1458.820 \pm 6 \times 10^{-3}) \text{ MeV}$,

$$\Gamma_W(\text{hadronic}) = (1428.803 \pm 0.030_{\text{theor.}} \pm 22.608_{\text{param.}}) \text{ MeV},$$

$$\Gamma_W(\text{hadronic}, V_{ij} V_{kj} = \delta_{ik}) = (1411.546 \pm 0.030_{\text{theor.}} \pm 0.742_{\text{param.}}) \text{ MeV}.$$

Partial width	$\Gamma_{\text{QCD}}^{(0)}$	$\Gamma_{\text{QCD}}^{(1)}$	$\Gamma_{\text{QCD}}^{(2)}$	$\Gamma_{\text{QCD}}^{(3)}$	$\Gamma_{\text{QCD}}^{(4)}$	$\Gamma_{\text{EW}}^{(1)}$	$\Gamma_{\text{Mixed}}^{(2)}$
$\Gamma_W(\text{hadronic})$ of [2]	1408.980	54.087	2.927	-1.018	-0.245	-5.132	-0.779
$\Gamma_W(\text{hadronic})$	1379.851	53.080	2.873	-1.000	-0.241	-5.002	-0.757
$\Gamma_W(\text{hadronic}, V_{ij} V_{kj} = \delta_{ik})$	1363.186	52.439	2.838	-0.988	-0.238	-4.942	-0.749


Numerical values of the partial decay widths. All values given in MeV.

The following uncertainties are present in $\Gamma_W(\text{hadronic})$:

- **parametric uncertainty** (modifying all PDG parameters by $\pm\sigma$, adding changes in quadrature)
 - $\pm 22.608 \text{ MeV}$ (the dominant parametric uncertainty is V_{cs}),
 - $\pm 0.742 \text{ MeV}$ for $V_{ij} V_{kj} = \delta_{ik}$ (dominated by m_W),
- **higher-order corrections** (assumed equal to Γ_Z , from $N^3\text{LO}$ to $N^4\text{LO}$): $\pm 0.0195 \text{ MeV}$,
- **non-perturbative uncertainties** (assuming power-corrections of order $\mathcal{O}\left(\frac{\Lambda^4}{m_W^4}\right)$): $\pm 7 \times 10^{-8} \text{ MeV}$,
- **finite quark masses beyond LO corrections**: $\pm 0.0042 \text{ MeV}$,
- **mixed corrections from [2]**: $\pm 0.006 \text{ MeV}$.


$\Gamma_W(\text{hadronic})$: Historical Data Versus Theory

Evolution of the PDG world average value of $\Gamma_W^{\text{EXP}}(\text{hadronic})$ (current value is (1405 ± 29) MeV) by year compared to theoretically predicted decay widths.



Extraction of α_s from $\Gamma_W(\text{hadronic})$

Using the $\mathcal{O}(\alpha_s^4)$ W-boson decay width formula we can extract α_s by comparing it to the experimental value which is $\Gamma_W^{\text{EXP}}(\text{hadronic}) = (1405 \pm 29) \text{ MeV}$.



\Rightarrow Current large parametric ($\pm 23 \text{ MeV}$) and experimental ($\pm 29 \text{ MeV}$) uncertainties on $\Gamma_W(\text{hadronic})$ propagate into a **huge α_s uncertainty $\sim 60\%$** .

Experimental priorities should be:

- measure $|V_{cs}|$ with better precision (current **1.6%**),
- significantly reduce uncertainty of $\Gamma_W(\text{hadronic})$ measurement to a **few MeV**,
- reduce m_W uncertainty (now it propagates to $\pm 0.8 \text{ MeV}$ on $\Gamma_W(\text{hadronic})$).

	$\alpha_s(m_W^2)$	$\alpha_s(m_Z^2)$
Experimental CKM	$0.0691 \pm 0.0513_{\text{param.}} \pm 0.0644_{\text{exp.}}$	$0.0683 \pm 0.0509_{\text{param.}} \pm 0.0638_{\text{exp.}}$
Unit CKM, $V_{ij}V_{kj} = \delta_{ik}$	$0.1071 \pm 0.0017_{\text{param.}} \pm 0.0664_{\text{exp.}}$	$0.1053 \pm 0.0016_{\text{param.}} \pm 0.0657_{\text{exp.}}$

Alternative Approach: α_s Extraction via $BR_W(\text{hadronic}) = \frac{\Gamma_W(\text{hadronic})}{\Gamma_W(\text{total})}$

$\Gamma_W(\text{hadronic})$ has a 2% experimental uncertainty, $BR_W(\text{hadronic})$ has an uncertainty of 0.4%.

\Rightarrow We try $BR_W(\text{hadronic}) = \frac{\Gamma_W(\text{hadronic})}{\Gamma_W(\text{total})}$ to extract α_s instead of $\Gamma_W(\text{hadronic})$.

For the total decay width $\Gamma_W(\text{total})$ we use the ZFitter NNLO (includes up to $\mathcal{O}(\alpha_s^3)$ QCD, $\mathcal{O}(\alpha)$ electroweak and $\mathcal{O}(\alpha\alpha_s)$ mixed corrections) fitted result by [3] which is parametrized as

$$\Gamma_W(\text{total}) = G_W^0 m_W^3 \quad \text{and} \quad G_W^0 = 4.0279 \times 10^{-6} (1 + 0.00095x_H - 0.0024x_H^2 + 0.0016x_H^3 + 000065x_s) \text{ GeV}^{-2},$$

where $x_s = f(\alpha_s)$, $x_H = f(m_H)$.


We also computed the associated parametric uncertainties as done for $\Gamma_W(\text{hadronic})$.

$\Gamma_W(\text{hadronic})$	$(1428.803 \pm 22.638) \text{ MeV}$
$\Gamma_W(\text{hadronic}, V_{ij}V_{kj} = \delta_{ik})$	$(1411.546 \pm 0.772) \text{ MeV}$
$\Gamma_W(\text{total})$	$(2093.591 \pm 1.172_{\text{param.}}) \text{ MeV}$
$BR_W(\text{hadronic})$	$0.6825 \pm 0.0108_{\text{param.}}$
$BR_W(\text{hadronic}, V_{ij}V_{kj} = \delta_{ik})$	$0.67422 \pm 0.00003_{\text{param.}}$
$BR_W(\text{hadronic})^{\text{EXP}}$	0.6741 ± 0.0027

[3] - G. C. Cho, K. Hagiwara, Y. Matsumoto and D. Nomura, JHEP 111, 068 (2011)

$BR_W(\text{hadronic})$: Historical Data Versus Theory


Evolution of the PDG world average value of $BR_W(\text{hadronic})$ (current value is 0.6741 ± 0.0027) by year compared to theoretically predicted decay widths.



Alternative Approach: α_s Extraction via $BR_W(\text{hadronic}) = \frac{\Gamma_W(\text{hadronic})}{\Gamma_W(\text{total})}$

We can extract α_s by comparing the theoretical hadronic branching ratio formula to the experimental world average value

$$\frac{\Gamma_W(\text{hadronic})(\alpha_s)}{\Gamma_W(\text{total})(\alpha_s)} = BR_W(\text{hadronic})^{\text{EXP.}}$$



⇒ Setting the CKM matrix to unit matrix instead of using experimental values, we can extract α_s with **35%** uncertainty.



To extract α_s with a higher precision:

- reduce the uncertainty of $|V_{cs}|$ as mentioned earlier,
- reduce the uncertainty of m_W (measured to **0.02%**) which becomes dominant once the $|V_{cs}|$ uncertainty is reduced below **0.05%**,
- measure $BR_W(\text{hadronic})$ with a better precision than today (**0.4%** now).

	$\alpha_s(m_W^2)$	$\alpha_s(m_Z^2)$	$\Delta\alpha_s(m_W^2)$
Experimental CKM	$-0.0030 \pm (> 0.1505_{\text{param.}}) \pm 0.0353_{\text{exp.}}$	/	/
Unit CKM	$0.1189 \pm 0.0004_{\text{param.}} \pm 0.0433_{\text{exp.}}$	$0.1167 \pm 0.0004_{\text{param.}} \pm 0.0430_{\text{exp.}}$	$\pm 37\%$

Extra: Indirect Determination of $|V_{cs}|$

The large experimental $BR_W(\text{hadronic})^{\text{EXP}}$ uncertainty precludes an accurate extraction of α_s , but we can use $BR_W(\text{hadronic})^{\text{EXP}}$ to determine $|V_{cs}|$ (fixing α_s to world average).



Extraction method	$ V_{cs} $ value
$\Gamma_W(\text{hadronic})$	$0.969 \pm 0.002_{\text{param.}} \pm 0.021_{\text{exp.}}$
$BR_W(\text{hadronic})$	$0.973 \pm 0.002_{\text{param.}} \pm 0.004_{\text{exp.}}$

\Rightarrow We can extract $|V_{cs}|$ with an uncertainty of **0.6%** compared to **1.6%** of the experimental measurement $|V_{cs}|^{\text{EXP}} = 0.986 \pm 0.016$.

Future Perspectives: LHC & FCC-ee

To determine α_s with a **higher precision** we need more precise measurement of $\Gamma_W(\text{total})$ and/or $BR_W(\text{hadronic})$ with reduced uncertainties.

- Uncertainties at LHC (5×10^5 high- m_T W's at $\sqrt{s} = 8$ TeV, 20 fb^{-1}):
 - Statistical: ~ 3 MeV (30 MeV at Tevatron, with 5×10^3 high- m_T W's)
 - Systematics: ~ 15 MeV (down from ~ 40 MeV at Tevatron, reduced PDF uncertainties)
- Improved result:** $\Gamma_W(\text{hadronic}) \sim (1429 \pm 12)$ MeV (i.e. **0.8%** uncertainty instead of **2%**)

	$\alpha_s(m_W^2)$	$\Delta\alpha_s(m_W^2)$
Unit CKM	$0.1208 \pm 0.0004_{\text{param.}} \pm 0.0275_{\text{exp.}}$	$\pm 23\%$

\Rightarrow Improved $\Gamma_W(\text{hadronic})$ at LHC allows to extract α_s with $\sim 23\%$ uncertainty.



- Uncertainties at FCC-ee (5×10^8 W's at $\sqrt{s} = m_W$):
 - Statistical: $\sim 0.005\%$ (0.4% at LEP with 8×10^4 W's)

Final result: $BR_W(\text{hadronic}) \sim 0.67410 \pm 0.00003$

	$\alpha_s(m_W^2)$	$\Delta\alpha_s(m_W^2)$
Unit CKM	$0.1208 \pm 0.0004_{\text{exp.}}$	$\pm 0.3\%$

\Rightarrow FCC-ee would provides us a value for α_s with a relative uncertainty of $\sim 0.3\%$.



Summary



⇒ We computed $\Gamma_W(\text{hadronic})$ and $BR_W(\text{hadronic})$ using state-of-the-art $N^3\text{LO}/\text{NNLO}$ calculations, removing some of the previously applied approximations:

$$\Gamma_W(\text{hadronic}) = (1411.546 \pm 0.772) \text{ MeV},$$

$$BR_W(\text{hadronic}) = 0.67422 \pm 0.00003.$$



⇒ Current experimental $\Gamma_W(\text{hadronic})$, $BR_W(\text{hadronic})$ and $|V_{CS}|$ uncertainties preclude precise extraction of α_S .

⇒ Improvements at LHC, and in particular FCC-ee, will allow one to incorporate $\Gamma_W(\text{hadronic})$ and $BR_W(\text{hadronic})$ into the PDG α_S .

	$\alpha_S(m_W^2)$	$\alpha_S(m_Z^2)$	$\Delta\alpha_S(m_W^2)$
Today (Unit CKM)	$0.1189 \pm 0.0004_{\text{param.}} \pm 0.0433_{\text{exp.}}$	$0.1167 \pm 0.0004_{\text{param.}} \pm 0.0430_{\text{exp.}}$	$\pm 37\%$
LHC (Unit CKM)	$0.1208 \pm 0.0004_{\text{param.}} \pm 0.0271_{\text{exp.}}$	$0.1185 \pm 0.0004_{\text{param.}} \pm 0.0260_{\text{exp.}}$	$\pm 23\%$
FCC-ee	$0.1208 \pm 0.0004_{\text{exp.}}$	$0.1185 \pm 0.0004_{\text{exp.}}$	$\pm 0.3\%$

⇒ FCC-ee will allow us to measure α_S with $\sim 0.3\%$ uncertainty.

Thank you. Any questions?

Backup slides

Experimental Hadronic Width Measurement

- Experimental hadronic $\Gamma_W(\text{hadronic})$ from $\Gamma_W(\text{total})$ via $\text{BR}_W(\text{hadronic})^2$:

$$\begin{aligned}\Gamma_W(\text{hadronic}) &= \Gamma_W(\text{total}) \times \text{BR}_W(\text{hadronic}) \\ &= (2085 \pm 42) \times (0.6741 \pm 0.0027) \text{ MeV} \\ &= (1405 \pm 29) \text{ MeV} \text{ (i.e. 2\% uncertainty)}\end{aligned}$$

- $\Gamma_W(\text{total})$ in $e^+e^- \rightarrow W^+W^- \rightarrow 4q, 2q + \ell\nu$

Where: LEP ($\sqrt{s} = 161 - 209 \text{ GeV}$) & FCC-ee ($\sqrt{s} = 161, 240, 350 \text{ GeV}$)

How: **Maximum-likelihood fit of m_W Breit-Wigner** with $\Gamma_W(\text{total})$ as free parameter.

- $\Gamma_W(\text{total})$ in $p\bar{p}, pp \rightarrow W + X$, with $W \rightarrow e\nu, \mu\nu$


Where: Tevatron ($\sqrt{s} = 1.8, 1.96 \text{ TeV}$) & LHC ($\sqrt{s} = 7, 8, 13, 14 \text{ TeV}$)

How: **Maximum likelihood fit of high $m_T(W)$ tail** with $\Gamma_W(\text{total})$ as free parameter (and via $\sigma(W)/\sigma(Z)$ ratios).

²0.4 % uncertainty measured in e^+e^-

LEP-2 W-boson Width Measurement [arXiv:1302.3415]

- $e^+e^- \rightarrow W^+W^- \rightarrow 4q, 2q + \ell\nu$ at $\sqrt{s} = 161 - 209$ GeV
- Statistics:
N(W's in all channels / experiments) ~ 40.000 pairs
- Binned likelihood fit to m_W Breit-Wigner with Γ_W (total) as free parameter:



Source	Systematic Uncertainty in MeV			
	on m_W			on Γ_W
	$q\bar{q}\ell\nu_\ell$	$q\bar{q}q\bar{q}$	Combined	
ISR/FSR	8	5	7	6
Hadronisation	13	19	14	40
Detector effects	10	8	9	23
LEP energy	9	9	9	5
Colour reconnection	–	35	8	27
Bose-Einstein Correlations	–	7	2	3
Other	3	10	3	12
Total systematic	21	44	22	55
Statistical	30	40	25	63
Statistical in absence of systematics	30	31	22	48
Total	36	59	34	83

- Final result: Γ_W (total) = $(2495 \pm 63_{\text{stat.}} \pm 55_{\text{syst.}})$ MeV

CDF W-boson Width Measurement [PRL 100, 071801 (2008)]

- $p\bar{p} \rightarrow W + X$, with $W \rightarrow e\nu, \mu\nu$ at $\sqrt{s} = 1.96$ TeV ($L_{int.} \sim 350$ pb $^{-1}$)
- Statistics:
N(W's)=3.436+2.619 with $90 < M_T < 200$ GeV
- Binned likelihood fit to ($90 < M_T < 200$ GeV) spectra with Γ_W (total) as free parameter:

$$M_T = \sqrt{2(p_T^\ell p_T^\nu - \vec{p}_T^\ell \cdot \vec{p}_T^\nu)}$$




TABLE I. The sources of uncertainty (in MeV) on Γ_W for the $W \rightarrow e\nu$ and $W \rightarrow \mu\nu$ measurements. If there is a correlated source of error between the two measurements its contribution to each measurement is listed in the third column, labeled C.

Source	$\Delta\Gamma_W^{e\nu}$	$\Delta\Gamma_W^{\mu\nu}$	C
Statistics	60	67	
Lepton E or p scale	21	17	12
Lepton E or p resolution	31	26	
Electron energy loss simulation	13		
Recoil model	54	49	
p_T^W	7	7	7
Backgrounds	32	33	
PDFs	20	20	20
M_W	9	9	9
EW radiative corrections	10	6	6
Lepton ID/acceptance	10	7	
Total systematic	79	71	27
Total (statistic + systematic)	99	98	27

- Final result: Γ_W (total) = $(2032 \pm 45_{\text{stat.}} \pm 57_{\text{sys.}})$ MeV

Tevatron W-boson Width Combined [arXiv: 1003.2826]


- CDF + D0 combined (BLUE method) $p\bar{p} \rightarrow W + X$, with $W \rightarrow e\nu, \mu\nu$ at 1.8 and 1.96 TeV.
- Improved likelihood fits to M_T spectra with updated underlying parameters:

	Run-I			Run-II	
	CDF-Ia	CDF-Ib	D0-Ib	CDF	D0
Γ_W (published)	2,110	2,042.5	2,231	2,032	2,028.3
Total uncertainty (published)	329	138.3	172.8	72.4	72
M_W used in publication	80,140	80,400	80,436	80,403	80,419
Correction to Γ_W from M_W	-78	0.3	11.1	1.2	6.0
Γ_W (corrected)	2,032	2,042.8	2,242.1	2,033.2	2,034.3
Total uncertainty (corrected)	329.3	138.3	172.4	72.4	71.9
Uncorrelated uncertainty (corrected)	327.6	136.8	167.4	68.7	68.5
PDF uncertainty (published)	0	15	39	20	20
PDF uncertainty (this analysis)	15	15	39	20	20
EWK RC uncertainty	28	10	10	6	7
M_W uncertainty (published)	0	10	15	9	5
M_W uncertainty (this analysis)	7	7	7	7	7
M_W extrapolation	26	0	4	0	2

- Final result: $\Gamma_W(\text{total}) = (2046 \pm 49) \text{ MeV} \begin{matrix} \pm 39 \text{ MeV (stat.)} \\ \pm (20+7.4+7.4) \text{ MeV (PDF + } m_W\text{+EW. corr.)} \end{matrix}$

W-boson Width: Tevatron + LEP Combined [PDG]

- World average of all LEP + Tevatron measurements:



Final result: $\Gamma_W(\text{total}) = (2.085 \pm 0.042)$ GeV (i.e. 2% uncertainty)

Note: $\Gamma_Z(\text{total}) = (2.4952 \pm 0.0023)$ GeV (i.e. 0.1 % uncertainty)

List of References



A. Denner, B. Kniehl, J. Kühn, K. Chetyrkin, ...



Dominik Kara *Corrections of Order $\alpha\alpha_s$ to W boson decays*, Nucl. Phys. B 877, 3 (2013)



G. C. Cho, K. Hagiwara, Y. Matsumoto and D. Nomura, *The MSSM confronts the precision electroweak data and the muon $g - 2$* , JHEP 111 (2011), 068



K.A. Olive et al. (Particle Data Group), Chin. Phys. C, 38, 090001 (2014)