

Double Beta Decay of Zr96 using NEMO-3 and Calorimeter R&D for SuperNEMO

Wednesday, April 8, 2009 12:15 PM (15 minutes)

Using 911 days of data from NEMO-3, a world best $2\nu\text{BB}$ decay half-life of Zr96 has been measured to be $[2.36 \pm 0.17(\text{stat}) + 0.17 - 0.14(\text{syst})] \times 10^{19}$ yr. The obtained limit on the $0\nu\text{BB}$ decay half-life at the 90% confidence level is 8.5×10^{21} yr which leads to the limit on the effective Majorana neutrino mass $< 7.4 - 20.3$ eV, using the RQRPA and pnQRPA nuclear models. SuperNEMO is a next-generation double beta decay experiment based on the successful tracking plus calorimetry design approach of the NEMO-3. SuperNEMO can study a range of isotopes, the baseline isotopes are Se82 and possibly Nd150. The total isotope mass will be 100-200 kg. A sensitivity to a $0\nu\text{BB}$ half-life greater than 10^{26} years can be reached which gives access to Majorana neutrino masses of 50-100 meV. One of the main challenges of the SuperNEMO R&D is the development of the calorimeter with an energy resolution of 4% FWHM at 3 MeV ($Q(\text{bb})$ value of Se82). This unprecedented milestone has been achieved using low density plastic scintillator coupled to high quantum efficiency photomultiplier tubes.

Primary author: KAUER, Matthew (University College London)

Presenter: KAUER, Matthew (University College London)

Session Classification: Parallel Session 3 C - Neutrinos and Double Beta Decay

Track Classification: Neutrinos