



Contribution ID: 82

Type: **Contributed Talk**

## Heavy-flavour production in pp collisions and correlations in pp and p-Pb collisions with ALICE at the LHC

*Tuesday 28 June 2016 16:00 (20 minutes)*

Heavy quarks (charm and beauty) are produced through hard parton scatterings in the initial stages of hadronic collisions. The study of their production in pp collisions constitutes a powerful tool to test perturbative QCD calculations at the LHC energies, where a region of phase space at low Bjorken- $x$  can be accessed. Moreover, these measurements provide a reference for the study of nuclear matter effects on heavy quarks in Pb-Pb collisions, where a Quark-Gluon Plasma (QGP) is produced.

Further insight into the mechanisms through which heavy quarks are produced and fragment into hadrons can be obtained by studying the angular correlations between heavy-flavour particles and charged hadrons produced in pp collisions. The comparison of the angular correlation distributions with those measured in p-Pb collisions can also help to investigate possible modifications of the heavy-quark production and hadronization induced by the presence of cold nuclear matter effects in the latter collision system, besides constituting a reference for the interpretation of Pb-Pb measurements. A comparison of correlation measurements with theoretical predictions can also allow us to put constraints on the models describing heavy-quark production and hadronization.

Thanks to its excellent tracking and particle identification performance, the ALICE detector is capable of measuring D mesons at central rapidity via their hadronic decay channels down to very low transverse momentum, as well as leptons from heavy-flavour hadron decays at central and forward rapidity. We will present a review of ALICE measurements of heavy-flavour production in pp collisions at  $\sqrt{s} = 7$  TeV and at  $\sqrt{s} = 2.76$  TeV. In particular, we will show an extension of the prompt  $D^0$   $p_T$ -differential cross section down to zero transverse momentum, which allows us to measure the  $p_T$ -integrated charm production cross section at central rapidity, in pp collisions at  $\sqrt{s} = 7$  TeV. In addition, we will discuss measurements of azimuthal correlations of prompt  $D^0$ ,  $D^+$  and  $D^{*+}$  mesons with charged hadrons in pp collisions at  $\sqrt{s} = 7$  TeV and p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV and compare the results with expectations from models.

### On behalf of collaboration:

ALICE

**Primary author:** COLAMARIA, Fabio Filippo (Universita e INFN, Bari (IT))

**Presenter:** COLAMARIA, Fabio Filippo (Universita e INFN, Bari (IT))

**Session Classification:** Heavy Quark Production