Strangeness in Quark Matter 2016



Contribution ID: 105

Type: Contributed Talk

Strange partial pressures from Lattice QCD

Thursday, 30 June 2016 10:00 (20 minutes)

The precision reached by recent lattice QCD results allows for the first time to investigate whether the measured hadronic spectrum is missing some additional strange states, which are predicted by the Quark Model [1,2] but have not yet been detected. This can be done by comparing some sensitive thermodynamic observables from lattice QCD to the predictions of the Hadron Resonance Gas model (with the inclusion of decays [3]). We propose a set of specific observables [4], defined as linear combinations of conserved charge fluctuations, which allow to investigate this issue for baryons containing one or more strange quarks separately. Applications of these observables to isolate the multiplicity fluctuations of kaons from lattice QCD, and their comparison [5] with the experimental results, are also discussed.

[1]S. Capstick and N. Isgur, Phys. Rev. D 34, 2809 (1986)

[2] D.Ebert, R.N.Faustov and V.O.Galkin, Phys. Rev. D 79, 114029 (2009)

[3] P.Alba, R.Bellwied, M.Bluhm, V.Mantovani Sarti, M.Nahrgang and C.Ratti, Phys. Rev. C 92, no. 6, 064910 (2015)

[4] A.Bazavov et al., Phys. Rev. Lett. 111, 082301 (2013)

[5] R.Bellwied, S.Borsanyi, Z.Fodor, S.D.Katz and C.Ratti, Phys. Rev. Lett. 111, 202302 (2013)

On behalf of collaboration:

None

Primary author: NORONHA-HOSTLER, Jacquelyn (University of Houston)

Co-authors: PASZTOR, Attila (Wuppertal University); RATTI, Claudia (University of Houston); ALBA, Paolo Giuseppe; BELLWIED, Rene (University of Houston (US)); KATZ, Sandor (Eotvos University); BORSANYI, Sz-abolcs (University of Wuppertal); MANTOVANI SARTI, Valentina (INFN Torino); FODOR, Zoltan (BUW)

Presenter: NORONHA-HOSTLER, Jacquelyn (University of Houston)

Session Classification: Strangeness in Astrophysics