Strangeness in Quark Matter 2016



Contribution ID: 73

Type: Contributed Talk

Measurement of higher harmonic flow coefficients of identified hadrons in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV

Thursday, 30 June 2016 09:00 (20 minutes)

The latest ALICE results on the centrality and transverse momentum dependence of v_2 , v_3 , v_4 and v_5 for π^{\pm} , \mathbf{K}^{\pm} and $\mathbf{p}+\overline{\mathbf{p}}$ in Pb-Pb collisions at $\sqrt{s_{\mathrm{NN}}} = 2.76$ TeV are presented. The values of these flow coefficients exhibit a clear mass ordering for $p_{\mathrm{T}} < 3$ GeV/c for all harmonics. For transverse momentum values larger than about $3 \,\mathrm{GeV}/c$, mesons exhibit distinctive flow values compared to baryons, suggesting that coalescence might be the relevant particle production mechanism in this region. The experimental data for $p_{\mathrm{T}} < 3 \,\mathrm{GeV}/c$ are described fairly well by the hydrodynamical model iEBE-VISHNU, which models the hydrodynamical expansion of the fireball using a value of $\eta/s = 0.08$, coupled to a hadronic cascade model (UrQMD). Finally, we find that predictions from A Multi-Phase Transition Model (AMPT) indicate the late hadronic rescattering stage contributes significantly to the mass splitting of azimuthal flow (n=2-4). These predictions are compared to our data.

On behalf of collaboration:

ALICE

Primary author: MOHAMMADI, Naghmeh (Nikhef National institute for subatomic physics (NL))Presenter: MOHAMMADI, Naghmeh (Nikhef National institute for subatomic physics (NL))Session Classification: Flow