



# Results from the OPAL Experiment

Richard Hemingway

IPP/Carleton University

On behalf of the OPAL Collaboration

LEP Fest, 10th October 2000

- Introductory Remarks

Luminosity .. Y2K Data

- Physics Results

PRELIMINARY LEP2, including  $\sqrt{s} > 202 \text{ GeV}$

- Standard Model cross-sections and couplings
- Indirect limits on new physics
- Search for SM and MSSM Higgs
- Searches for new particles

- Where do we go from here?



## Apologies .. Consult our Web site

30 minutes is too short to give adequate recognition to the intensive work over the past few weeks (months) (years) ..

Please consult our public web site

'The OPAL experiment at LEP'

<http://opal.web.cern.ch/Opal/PPwelcome.html>

where you will find all OPAL physics results and, in particular, 3 Collective Physics Notes prepared for this LEP Fest

- Measurement of Standard Model Processes in e+e- Collisions at  $\sqrt{s} > 202$  GeV
- Updated Results of Higgs Boson Searches in e+e- Collisions at the Highest LEP Energies
- New Particle Searches in e+e- Collisions at  $\sqrt{s} = 200-209$  GeV



# Special People .. Many Thanks

- OPAL Spokesmen

- Aldo Michelini, early beginnings - 1993

- Design, construction, installation, exploitation LEP1

- Rolf Heuer, 1994 - mid 1998

- LEP1 to LEP2 transition, precision physics

- Dave Plane, mid 1998 - present

- LEP2 highest energies/luminosities

- OPAL Secretary

- Mette Stuwe, early beginnings - present




# The OPAL Collaboration .. past/present

- **CANADA**  
Alberta, Carleton, CRPP/NRC, Montreal, UBC, Victoria
- **FRANCE**  
Saclay
- **GERMANY**  
Aachen, Bonn, Freiburg, Hamburg/DESY, Heidelberg, LMU-Munich, MPI-Munich
- **HUNGARY**  
Budapest, Debrecen
- **ISRAEL**  
Technion, Tel Aviv, Weizmann
- **ITALY**  
Bologna
- **JAPAN**  
ICEPP-Tokyo/Kobe
- **CERN**
- **UK**  
Birmingham, Brunel, Cambridge, Manchester, QMW, RAL, UCL/Birkbeck
- **USA**  
Chicago, Duke, Indiana, Maryland, Oregon, Riverside, Yale



# OPAL Luminosity Summary

- Special thanks to the LEP Division and all the technical staff associated with the LEP program. Each year, 1989-2000, has been great!




Today: Total luminosity recorded in Y2K almost 200 pb<sup>-1</sup>

OPAL data taking efficiency in Y2K = 92%



# OPAL Data Quality

- Calibration procedures for OPAL have maintained steady and robust resolutions throughout the LEP2 period








# A Bhabha event at 209 GeV

- 10 August 2000, LEP sets energy record at 209 GeV
- Run lasted less than 2 minutes. Pity!  $80 \text{ nb}^{-1}$


```
Run:event 14880: 285      Ctrk(N= 2 Sump=162.1) Ecal(N= 14 SumE=216.5)
Ebeam 104.50 Vtx ( -.02, .04, .53) Hcal(N= 0 SumE= .0) Muon(N= 0)
```





# Standard Model Cross-sections

- from CERN Yellow Report, LEP2 Physics








# A Multihadron event at 208 GeV

- One person's signal is another person's background!

Run:event 14590: 1495      Ctrk(N= 52 Sump=325.9) Ecal(N= 71 SumE=112.2)  
 Ebeam 104.10 Vtx ( -.02, .03, .01) Hcal(N=30 SumE= 24.6) Muon(N= 1)




# Cross-section for hadrons

- separate full-energy annihilation events from  $Z^0$  radiative return

1478 non-radiative events at  $\sqrt{s} = 205$  GeV

1709 non-radiative events at  $\sqrt{s} = 207$  GeV




Curve is ZFITTER prediction

# Cross-section for mu-pairs

- separate full-energy annihilation events from  $Z^0$  radiative return

211 non-radiative events at  $\sqrt{s} = 205$  GeV

225 non-radiative events at  $\sqrt{s} = 207$  GeV




Curve is ZFITTER prediction



# Forward-backward Asymmetries

- Measure  $A_{fb}$  for lepton pairs  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$




Curve is BHWIDE ( $e^+e^-$ ), ZFITTER ( $\mu^+\mu^-$ ,  $\tau^+\tau^-$ ) prediction



# Fine Structure Constant

- Use non-radiative cross-sections and asymmetries
- ZFITTER, with all other pars fixed, gives  $\alpha_{em}(\sqrt{s})$


$$\alpha_{em}(\sqrt{s} = 190.6 \text{ GeV}) = 128.4^{+2.5}_{-2.3} \text{ (SM: 127.9)}$$





# QCD at 206 GeV

- All QCD observables well represented by PYTHIA, HERWIG
- No evidence for anomalous 4-jet production






# Strong Coupling Constant

- Fit distributions of 1-T,  $M_H$ , C,  $B_W$ ,  $B_T$ , and  $y_{23}^D$  to NLLA  $\mathcal{O}(\alpha_s^2)$  QCD calculations
- Results consistent with running of  $\alpha_s$

$$\alpha_s(\sqrt{s} = 205.9 \text{ GeV}) = 0.107 \pm 0.002 \pm 0.004$$

## Summary of OPAL measurements of $\alpha_s(Q)$






# Cross-section for photon-pairs

- A pure QED process at tree level

At  $\sqrt{s}=205$  GeV events observed/expected = 467/463

At  $\sqrt{s}=207$  GeV events observed/expected = 534/549








# Cross-section for WW-pairs

- Isolate all 3 decay channels  $WW \rightarrow q\bar{q}q\bar{q}, q\bar{q}l\bar{\nu}_e, l\bar{\nu}_el\bar{\nu}_e$
- Cross-sections assume SM W decay fractions

At  $\sqrt{s}=205$  GeV obtain 651, 545, 125 events resp.


At  $\sqrt{s}=207$  GeV obtain 887, 708, 162 events resp.



SM prediction via RACOONWW and YFSWW



# A $WW \rightarrow q\bar{q}q\bar{q}$ event at 208 GeV




# Cross-section for ZZ-pairs

- Isolate decay channels  $llll, ll\nu\nu, qqll, qq\nu\nu, qqqq$
- Cross-sections assume SM Z decay fractions

At  $\sqrt{s}=205$  GeV obtain 77 candidates, expected SM bkgd = 37

At  $\sqrt{s}=207$  GeV obtain 85 candidates, expected SM bkgd = 45



No evidence for non-zero neutral TGCs ( $ZZ\gamma, ZZZ$ )




# A ZZ event at 205 GeV

Run:event13394: 5924 Date:\*\*\*\*\* Time 105638 Ctrk(N= 25 Sump=174.8) Ecal(N= 36 SumE= 63.2) Hcal(N=17 SumE= 31.3)  
 Ebeam102.699 Evis 213.8 Emiss -8.4 Vtx ( -0.05, 0.04, -0.62) Muon(N= 2) Sec Vtx(N= 2) Fdet(N= 0 SumE= 0.0)  
 Bz=4.023 Bunchlet 1/1 Thrust=0.7139 Aplan=0.0045 Oblat=0.4498 Spher=0.1625



$M_{mm} \sim 93.1 \text{ GeV}$   $M_{qq} \sim 90.6 \text{ GeV}$



Centre of screen is ( -47.1442, -3.7475, 0.0000 )


200. cm.      5 10 20 50 GeV



# Charged Current TGCs

- Combine WW cross-section and angular distributions with single-W cross-section

## OPAL Preliminary



No anomalous behaviour: SM values look OK.

Obtain limits on CP-violating TGCs via spin density matrix.

No evidence for anomalous QGCs, eg  $WW\gamma\gamma$ .


# Precision $M_W$ Measurement

- With  $480\text{pb}^{-1}$  (prior to Y2K)

$$m_W = 80.485 \pm 0.052(\text{stat}) \pm 0.039(\text{sys}) \text{ GeV}$$

- With final statistics, expect

$$m_W = 80.xxx \pm 0.040(\text{stat}) \pm 0.025(\text{sys}) \text{ GeV}$$



Data shows no FSI (BEC,CR),  $M(qqqq) = M(qq\nu\nu)$ .

Emphasise need for good LEP energy determination.

Estimate all-LEP error 30-35 MeV (SM indirect = 26 MeV).



## Non-SM Physics ...indirect limits


$f\bar{f}, \gamma\gamma, ZZ, \dots$  cross-sections and couplings in agreement with SM. They provide limits on possible new physics (generally model dependent).

- **4-fermion Contact Interactions**  
Mass limits 8-15 TeV with  $g^2/4\pi = 1$
- **Sneutrino exchange (RPV) in s-channel**  
 $\lambda_{131}, \lambda_{121}$  coupling limits within 100-300 GeV range
- **$Z'$  exchange in s-channel**  
Exclusion limits in 400-750 GeV range
- **QED  $\Lambda$  cut-off parameters in  $e^+e^- \rightarrow \gamma\gamma$**   
Both  $\Lambda_+, \Lambda_-$  above 330 GeV
- **Excited electron in t-channel  $\gamma\gamma$**   
Mass limit  $\sim 300$  GeV assuming  $e^*e\gamma = ee\gamma$  coupling
- **Low scale quantum gravity in  $\mu^+\mu^-, \tau^+\tau^-, \gamma\gamma, ZZ$**   
Mass limits 830-900 GeV

**General conclusion: New Physics is far beyond EW scale**




# Contact Interactions








# Limits on possible $Z'$




# Gravity in Extra Dimensions



95% CL lower limit on  $M_S$  is

0.83 TeV with  $\lambda = -1$  and 0.90 TeV with  $\lambda = +1$

# Standard Model Higgs Decays



4-jet channel, data=18, bkgd=19.1+-2.4

Missing E ch, data=26, bkgd=28.1+-2.9

Tau channel, data=4, bkgd=2.7+-0.4





e/mu channel, data=9, bkgd=5.9+-1.0

TOTAL, data=57, bkgd=55.7+-5.8




# Higgs: Individual Channel Mass Distributions

## OPAL Preliminary






# Higgs: Mass Distribution, all Y2K data






# Higgs: Mass Distribution, only 207 GeV data






# Higgs: $1-CL_b$ : background-only hypothesis





# SM Higgs Mass .. 95% CL lower limit



$m_H > 107.9$  GeV at 95% CL





# More Higgs Searches

- SUSY Higgs

- $e^+e^- \rightarrow h^0 Z^0$ : as HZ search
- $e^+e^- \rightarrow A^0 h^0 \rightarrow b\bar{b}b\bar{b}$ : Data=7, Bkgd=8.4+-1.3
- $e^+e^- \rightarrow A^0 h^0 \rightarrow \tau^+ \tau^- b\bar{b}$ : Data=5, Bkgd=3.4+-0.5

For  $\tan(\beta) > 1.2$ , determine  
 $m_h > 80.7 \text{ GeV}$ ,  $m_A > 83.2 \text{ GeV}$ .

- $e^+e^- \rightarrow H^+ H^- \rightarrow \tau^+ \nu_\tau q\bar{q}'$ : Data=160, Bkgd=168
- $e^+e^- \rightarrow H^+ H^- \rightarrow q\bar{q}' q\bar{q}'$ : Data=279, Bkgd=302

$m_{H^\pm} > 72.3 \text{ GeV}$  assuming  $\text{Br}(\tau\nu + q\bar{q}) = 1$

- Special Higgs Decays

- Invisible  $h^0$ , eg.  $\rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0$ : Data=35, Bkgd=53


$m_{h^0} > 107.2 \text{ GeV}$  assuming SM prod rate

- Fermiophobic  $h^0 \rightarrow \gamma\gamma$ : Data=16, Bkgd=19

$m_{h^0} > 104.6 \text{ GeV}$  assuming SM prod rate

# Limits in CMSSM parameter space

$m_h$ -max: A specific benchmark MSSM scan which provides the most conservative range of excluded  $\tan(\beta)$  values.



Exclusion:  $\tan(\beta)$  region 0.8-2.0



## Search Channel List


$e^+e^- \rightarrow hZ$	$h \rightarrow b\bar{b}$	$b\bar{b}q\bar{q}, b\bar{b}\nu\bar{\nu}, b\bar{b}l^+l^-$
	$h \rightarrow \gamma\gamma$	$(q\bar{q}, l^+l^-, \nu\bar{\nu}) + \cancel{E}$
$e^+e^- \rightarrow hA$	$h \rightarrow \tilde{\chi}^0\tilde{\chi}^0$	$q\bar{q}, l^+l^- + \cancel{E}$
	$h, A \rightarrow b\bar{b}, \tau\tau$	$b\bar{b}b\bar{b}, b\bar{b}\tau^+\tau^-$
	$h \rightarrow AA$	$b\bar{b}b\bar{b}b\bar{b}$
$e^+e^- \rightarrow H^+H^-$	$H^+ \rightarrow q\bar{q}, \tau\nu$	$q\bar{q}q\bar{q}, q\bar{q}\tau\nu, \tau\nu\tau\nu$
$e^+e^- \rightarrow \tilde{\chi}^+\tilde{\chi}^-$	$\tilde{\chi}^- \rightarrow W^*\tilde{\chi}^0$	$\text{jets } (+l^\pm), l^+l^- + \cancel{E}$
	(In)Direct RPV	$\text{jets}, l^\pm, \nu$
	$(\tilde{\chi}^0 \rightarrow \gamma\tilde{G})$	$\text{jets}, l^+l^-, \gamma\gamma + \cancel{E}$
$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_1^0$	$\tilde{\chi}_2^0 \rightarrow Z^0\tilde{\chi}_1^0$	$2 \text{ jets} + \cancel{E}$
	$\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$	$\gamma + \cancel{E}$
$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0$	$\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$	$\gamma\gamma + \cancel{E}$
	$\tilde{\chi}_1^0$ Lifetime	non-pointing $\gamma$
$e^+e^- \rightarrow \tilde{l}^+\tilde{l}^-$	RPV Decays	$\text{jets}, l^\pm, \nu$
	$\tilde{l}^- \rightarrow l^-\tilde{\chi}_1^0$	$l^+l^- + \cancel{E}$
	$(\tilde{\chi}^0 \rightarrow \gamma\tilde{G})$	$l^+l^-\gamma\gamma + \cancel{E}$
	(In)Direct RPV	$2,4,6 \times l^\pm + \cancel{E}$
	$\tilde{l}^\pm$ Lifetime	Kinked Tracks
		Stable, Charged
$e^+e^- \rightarrow \tilde{\nu}\tilde{\nu}$	(In)Direct RPV	$l^+l^-l^+l^-$
		$\text{jets} + \cancel{E}$
$e^+e^- \rightarrow \tilde{t}_1\tilde{t}_1$	$\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$	$2 \text{ jets} + \cancel{E}$
	$\tilde{t}_1 \rightarrow bl^+\tilde{\nu}$	$2 \text{ jets} + l^+l^- + \cancel{E}$
	(In)Direct RPV	$l^+ql^-q$
$e^+e^- \rightarrow N\bar{N}$	$N \rightarrow lW$	$\text{jets} + l^\pm$
$e^+e^- \rightarrow L^+L^-$	$L^+ \rightarrow \nu W$	$\text{jets}, l^\pm + \cancel{E}$
$e^+e^- \rightarrow l^{*+}l^{(*)-}$	$l^{*+} \rightarrow l^+\gamma$	$l^+l^-\gamma(\gamma)$
$e^+e^- \rightarrow \nu^*\bar{\nu}^*$	$\nu^* \rightarrow \nu\gamma$	$\gamma(\gamma) + \cancel{E}$
$e^+e^- \rightarrow l^*l, \nu^*\nu$	$l^* \rightarrow lZ$	$\text{jets}, l^\pm, \nu$

# Single Photon Recoil Mass

Sensitive to Gravity in Extra Dimensions, GMSB scenarios, eg.

$\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$ , and MSSM  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \gamma$ , and excited neutrinos, eg.  
 $\nu^* \rightarrow \nu \gamma$ .

Standard Model process is  $e^+e^- \rightarrow \nu\bar{\nu}\gamma(\gamma)$ .





# Non-SM Physics ...direct searches

In general a null search provides limits on cross-sections, couplings, masses, ...

- **Leptoquarks**

$$e^+e^- \rightarrow L_q \bar{L}_q, L_q \rightarrow lq \text{ Data}=54, \text{ Bkgd}=55$$

- **Heavy Leptons**

$$e^+e^- \rightarrow N \bar{N}, N \rightarrow lW \text{ Data}=63, \text{ Bkgd}=51$$

- **Excited Leptons**

$$e^+e^- \rightarrow l^{*+} l^{*-}, l^* \rightarrow l\gamma \text{ Data}=6, \text{ Bkgd}=4$$

$$e^+e^- \rightarrow l^{*+} l^-, l^* \rightarrow l\gamma \text{ Data}=642, \text{ Bkgd}=691$$

- **Stable, long-lived, massive particles**

$$\text{Sensitive to } Q/e = \pm 1, \pm 2/3 \text{ Data}=0, \text{ Bkgd}=1$$

- **Single top via FCNC**

$$e^+e^- \rightarrow t\bar{c}(\bar{u}) \text{ Data}=21, \text{ Bkgd}=23$$

$$M(l^*) > 103 \text{ GeV}$$

$$M(\tilde{\ell}) > 97 \text{ GeV}, M(\tilde{\chi}^\pm) > 101 \text{ GeV}, \text{ for long-lived}$$

$$\sigma_{top} < 0.36 \text{ pb, assuming } \text{Br}(t \rightarrow bW) = 1$$



## Non-SM Physics ...direct SUSY searches

MSSM searches of two types are conducted

(A) MSUGRA, with/without RPC, where LSP =  $\tilde{\chi}_1^0$  (stable),  
which leads to topologies with jets, leptons, AND Missing Energy

(B) GMSB, where LSP =  $\tilde{G}$  [ $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ ,  $\tilde{\ell} \rightarrow l\tilde{G}$ ], which give  
MORE leptons and photons

- **Scalar Leptons**

$$e^+e^- \rightarrow \tilde{\ell}^+\tilde{\ell}^-, \tilde{\ell} \rightarrow l\tilde{\chi}_1^0$$

- **Scalar top/bottom quarks**

$$e^+e^- \rightarrow \tilde{t}\tilde{t}^-, \tilde{t} \rightarrow c\tilde{\chi}_1^0, b\tilde{\chi}_1^\pm$$

$$e^+e^- \rightarrow \tilde{b}\tilde{b}^-, \tilde{b} \rightarrow b\tilde{\chi}_1^0$$

- **Charginos**

$$e^+e^- \rightarrow \tilde{\chi}^\pm\tilde{\chi}^\mp, \tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 W^\pm$$

- **Neutralinos**


$$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_1^0, \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 Z^0$$

No compelling evidence for data in excess of SM bkgd

Cross-section limits are determined

Calculate exclusion regions in MSSM parameter space


# MSSM exclusion limit for stop





# MSSM exclusion limit for gauginos


## OPAL Preliminary







# MSSM exclusion limit for neutralino



**Absolute lower limit on lightest neutralino**

$$m_{\tilde{\chi}_1^0} > 39.0 \text{ GeV for } m_0 > 500 \text{ GeV}$$

$$m_{\tilde{\chi}_1^0} > 36.0 \text{ GeV for any } m_0$$



## Summary

- OPAL data taking in 2000 very successful  
 $\int \mathcal{L} dt$  almost  $200 \text{ pb}^{-1}$
- Many ongoing physics analyses, both LEP1 and LEP2
- All results from LEP data in good agreement with SM predictions.
- We look forward to several years of continuing physics analysis and  
.....perhaps, in some forgotten corner, unexpected new physics